

COUNTWAY LIBRARY



HC 4YRT .

COSMETICS

THEODOR KOLLER

BOSTON
MEDICAL LIBRARY
& THE FENWAY



57- net

3/6

(2)

C

COSMETICS



Digitized by the Internet Archive
in 2024

COSMETICS

A HANDBOOK

OF THE MANUFACTURE, EMPLOYMENT, AND
TESTING OF ALL COSMETIC MATERIALS AND
COSMETIC SPECIALTIES

BY

DR. THEODOR KOLLER

TRANSLATED FROM THE GERMAN

BY

CHAS. SALTER

LONDON.

SCOTT, GREENWOOD & CO.

19 LUDGATE HILL, E.C.

1902

[The sole right of translation into English rests with the above firm]



27.C.33.

PREFACE.

THE cosmetic industry is mainly indebted to the researches of chemists, who have first pointed out the right paths to follow, then introduced new and valuable substances to employ in the preservation of the hair and teeth, and also obviated certain injurious influences by the aid of antiseptics. Furthermore, dangerous substances have been replaced by innocuous substitutes; and, in a word, the technology of cosmetics has been undoubtedly elevated to the position of a branch of natural science.

The manufacture of cosmetics is still, in many instances, suffering from too close an adherence to ancient tradition, whereby entirely useless substances are accredited with healing powers; and even—which is far more serious—directly poisonous and injurious materials are employed, the use of which might entail great danger to health.

In many quarters too little importance is still attached to the consideration of the first question that should arise in connection with any cosmetic, namely, whether the same contains any ingredients capable of acting beneficially in the cases for which it is designed, and without leading to injurious results on prolonged employment.

Now, it is only when this primary question has been put, and satisfactorily answered, that the rational stage of cosmetic technology is reached; and to answer this question it is

necessary to have an accurate knowledge of the materials under consideration. This forms the basis of the present work, which is an endeavour to afford guidance and advice to the maker of cosmetic preparations; and to point out the best as well as the simplest methods of procedure. The fundamental principles to be followed in making up the various preparations are fully explained. Where the habits and whims of the public demand remedies where none exist, the question of how to satisfy this demand has received consideration, without, however, omitting to call attention to the inadequacy of the preparations themselves. All injurious ingredients have been strongly banned.

It is highly desirable that the mystery of "secret preparations" should be abandoned in cosmetics, and all endeavours concentrated on the attainment of perfection.

Such a course would soon gain the confidence of the public, and would be of the greatest advantage to the cosmetic industry itself, which would then take up a position beside the sister industry of perfumery; and, within its proper limits, be just as useful as other branches of technology.

That the present work may incite and conduce to the establishment of cosmetic technology on a rational basis, and may prove useful to all engaged in this really profitable branch, is the sincere wish of

THE AUTHOR.

CONTENTS.



CHAP.	PAGE
I. PURPOSES AND USES OF COSMETICS, AND INGREDIENTS USED IN THEIR PREPARATION	1
II. PREPARATION OF PERFUMES	11
III. CHEMICAL AND ANIMAL PRODUCTS USED IN THE MANU- FACTURE OF COSMETICS	19
IV. OILS AND FATS USED IN THE PREPARATION OF COSMETICS	54
V. GENERAL COSMETIC PREPARATIONS	85
VI. MOUTH WASHES AND TOOTH PASTES	101
VII. HAIR DYES, HAIR RESTORERS, AND DEPILATORIES	116
VIII. COSMETIC ADJUNCTS AND SPECIALTIES	142
IX. ANTISEPTIC WASHES AND SOAPS, TOILET AND HYGIENIC SOAPS	179
X. COSMETIC SECRET PREPARATIONS FOR THE SKIN, COM- PLEXION, TEETH, MOUTH, ETC.	208
XI. TESTING AND EXAMINING THE MATERIALS EMPLOYED IN THE MANUFACTURE OF COSMETICS	232
INDEX	243



COSMETICS

CHAPTER I.

PURPOSES AND USES OF COSMETICS, AND INGREDIENTS USED IN THEIR PREPARATION.

THE subject of cosmetics is closely allied to perfumery; but whilst the latter deals merely with the more external question of the dispersion of odour and aroma, and the masking of less pleasant smells, the domain of cosmetics is more of an internal character.

True, the two often join hands, as it were, and in many points it is difficult to maintain the line of demarcation between the two terms. This is particularly the case where it is a question of methods of preparation, application, and ingredients bordering on the domain of medicine. Here perfumery exhibits a certain independence, though not always to its advantage or repute; since in many instances preparations are made and sold which really belong almost entirely to the department of medicine, but contain substances either in themselves poisonous or else capable of producing injurious effects on the human organism by protracted use.

When rationally practised, the art of cosmetics should, however, be the means of preventing such encroachments, proving by critical analysis that such and such ingredients are worthless, and contain nothing capable of producing the effect

in view; or that the means proposed for a given purpose contain substances either prejudicial or directly inimical to health; and indicating means and ways for attaining the desired aim without the slightest detriment to health.

Thus the cosmetic art is primarily of hygienic importance. In this connection its task is to assist medicine in a friendly manner, and not to oppose the same. It should, by the means at its disposal, support and assist the functions of certain parts of the human body, protect them from injury, and remedy injuries already existing. This, however, is possible only when all the materials employed have been carefully tested in order to ascertain whether they are actually capable, by their nature and properties, of exerting a certain definite action, and seem to be unobjectionable from a hygienic point of view.

Furthermore, the art of cosmetics plays the part of an impartial adviser to the laity, whose confidence it enjoys in certain matters dealing with the care of the body; it opens and points the way for those engaged in the manufacture of cosmetic preparations; and, finally, constitutes a link in the chain of hygienic regulations, inasmuch as, in its capacity as a branch of hygienic science, it endeavours to remove or frustrate everything that is contrary or injurious to health.

The second task to be performed by cosmetics may be regarded as of an æsthetic character, namely, the beautifying of certain parts of the body. Even in this case, however, the main point of hygiene cannot be put out of sight; and this, indeed, is the crux which forms the stumbling-block of many thousands of ignorant or intentional deceivers, who, in their eagerness for profit, disregard the question whether what they offer the public may be detrimental to health.

Consequently, when the rational art of cosmetics deals also here with the analysis and testing of the means proposed, employed, or destined to be employed, for the beautifying of

certain parts of the body, it not only fulfils its hygienic mission, but is also at the same time explanatory and instructive.

Then, again, the third task of our subject is instructive as regards the elucidation and valuation of old and new preparations, and the proper means of producing suitable beautifying preparations.

The entire domain of cosmetics, in all three respects mentioned, is concerned with the rational care of the skin, the hair, the teeth, and the mouth. Moreover, it is not merely restricted to the indication of means capable of attaining the end in view, but also points and elucidates the manner in which those means can be prepared.

The rational care of the skin, hair, teeth, and mouth constitutes an important task, owing to the fact that the maintenance of these organs in a fit state to discharge their functions as they should be, is a fundamental condition for the health of the individual. At the same time, the æsthetic importance of the matter should be by no means despised, the progress made by chemistry having placed at disposal numerous means capable of furnishing successful results, without the slightest danger to health.

The art of cosmetics, however, will not attain its full repute and the complete appreciation of its importance, until it has been freed from the dross still adhering to it from times long past; until it takes its position as an independent technology, and never leaves out of sight—in its methods and the selection of materials—the hygienic mission with which it is charged. It will then also occupy the position of an assistant to medicine and a profitable patron of the art of perfumery.

Meanwhile it must not be forgotten that the need for cosmetic appliances has not arisen merely in our own times; on the contrary, this need extends backward to the earliest ages, and, moreover, is manifested among peoples whose standard of civilisation would not lead one to suppose such to be the

case. The cosmetic art of the present day, however, differs in the manner in which this need can be supplied, by the practical application of the progress of science to the production of cosmetic preparations, by rational processes and reliable ingredients, from that of older times, wherein chance, prejudice, ignorance, and even superstition were all convoked.

INGREDIENTS USED IN THE PREPARATION OF COSMETICS.

To a certain degree the preservation of scent and aroma is at least as important as the preservation of bodies occupying space and possessing form ; but, in consequence of the extraordinary volatility of the first named, is an exceedingly difficult task. Moreover, the external effect of scents and aromatics resides entirely in this volatility, which, besides, affords the sole means whereby their presence can be detected in association with other substances.

On the other hand, the retention of these readily vanishing products is one of the chief tasks of the cosmetic art, and the practitioner of this art must attempt to impart to his products a certain permanence and durability, despite the natural difficulties with which he is here confronted.

In the preservation of foodstuffs, wood, and metal, the conditions are, in some respects, simpler, it being for the most part merely a question of preventing decomposition, from internal or external sources, by the addition of suitable antiseptics, in the case of wood or foodstuffs ; in metals, external influences alone have to be guarded against ; whereas in preserving scent and aroma, the only point under consideration is the binding and fettering of the volatile substance.

The methods of preparing scents are various ; and the choice depends, not merely on the properties of the raw material from which the scent is to be recovered, nor on the yield and profit attainable, but more particularly on whether the scent can be

obtained in a pure state, and free from all extraneous admixtures capable of injuriously affecting its purity in any way.

The methods employed for the recovery of scents are: pressing, distillation, maceration, absorption, and, finally, extraction. Before proceeding to deal with these various methods and the details of their performance, it will perhaps not be devoid of interest and practical utility to regard them critically, and ascertain how they have behaved in practice. The practical man will then be in a position to compare them with his own experience, and thus recognise whether his own methods are rational or not.

On a small scale, the method of winning perfumes by pressure suffers from the drawback that, even when the strongest screw presses are used, a considerable portion of the oil is lost owing to a very large number of the oil vessels in the shells remaining unbroken. For this reason it is always advisable to employ hydraulic presses, since these give an amount of pressure unattainable by other means. Hence this method is out of the running for workers on the small scale. Furthermore, it is merely of low value to the maker of cosmetic preparations, being only suitable for application to a few substances yielding cheap scents. The employment of a hydraulic press is only profitable for working on the large scale, and especially for the production of various fatty oils (which, besides, are frequently used in compounding cosmetics).

The distillation method is extensively practised, and is capable of application to a large number of substances, such as the ethereal oils present in caraways, aniseed, lavender, mace, fennel, nutmegs, cinnamon, etc., all of which are obtained by this means.

Though suitable forms of apparatus exist for the performance of this method on the small scale, for those who are unprovided with a steam boiler, and are obliged to use fire heat, nevertheless the products for the most part do not come up to the high

standard of purity now required. In fact, steam distillation is indispensable for working on the large scale, which latter is the only profitable way of conducting these operations. Despite sundry "ifs" and "buts," the distillation method must always be retained, more especially for dealing with vegetable matter rich in scented constituents, or voluminous in bulk, and with woods. On the other hand, distillation is not recommended for working on the small scale, or where very fine perfumes are in question.

The maceration process, which is chiefly practised in France, is based on the property of fatty substances of absorbing perfumes with avidity, and of parting with a large proportion of same when treated with very strong alcohol. According as the maceration of the flowers is effected by the aid of fats of high or low melting point, the products are classed as pomades or scented oils. By using the same fatty matter for repeatedly treating fresh quantities of flowers, the manufacturer has it in his power to perfume the pomade or oil to any desired extent. These products are indicated by numbers, the higher numbers expressing the stronger products. Fats may be advantageously replaced for this purpose by glycerin, which also absorbs the perfume from the flowers in a very rapid manner.

The first point to be mentioned in connection with this method is, that however long the fat is exposed to the action of alcohol, the extraction of the perfume remains incomplete, a small portion being always left behind. True, these residual perfumed fats and oils can be sold as such; but this circumstance is immaterial so far as concerns the appreciation of the value of the method, the point at issue being to determine whether the method attains the object for which it is employed, namely, the complete and pure extraction of the perfume. In addition, it must be remembered that some vegetable perfumes are unable to stand even the slight increase of temperature entailed by the maceration process.

A modification of the maceration method consists in treating the flowers with fats or oils at the ordinary temperature, and is known as "enfleurage." Again, fat may be replaced by carbon dioxide for this purpose; but this treatment, which is troublesome and lengthy, exhibits no special advantage, except perhaps in the avoidance of high temperatures.

The extraction method is based on the capacity exhibited by certain volatile liquids, such as sulphuric ether, chloroform, petroleum ether, and carbon disulphide, for quickly extracting the perfumed constituents from flowers, and then leaving the extractives behind in a pure state when themselves evaporated by the application of heat.

This method offers the undeniable advantage of being equally useful for working on the small scale as the large; and this is a point worthy of mention. It is, however, attended with the inconvenience of having to get rid of the solvent after the latter has performed its allotted task, and become merely a source of trouble. True, the solutions of perfume in such solvents can be volatilised at a very low temperature, and the solvents themselves can be recovered in the condenser for use over again; but the method does not preclude the volatilisation of the perfume, and its alteration by the action of atmospheric oxygen, even when the temperature of 36° C. is employed in the case of sulphuric ether, 65° C. with chloroform, 50° C. with petroleum ether, and 48° C. with carbon disulphide.

It is thus evident that each method is defective; and, in fact, there is really no method known which could be characterised as the best, and most commendable for use in all cases. This is probably the reason why attempts are continually made to increase the production of perfumes by artificial means, *i.e.* the synthetic process.

Endeavours were made for a long time to isolate the perfumes of sweet-scented flowers by using readily volatile

solvents, in place of the earlier method of impregnating fats and oils with the corresponding perfumes by maceration or enfleurage. These attempts, however, were unattended by any striking amount of success, although the utility of the method is beyond doubt. The chief drawback resides in the insufficiency of the solvents employed, whether owing to the fact that their boiling points are too high to permit the recovery of the extracted perfumes in an unaltered or pure condition, or that a small proportion of the solvent remains obstinately adherent to the extracted matter, its removal then entailing the application of such a high temperature that the perfume suffers alteration. In the year 1891 it was widely stated in the technical press that a solvent free from the aforesaid defects had been discovered in methylal. The low boiling point (42° C.) of this substance, coupled with the readiness with which it volatilised without leaving a trace behind, led to the expectation that it would prove suitable for isolating even the most delicate perfumes; and these hopes were said to have been confirmed by practical experiments, for example, with violet blossoms. Nevertheless, this solvent does not appear to have been fully adopted in practice, no further experiences with it having been made public since that time.

A point specially worthy of mention is the circumstance that in the preparation of perfumes the most divergent gradations can be obtained, often in a surprisingly simple manner. Apart from this, however, entirely new perfumes can be prepared by the mixing and suitable treatment of various components.

Thus it was observed by E. Soxhlet that a mixture of Peruvian balsam with traces of carbolic acid produced the same scent as castoreum; whilst acetic acid and traces of clove oil closely imitate the smell of fresh strawberries. The odour of fresh poplar buds can be easily produced by suitably mixing an alcoholic solution of guaiacum with lavender oil. An alcoholic solution of rose oil with traces of patchouli

tincture, gives the well-known "white rose" perfume; and rose oil with traces of musk tincture gives moss rose. A close imitation of the scent of lilies of the valley is obtained by mixing ylang-ylang extract with traces of tincture of cardamoms; and elder flower extract is prepared by adding ylang-ylang and linaloe to turpentine oil. Eau de Cologne can be produced by adding 1 gram of neroli oil to 100 grams of alcohol. Every lover of this scent will admit that the product obtained in this manner is superior to all others, and, in fact, when kept for some time, it is really excellent, the only objection being that it is more expensive than mixtures prepared from the ordinary recipes, which, in addition to small quantities of neroli oil, contain such oils as lavender, rosemary, lemon, etc. The well-known and highly appreciated handkerchief scent, Ess Bouquet, can be imitated by the following recipe: Jasmine extract, reseda extract, and violet extract, 1000 grams each; orris root, 1500 grams; liquid storax, 5 grams; and ambergris, 2 grams: the whole digested together for a month and then filtered.

English people prefer strong scents, so for this market the flower perfumes are usually strengthened by additions of ambergris, musk, storax, benzoe, cumarin, and patchouli.

A number of new scents have been discovered by chemists, and these are now obtainable so cheaply as to make it worth while attempting to use them in the preparation of extracts and fancy scents. To this class belong heliotropin, cumarin, vanillin, nerolin, terpinol, etc.; and these, when dissolved in alcohol and mixed with a suitable fixing agent, such as musk, ambergris, etc., furnish really excellent and cheap handkerchief perfumes. An excellent heliotrope extract can be prepared from heliotropin by digesting 1 gram of heliotropin, 0.2 gram of vanillin, 3 grams of neroli oil, 1 gram of oil of bitter almonds, 0.1 gram of musk, and 5 grams of benzoin, in 1 litre of alcohol for a month.

It is well known that the addition of just sufficient fresh milk to a solution of ethereal oils as to cause incipient separation of the oil, contributes to the durability and intensity of the perfume of the mixture, the small amount of butter fat in the milk sufficing to retain the aroma, which would be lost by the volatilisation of the alcohol. The effect is still greater when hot milk is used; but it is always necessary to know in advance just how much milk is required to produce the incipient separation of the perfume. Consequently the following method must be adopted:—

The ethereal oil preparation being ready, a small quantity is taken, say about 30 c.c., and placed in a graduated burette, where sufficient water is added—drop by drop towards the end—to produce a distinctly opalescent mixture, without causing the separation of any drops of oil after standing for an hour. It then becomes a question of ascertaining how much milk will be necessary to displace the quantity of water used. Cow's milk contains on the average 80 per cent. of water; then, assuming we have used, in the above case, 15 minims of water (30 c.c. being equal to 1 fluid ounce), it will be necessary to multiply this figure by 1.055 in order to find the quantity of milk required. The multiplier is found in the following manner:—1 fluid ounce (=30 c.c.) of milk weighs about 470 grains (30.4522 grams); and the sp. gr. of milk is 1.030 at 16° C. These 470 grains contain 433.6 grains of water (88 per cent.), and 36.4 (12 per cent.) of solids, the 433.6 grains of water corresponding to 455 minims. Hence we have the sum: $455 : 480 = 15 : x$; $x = 15 \times 1.055$. In our special case x is equivalent to 15.25; or, in round numbers, 16 minims. To ascertain the weight, instead of the volume, of milk required, all that is necessary is to multiply the volume in minims, etc., by 1.033. (1 pound is equivalent to 12 ounces of 8 drams, of 3 scruples, of 29 grains or minims.)

The various methods of preparing scents will now be described.

CHAPTER II.

PREPARATION OF PERFUMES.

THE PRESSURE METHOD.

THIS method is only applicable to certain perfumes occurring abundantly in various parts of plants. Thus the skins of oranges, lemons, quinces, etc., contain considerable amounts of ethereal oils, enclosed in oil vessels which can be easily detected under the microscope. On subjecting the said parts to powerful pressure, the oil vessels are ruptured and the contents escape. As a rule, the operation is conducted in screw presses, provided with strong iron spindles, the materials being placed in strong press cloths made of horse hair or linen cloth, these being then laid between iron plates and exposed to gradually increasing pressure. Comparative tests have shown that, even when the strongest screw presses are used, a considerable proportion of the oil is lost, through a large number of the oil vessels being left unbroken. For this reason it is always preferable to employ hydraulic presses, since these furnish a degree of pressure unattainable by any other means. In hydraulic presses for this class of work, the piston or plunger fits exactly in a hollow iron cylinder, which is perforated all round like a sieve, and contains the substances to be pressed. When pressure is applied, the matter expressed from the materials flows through the perforations in the cylinder, leaving the residue behind in the form of a compact cake, which latter is quite free from oil.

The expressed liquid contains an aqueous fluid in addition to the oil, the whole being rendered milky by the presence of vegetable fibres, mucilage, etc. This is collected in a high glass cylinder, which is then set aside in a place where it will be quite free from vibration. After a few hours' rest the liquid will be found to have separated into two layers, the lower one consisting of the water, vegetable mucilage, etc., and the upper one of nearly pure oil. This is carefully decanted and passed through a filter, until perfectly clarified, a double layer of filter paper in a glass funnel covered with a glass plate being used.

A better method of separating the oil and water is by means of a separating funnel, which can be made by cutting the bottom off a bottle and passing a glass tube, about 6–10 mm. in diameter, through the cork. By attaching to this tube a short length of indiarubber tubing, fitted with a pinchcock, this extemporised funnel can be used for separating even the last drops of water from the press runnings.

For the manufacturer of cosmetics, the method of extracting perfumes by pressure is of little value, since it is only applicable to a few materials which furnish cheap scents.

THE DISTILLATION METHOD.

The vapours of many perfumes and ethereal oils have the property of being carried away in such quantity by steam that they can be extracted from the parent plant by this means, though as a rule the ethereal oils have a higher boiling point than water. This method of volatilising any substance and then recovering it by condensation is termed steam distillation, and is capable of application to a large number of substances, the ethereal oils contained in caraways, aniseed, lavender, fennel, mace, nutmegs, cinnamon, etc., being obtained exclusively by this means.

The forms of apparatus used in carrying out this method

vary in accordance with the amount of material to be treated ; and a few of the chief types will now be described.

For those who are unprovided with a boiler, and have therefore to use direct fire heat, an advantageous form of still is one set in brickwork in such a manner that the hot gases from the fire come in contact, not only with the concave bottom of the vessel, but also with the sides, by means of flues arranged in the brickwork. The upper part of the still projects above the setting, and is provided, at one side, with an opening fitted with a screw cover, and serving to fill up the still with water whilst distillation is proceeding. A copper or tin hood is fitted, steamtight, over the edge of the still, and terminates in a pipe leading to the condenser, which consists of a coiled tin pipe fixed in a wooden or metal vessel. The latter is provided with an elbow pipe at the top, and with a tube, bent at right angles and debouching immediately over the bottom of the vessel, its length being such that it projects beyond the edge of the vessel.

The materials to be distilled are introduced into the still and suffused with water ; it is also advisable to provide the still with stirrers, which can be kept in motion until the water actually boils, since otherwise there is a risk of local overheating and burning against the walls and bottom. Such a risk may also be counteracted by introducing into the still a perforated false bottom, placed above the level of the lateral flues ; or by putting the materials under treatment into a wire basket and suspending the latter in the still.

The employment of steam for the recovery of perfumes by the distillation method is highly advantageous, and indeed indispensable for working on the large scale. The stills, which in this case may be constructed of strong tin plate, are unmounted, and are lagged with wood to prevent loss of heat by radiation. Immediately above the concave bottom is fitted a perforated false bottom, on which the charge is laid.

A pipe, passing through the bottom of the still, is connected with a boiler supplying steam under low pressure. This steam-pipe is fitted with a tap, and a second tap serves to run off the water left in the still after the distillation is at an end. When the still is charged, the steam tap is turned on by degrees, and the passage of this current is continued until the operation is complete.

During distillation a mixture of pure water and ethereal oil escapes from the mouth of the condenser. Where the quantities to be dealt with are large, it is advisable to effect the separation of the distilled oil and water by means of a so-called Florentine bottle. This consists of a glass bottle, from the bottom of which an upwardly bent tube projects to such a height that the bend in the tube is on a slightly lower level than the neck of the bottle. During distillation the bottle fills with water, on the top of which floats the separated oil; but when the water reaches a certain level the excess runs away through the bent tube, so that, towards the end, the bottle contains only oil and a very little water.

For work on the large scale this somewhat fragile appliance is replaced by separating bottles consisting of glass cylinders, tapering at the top and bottom, and mounted on a suitable frame. The water settling down below the oil is drawn off through a tap at the bottom; and as soon as the first bottle of the series is full of oil, the liquid coming from the condenser is diverted through a horizontal pipe into the next, and so on.

The complete separation of the oil from water is effected in a separating funnel: a glass funnel, closed by a ground glass plate at the top, and drawn out to a fine point at the bottom of the tapped limb. The contents of the Florentine bottle are transferred to the separating funnel, which is then covered up and left until a clear line of demarcation is observed between the oil and the water. By carefully opening the tap, the whole of the water can be drawn off; the oil is then

transferred to bottles, which are fitted with air-tight stoppers, and stored in a cool dark place.

THE MACERATION METHOD.

Certain perfumes, such as those of the acacia, rose, mignonne, orange blossom, violet, and many other scented flowers, can be less perfectly or less purely recovered by distillation than by maceration, a process based on the property of fats for greedily absorbing perfumes, which they subsequently cede to very strong alcohol.

The maceration process is a very simple operation.

The fat is placed in a porcelain or enamel-lined iron pan, which is then warmed to about 40° (highest 50°) C., by setting it in a shallow vessel containing hot water, the flowers to be extracted being enclosed in bags of fine linen and placed in the fat, where they are allowed to remain for one-half to two days. The bags are then taken out, drained, pressed, refilled with fresh flowers, and returned to the pan. This operation is repeated twelve to sixteen times, or even more, and furnishes a highly scented pomade or oil.

The perfumes are extracted from the pomades or "huiles antiques" by treating the latter with 90-95 per cent. alcohol, which dissolves out the perfumes but leaves the fat untouched. In the case of oils, the extraction is effected by shaking them up with the alcohol in large glass bottles; but the pomades have to be cooled down until solidified, and then cut up into very small pieces for agitation with the spirit. A better plan is to put the pomade in a cylinder with a fine aperture at the bottom, through which, by the aid of a close-fitting piston, the fat is expressed in the form of a fine thread, thus exposing a very large surface and facilitating extraction by the alcohol. The solution obtained as the result of an exposure of several weeks' duration in the above manner is then distilled at a low temperature.

When glycerin is used instead of fat, the method of treatment is the same, except that benzol is employed as the extracting solvent in place of alcohol, owing to the circumstance that the latter dissolves glycerin as well as ethereal oils.

THE ABSORPTION OR ENFLEURAGE METHOD.

In this method also the absorbent property of the fats and oils for perfumes is utilised, but the treatment of the flowers is carried on at the ordinary temperature. In this case the fat—lard—is spread out about $\frac{1}{4}$ inch thick, on glass plates, some 40 inches long and 24 inches wide, which are then laid on wooden frames and covered over with the flowers. The frames are placed one above another and left for one to three days, after which the withered flowers are replaced by fresh ones, and this operation is repeated until the pomade has attained the desired strength.

A useful modification of this method consists in introducing into a tightly closed case a large number of glass plates, covered with lard that has been converted into thin threads by means of a jet. This case is placed in communication with a second one, filled with flowers and provided with an aperture at top and bottom. One of these apertures communicates with the first case by means of a pipe, and a current of air is drawn through the whole apparatus by suitable means, such as a fan driven by clockwork. The air drawn out of the case containing the flowers is laden with perfume, which it surrenders to the fat in passing through the other case.

By the use of this apparatus several important advantages are secured. In the first place, the absorption proceeds at a rapid rate, only a small amount of motive power is required, and the flowers never come in direct contact with the fat, which therefore cannot absorb anything beyond the perfume floating in the air.

The place of fat can be filled by carbon dioxide, in which event the following apparatus is used. A number of lumps of white marble are placed in a large glass vessel, and to them is introduced, at intervals and by means of a funnel, a quantity of hydrochloric acid. The resulting stream of carbon dioxide gas is passed through a washing flask charged with water, and is then led through a chamber of sheet metal filled with the flowers to be extracted, whence it is delivered into a flask filled with strong alcohol and surrounded with cold water. The gas takes up the perfume from the flowers in the state of vapour, and gives up the same to the alcohol, being itself discharged through an effluent tube in the alcohol vessel.

THE EXTRACTION METHOD.

The most suitable solvents for this purpose are petroleum ether and carbon disulphide, these being cheaper than chloroform or ether.

The apparatus employed consists of a tinplate cylinder, which carries at its upper end an annular channel provided with a tap, and can be closed by means of a cover, also provided with a tap. At the bottom of the cylinder is a tube, which can be closed by means of this tap. The cylinder is charged with the flowers to be extracted, and with the solvent, —petroleum ether or carbon disulphide,—the lid is put on, and water is poured into the annular channel, thus forming a hermetical seal. After the extraction is completed, which takes about thirty to forty minutes, both taps are turned on, and the liquid is run into the retort attached to the apparatus. If it be desired to repeat the extraction, the water seal is removed and a fresh quantity of solvent is poured over the flowers through the lid of the vessel.

Having thus seen how the perfumes can be extracted and recovered from the materials in which they are contained, we will now proceed to consider the substances which possess

any importance in connection with the preparation of cosmetics.

It is advisable to distinguish between—

1. Chemical products.
2. Animal substances.
3. Oils and fats.

CHAPTER III.

CHEMICAL AND ANIMAL PRODUCTS USED IN THE MANUFACTURE OF COSMETICS.

CHEMICAL PRODUCTS.

ETHER, which is also known as "sulphuric ether," is prepared by distilling alcohol with sulphuric acid and rectifying the product by a second distillation. In the pure state, ether is a thin-flowing, strong-smelling, and inflammable liquid. The sp. gr. is 0.736, and the boiling point 35° C. Ether possesses high solvent powers for ethereal oils, resins, and fats. Being highly volatile, ether rapidly disseminates its vapour through the air; and as this vapour is exceedingly inflammable, it is necessary to keep all lights away from the vicinity of bottles containing the liquid in question.

It is well known that in mixing water and sulphuric acid, the latter should always be poured into the water, and not *vice versa*. According to Orlow, the reverse procedure must be followed in mixing ether and sulphuric acid, since he observed that when this acid is poured into ether, violent spurning, sometimes even accompanied by explosion, occurs; whereas, when the operation is performed in the converse manner, it is free from danger. The ether used must be water-white, exhibit the true odour and flavour of ether, and must not impart a red coloration to litmus paper. An addition of alcohol increases the sp. gr. Pure ether leaves no residue on evaporation, and should not turn milky when mixed with water, but float on the surface of the latter.

So-called pure commercial ether invariably contains certain

impurities, chiefly organic sulphur compounds, which are left behind as a malodorous residue on the spontaneous evaporation of the ether. The name of sulphuric ether is all the more applicable, owing to the presence of free sulphur in solution. This may be readily detected by agitating the suspected sample with a little pure bright metallic mercury in a test tube. If the amount of sulphur be only very small, the surface of the mercury merely turns a dull grey; but when a relatively large amount of sulphur is present, the entire liquid becomes grey or black from mercury sulphide.

ALCOHOL.

Pure alcohol is a thin liquid, of sp. gr. 0·8095, boiling at $78^{\circ}4$ C., but requiring a temperature far below -100° C. to effect its solidification. By evaporating solid carbon dioxide with ether, Dewar obtained a temperature which is estimated as about -200° C., and which caused alcohol to freeze as a crystalline solid mass, exhibiting the peculiarity of not suddenly becoming liquid when thawed, but of first passing through an intermediate viscous stage resembling glycerin. Alcohol is characterised by a high solvent power for resins, balsams, and ethereal oils. For the production of absolute alcohol, Yoon employed calcium carbide. When this last named substance is broken into lumps and treated with 90–95 per cent. alcohol, it is briskly attacked and continues to liberate acetylene so long as any water remains in the alcohol; as soon, however, as all the water has been absorbed by the carbide, the evolution of gas ceases. Hence calcium carbide affords a means of testing alcohol for the presence of water. The resulting alcohol no longer gives a precipitate with barium alcoholate; consequently calcium carbide is just as delicate a reagent as the latter, and enables absolute alcohol to be obtained from 95 per cent. (or even 90 per cent.) spirit in at most two distillations. For this purpose the alcohol to be rectified is mixed with 25 per cent. by weight

of calcium carbide, repeatedly agitated, left to settle until the liberation of gas ceases, and is then distilled. The distillate is freed from any accompanying acetylene by adding a small quantity of calcined copper sulphate and distilling anew.

For the preparation of absolute alcohol in glass vessels and on the small scale, Habermann has recommended that the lime used for this purpose should be added not to the still direct, but allowed to act in a vessel communicating therewith. This vessel consists of a cylinder, drawn out to a fine tube at the lower end, and inserted in the path of the distilled vapour on its way to the condenser. This cylinder contains a narrow cylindrical roll of iron-wire gauze, closed at the upper end, and resting with its lower end against the tapered portion of the glass cylinder, whilst the top extends to within a few centimetres of the stopper of the latter. The annular space between the gauze and the glass is filled with a weighed quantity of quicklime, in lumps as large as hazel nuts. The still being charged with a corresponding quantity of (at least 95 (vol.) per cent.) alcohol, the attachment containing the lime is tightly fixed on and connected with the condenser. The still is then heated on the water bath, slowly at first, so that only a small quantity of the alcohol distils over during the first two hours, at the end of which time the receiver is changed. This vessel is connected with the condenser by a cork provided with a double bore fitted with a calcium chloride tube open at both ends. Should the quicklime be found to have become wet and pappy during distillation, the operation must be suspended and the lime replaced by a freshly charged tube, since in the condition mentioned the lime has not only lost its power of absorbing moisture from the alcohol, but is capable of ceding such to the alcohol.

In order to deodorise alcohol for cosmetic purposes, 5 parts of alcohol should be left for several days in contact with a mixture of powdered charcoal 30 parts, quicklime 60, and

powdered alum 8 parts, the whole being frequently agitated, and finally filtered.

Potassium permanganate also is recommended as an excellent means for rectifying alcohol, 2-3 c.c. of an aqueous solution of permanganate—15.89 grams of this salt to 100 c.c. of water—being sufficient for 1 litre of badly flavoured spirit. The alcohol can be separated from the resulting black precipitate by decantation.

Another excellent means for rectifying once-distilled alcohol is found in sodium peroxide. The quantity employed per hecto-litre varies from 100 to 150 grams of sodium peroxide, added twenty-four hours previous to rectification. By this means the regular progress of the reaction is ensured, the separation of the impurities is accentuated, and a higher yield of alcohol of pure flavour is obtained.

The usual tables for preparing 90 per cent. wine spirit from stronger alcohol are arranged in percentages; but it is better to employ a table based on sp. gr., as follows:—

Sp. gr. at 15° C.	1000 Grams of 90 % Spirit contain		Sp. gr. at 15° C.	1000 Grams of 90 % Spirit contain	
	Alcohol Grams.	Water Grams.		Alcohol Grams.	Water Grams.
0.8070	894.6	105.4	0.8130	914.5	85.5
0.8075	896.2	103.8	0.8135	916.3	83.7
0.8080	897.8	102.2	0.8140	918.1	81.9
0.80833	898.88	101.12	0.8145	919.9	80.1
0.8085	899.4	100.6	0.8150	921.7	78.3
0.8090	901.05	98.95	0.8155	923.5	76.5
0.8095	902.65	97.35	0.8160	925.3	74.7
0.8100	904.3	95.7	0.8165	927.13	72.87
0.8105	905.95	94.05	0.8170	928.95	71.05
0.8110	907.6	92.4	0.8175	930.8	69.2
0.8115	909.3	90.7	0.8180	932.6	67.4
0.8120	911.0	89.0	0.8185	934.4	65.6
0.8125	912.75	87.25			

CHLOROFORM.

Chloroform is obtained by distilling wood spirit, common salt, and sulphuric acid, and rectifying the distillate. One method of preparing pure chloroform is based on the precipitation of chloroform salicylide from impure chloroform. The salicylide is obtained by the action of phosphorus oxychloride on salicylic acid, and this class of substances give, with chloroform, products that are only sparingly soluble, if at all, and crystallise in an excellent manner. The combination is very unstable, the chloroform being liberated on the application of gentle heat; and it can be prepared in this manner, in a perfectly pure state, just before use. Chloroform has an agreeable odour, a sweet taste, boils at 61° C., and is of sp. gr. 1.48. On account of its high solvent powers and low boiling point, it is largely used for the extraction of vegetable perfumes.

Numerous experiments have been made on the decomposition and preservation of chloroform. Allen showed that an addition of sulphur, in the proportion of 1 : 1000, retards the decomposition of chloroform in daylight. In this decomposition, oxygen plays a considerable part, and chlorine is always present in the first stage of the reaction; it would therefore seem as though the sulphur acts as a reducing agent, and that any readily oxidisable substance soluble in chloroform would have the same effect. With this object Dott divided a sample of chloroform into four portions, a very small amount of morphia being added to the first, gallotannic acid to the second, hypophosphorous acid to the third, and nothing at all to the fourth. After a few weeks, the pure chloroform exhibited signs of decomposition, as evidenced by the smell and on the addition of silver nitrate. The others remained entirely unaltered, even after exposure to light for a month longer.

BENZOIC ACID.

The preparation of benzoic acid from benzoin can be performed by any one in a very simple manner, by digesting this resin with 3-4 parts of strong acetic acid. The resulting brown solution is decanted and poured into 4 parts of boiling water. The resin is deposited as a greyish brown mass, and the liquid, after filtration and cooling, furnishes crystals of benzoic acid (most of that present in the solution), a further quantity being obtainable by concentrating the residual liquid after partial saturation with quicklime. In working on a large scale, the acetic acid would, of course, be recovered from the liquor.

The resin thrown down from the acetic solution has an agreeable and aromatic odour when dried and melted, and can be used up for other purposes. On account of the preservative qualities of benzoin, it is advisable to add this by-product to the pitch used for lining beer and other casks.

BENZOL AND BENZINE.

According to the raw materials from which it is prepared, namely, coal tar or petroleum, the product is known as benzol or benzine, the former having the sp. gr. 0·850, and boiling at 82° C., whilst the latter the sp. gr. 0·730, and a variable boiling point which does not fall below 130° C. Benzol from coal tar is exclusively used by the makers of coal-tar dyes; but for cosmetic purposes the petroleum benzine is the only one coming under consideration, as it is used for the extraction of perfumes from flowers.

A simple distinguishing test for benzol and benzine is afforded by iodine, a small crystal of which will impart a carmine red coloration to the former, and a violet tinge to the latter. This reaction is very reliable, and also applicable to mixtures of benzol and benzine. Another simple and reliable test is by agitation with a little alcohol, which produces considerable

turbidity in the case of benzol, but leaves benzine perfectly clear. To perform this test, a few drops of 95 per cent. alcohol are poured into a dry test tube and then thrown out again, so as to leave only a very minute quantity behind; 1–2 c.c. of the suspected benzine is then run in, and the whole well shaken up. A turbid, milky liquid indicates the presence of benzol; but if the liquid remain quite clear, the heavier drops quickly separating from the rest, only benzine is present. The alcohol test is simpler than that with iodine, and gives equally reliable indications, provided the method be performed as above.

To rectify benzine, Gawalowski recommends the following treatment:—

When a malodorous benzine is treated with an addition of about 1–2 per cent. of free fatty acids, the latter are dissolved. About $\frac{1}{4}$ per cent. of tannin is then added, and the whole well mixed together. Finally, an addition is made of sufficient potash or soda lye, or milk of lime, to saponify the fatty acids and neutralise the tannin. After shaking the mixture and leaving it to stand a while, the milky liquid separates into two layers, the mucilaginous soapy matters sinking to the bottom, and leaving the clear, colourless, and deodorised benzine floating above. This, when decanted and filtered, gives a product capable of immediate utilisation for many technical purposes, and convertible, by redistillation, into a perfectly pure benzine.

Beef fat, olive oil, or other fats and oils, may be used for the above treatment, provided the same be free from rancidity. The olein—oleic acid—used by candlemakers can also be employed, but should first be shaken up with a 0.1 per cent. solution of soda, in order to eliminate the malodorous liquid fatty acids, butyric acid in particular.

Another method of deodorising benzine is as follows:—

$4\frac{1}{2}$ litres of benzine are well mixed with $\frac{1}{4}$ litre of sulphuric acid, $1\frac{3}{4}$ litres of water, and 30 grams of potassium permanganate, then left for twenty-four hours, and the decanted benzine

afterwards shaken up with a solution of $7\frac{1}{2}$ grams of potassium permanganate and 15 grams of soda in 1 litre of water.

AMMONIUM CARBONATE.

Commercial ammonium carbonate is a mixture of ammonium mono- and bicarbonate, and may be prepared by heating an intimate mixture of 1 part of ammonium sulphate and 2 parts of chalk in horizontal cylindrical, cast-iron retorts, 7–10 feet long and 20 inches in diameter, the vapour being collected in stone chambers, where the salt settles down on the walls and is afterwards chipped off. Behind these chambers is a coke scrubber containing sulphuric acid for the vapours not condensed in the chambers, together with free ammonia and ammonium sulphate. Seventy-six parts of chalk are consumed for each 100 parts of ammonium sulphate, and 59–60 parts of ammonium carbonate are obtained. The crude salt is usually sublimed over again. The sublimation and purification are performed at a lower temperature, not exceeding $65\text{--}70^{\circ}\text{C.}$, the waste heat from the retorts being employed to heat a water bath for this purpose. The iron pans used are surmounted by unsoldered leaden cylinders formed by rolling sheet lead, and open at both ends, the upper ends being covered with leaden plates and a luting of paper. When the operation is complete, the cylinders are cooled down, the paper luting removed, and the sheets unrolled to enable the product to be taken out. When pure, ammonium carbonate will volatilise, without leaving any residue, if heated on a bright piece of platinum foil.

Ammonium carbonate is readily soluble in cold water,—25 : 100,—but is decomposed by hot water ; it is only sparingly soluble in alcohol.

OIL OF BITTER ALMONDS.

Oil of bitter almonds consists of benzoyl hydride or benzaldehyde. It may be prepared by artificial means, but this

method is not adopted in practice, the oil being obtainable more cheaply from bitter almonds.

Pure oil of bitter almonds is colourless, of high refractive power, and with a very intense odour of bitter almonds. By the action of the air it is gradually changed into an inodorous and colourless crystalline mass, consisting of benzoic acid. This change is facilitated by light, and therefore the oil should always be stored in hermetically closed bottles, protected by an opaque envelope.

The oil is adulterated with nitrobenzol, alcohol, and light ethereal oils. Nitrobenzol may be detected by placing 10 drops of the oil in a narrow test cylinder along with 5 c.c. of 90 per cent. alcohol and 5 c.c. of water (or 10 c.c. of 45 per cent. alcohol), closing the cylinder with the finger, and gently turning the mixture round twice. Under this treatment, pure oil of bitter almonds is immediately dissolved to a clear solution; but if even only one 1 per cent. of nitrobenzol be present, the liquid is clouded and the drops of nitrobenzol can be seen floating about previous to collecting at the bottom of the vessel. The formation of merely a turbid solution indicates adulteration with other ethereal oils. The test recommended by Maisch consists in dissolving 1 gram of oil in 8 grams of alcohol, adding one gram of caustic potash, and warming the whole until one-third of the alcohol has evaporated. Pure oil gives a brown liquid, but in presence of nitrobenzol a brown resin is deposited. According to Hofmann, a bluish violet coloration is obtained, in presence of nitrobenzol, on diluting the oil with ether, adding a little hydrochloric acid and sulphuric acid, and afterwards a little alcohol and a few granules of zinc, the mixture being shortly after neutralised with potash, shaken up, the ether allowed to elaborate spontaneously, and the liquid treated with a few drops of bleaching powder solution. A test, introduced by Wagner, is based on the different sp. gr. of mixtures of bitter almond oil and nitrobenzol, and on the

readiness with which the former combines with alkali bisulphites. Commercial oil of bitter almonds varies in sp. gr. between 1.040 and 1.044; nitrobenzol, between 1.180 and 1.201.

The weight per given volume of mixtures of oil of bitter almonds and nitrobenzol is as follows:—

5 c.c. of pure oil of bitter almonds weigh	5.29 grams.
5 do. do. containing 25 per cent.	
of nitrobenzol, weigh	5.39 do.
5 do. do. do. 50 do. do.	5.57 do.
5 do. do. do. 75 do. do.	5.75 do.
5 do. of pure nitrobenzol	do. 5.90 do.

If 5 c.c. of commercial oil of bitter almonds be shaken up in a graduated test glass with 35 c.c. of a concentrated solution of calcium- or sodium bisulphite, and the mixture is then made up to 50 c.c. with water, any nitrobenzol present will float on the top, and the amount can be approximately estimated. This treatment will also cause the separation of any extraneous ethereal oils. To facilitate the collection of the nitrobenzol, it is advisable to add 5 c.c. of benzine to the oil, on commencing this test, the added volume being deducted from the total in the supernatant layer of nitrobenzol.

Oil of bitter almonds is also adulterated with alcohol, and in such event exhibits a lower sp. gr. than usual. Should the alcohol amount to 8–10 per cent., the addition of a twofold volume of nitric acid (sp. gr. 1.42) causes the liquid to swell up in a few minutes and disengage nitrous fumes.

The presence of artificial oil of bitter almonds in the natural oil can be detected in the following manner:—A strip of filter paper, impregnated with a few drops of the suspected oil, is ignited, and the products of combustion are caught in a damp beaker. Water is then added to the beaker, and the liquid is filtered and treated with silver nitrate solution. Should the artificial oil in question be present, a white precipitate will

be obtained, owing to the chlorine invariably accompanying the said oil.

BORAX.

In preparing borax from boric acid, a quantity of soda crystals (*e.g.* 1300 kilos.) is dissolved in water, and the solution raised to 100° C. by the aid of steam, whereupon a corresponding quantity (1200 kilos.) of purified commercial boric acid is added in small portions (4–5 kilos.).

Borax is soluble to the extent of 6 parts in 100 of cold water, or 2 to 1 in hot water; it is insoluble in alcohol.

In purchasing borax, attention should be bestowed on the colour: pure borax is quite white, whereas a reddish coloration indicates contamination with ferric oxide.

CUMARIN.

Cumarin occurs in the Tonka bean, the blossoms of melilot trefoil, the leaves of giant clover, the fruit of *Myrosoton toluiferum*, in woodruff, rue, and sweet vernal grass. It is artificially prepared by treating sodium salicylal with acetic anhydride. Pure cumarin forms handsome white acicular crystals, which possess a very agreeable odour; they melt at 67° C. The rates of solubility are as follows:—

Cumarin dissolves in 400 parts of cold water (at 15° C.), or 45 parts of boiling water; is readily soluble in alcohol and ether, as well as acetic acid and ethereal oils.

ACETIC ACID (GLACIAL ACETIC ACID).

In its purest form acetic acid is a colourless liquid, with a strongly acid taste and smell; it solidifies to colourless crystals at 8° C. Like alcohol, this acid dissolves ethereal oils, and is employed in the preparation of numerous toilet vinegars and washes.

In chemical works, glacial acetic acid is prepared in a cheap and pure form by distilling sodium acetate with sulphuric acid.

The reagent for detecting mineral acids in acetic acid is rosaniline hydrochloride. One c.c. of the acetic acid to be examined is poured into a shallow porcelain dish, and treated with one drop of an alcoholic solution of rosaniline hydrochloride—25 grams of fuchsine in 100 c.c. of 90 per cent. alcohol. If the sample be pure, no alteration in the red-violet colour of the fuchsine occurs on mixing the two liquids; in fact, the colour is rather intensified. In presence of mineral acids, however, even as low as 1 per cent., the colour changes to a dirty yellow. The characteristic colour of the fuchsine is restored by neutralisation with pure alkali.

For detecting the presence of sulphuric acid or hydrochloric acid, Hager proposes warming 1 large or 2 small drops of gurjun balsam—East Indian copaiba balsam—with 35–40 drops of pure acetic acid, in a not too wide test tube, and warming the same with agitation. On then adding at most three drops of the suspected sample, a blue violet coloration ensues, at once if hydrochloric acid be present, or more slowly in presence of sulphuric acid. The development of this coloration may be accelerated by warming the mixture above 60° C. This reaction requires only two to three minutes for its performance.

NITROBENZOL.

Nitrobenzol is formed by treating benzol, or a mixture of this and toluol or higher homologues, with strong nitric acid or a mixture of nitric and sulphuric acids, washing the product with water and sodium carbonate, caustic soda, or ammonia, driving off the unaltered hydrocarbons with steam, and rectifying the residue.

The substances technically comprised under the name nitrobenzol generally form yellowish or brownish liquids, with a high refractometer index; they are of oleaginous consistence, and possess an intensely sweet, empyreumatic taste, are sparingly soluble in water (to which, however, they impart their

odour), are miscible to a clear liquid with alcohol, ether, chloroform, carbon disulphide, benzol, benzine, petroleum ether, and most of the fats and the ethereal oils. They can be distilled without alteration, but undergo decomposition, occasionally accompanied by explosion, when rapidly and strongly heated. According as the nitrobenzol is prepared from light or heavy benzol, the product is known as light or heavy, these terms applying to the boiling point and not to the density.

Light nitrobenzol, also known as nitrobenzol for blue and black (in the dyestuff trade), is an almost pure article, pale yellow in colour, and (in comparison with the others) of agreeable smell, recalling bitter almonds. The sp. gr. is 1.186–1.2; boiling point, 205–210° C.; solidifying (crystallising) point, 3° C. It serves as a perfume,—artificial oil of bitter almonds, mirbane oil, mirbane essence,—especially for fats and soaps. Only this light nitrobenzol, however, is suitable for this purpose. An extra fine quality, nearly or quite colourless, and water-white, is prepared, for this use, from pure crystallisable benzol, and purified by washing with potassium bichromate and sulphuric acid, followed by rectification with steam.

The detection of nitrobenzol as an adulteration in oil of bitter almonds has already been dealt with under the latter heading.

The following reaction is suitable for the detection of nitrobenzol in soaps, etc. Two drops of carbolic acid, three of distilled water, and a piece of caustic potash the size of a pea, are placed in a porcelain basin and warmed carefully until boiling point is reached. On then at once adding a few drops of a mixture of water and nitrobenzol and continuing to boil the liquid, it will be found that a carmine red coloration is produced around the edges, which colour is changed to emerald green by the addition of a saturated solution of bleaching powder. To detect nitrobenzol in liquids, and isolate it from same, the liquid in question is concentrated to a small volume, treated

with quicklime, cooled, and finally extracted with ether. The solvent is evaporated to dryness, taken up by shaking with a small quantity of distilled water, and treated as above. Soaps are dissolved in water, treated with an excess of quicklime, extracted with ether, and treated in the same manner.

With a few modifications, the following process can be utilised for our purposes. For detecting nitrobenzol in confectionery, fruits, etc., Dammer proposed the following method :—

Nitrobenzol can be readily detected by shaking up a sufficient quantity of the fruit with ether, evaporating the latter, treating the residue with a little powdered iron and an aqueous solution of sulphuric acid, concentrating, decomposing the residue with a little caustic soda solution, extracting the resulting aniline with alcohol, driving off the solvent after acidifying with sulphuric acid, and finally adding a little potassium chromate and sulphuric acid. A blue coloration indicates nitrobenzol.

Now, in many cases this process is too complicated, and, as for small quantities, unreliable; and as it is generally merely a question whether the smell of bitter almonds in a soap is due to the presence of natural or artificial oil of bitter almonds, the ready oxidisability of the true oil forms a convenient test. When crushed bitter almonds are stirred with water until they exhibit a powerful odour, the smell of hydrocyanic acid can be destroyed by warming them with a little dissolved soda and a few drops of a ferrous salt. On then adding sufficient potassium permanganate, the odour of oil of bitter almonds is also destroyed. This treatment gives no results in the case of nitrobenzol, and therefore affords a means of detecting the presence of this latter.

ACETIC ETHER.

Small quantities of acetic ether are found in vinegar, French brandy, and certain wines. It is formed by the action of acetic acid on alkalis, by treating an alcoholic solution of potassium

acetate with carbon dioxide, and is obtained by distilling on the water bath a cooled mixture of 1305 parts of concentrated sulphuric acid, 655 parts of 90 per cent. alcohol, and 1000 parts of anhydrous sodium acetate. The distillate is shaken up with lukewarm water, then separated therefrom, shaken up with anhydrous sodium carbonate, dehydrated with calcium chloride, and rectified.

It is a colourless liquid, with an agreeable, refreshing taste and smell. The sp. gr. is 0.924, the boiling point 72.8°C . It dissolves in 17 parts of water at 17.5°C .; is miscible with alcohol and ether, has a neutral reaction, is decomposed into alcohol and acetic acid when in the state of an aqueous solution, and when its vapours are mixed with air and heated. It burns with a smoky flame, and is decomposed by caustic alkalis, the same result being effected by alkali carbonates in the warm.

BUTYRIC ETHER.

Butyric ether (pineapple essence) is obtained by saponifying butter with caustic potash, distilling the soap with alcohol and sulphuric acid, and rectifying the distillate. It is a combustible liquid, boiling at 119°C ., and exhibiting a strong odour of pineapple.

VALERIANIC ETHER.

Valerianic ether is prepared by distilling sodium valerianate with alcohol and sulphuric acid; it boils at 133°C ., and disengages an odour of apples.

ESSENCE OF JARGONELLE PEAR.

This substance, chemically known as amyl valerianate, is prepared from the amyl alcohol so largely formed as a by-product of the manufacture of potato spirit. This amyl alcohol is carefully heated in a still with potassium bichromate and sulphuric acid. The distillate possesses an agreeable odour of pears, and boils at 196°C .

NITROUS ETHER.

Nitrous ether is obtained by the action of nitric or nitrous acid on alcohol. With this object, 8 parts of fuming nitric acid, 4 of water, and 9 of alcohol (90 per cent.), are poured, in superimposed layers, into a flask. This is left to stand for several days, after which time the ether is drawn off. Another method is to pass well-cooled nitrous acid, formed by the action of nitric acid (1·3) on starch, into a cold mixture of 2 parts (85 per cent.) alcohol and 1 of water, the liberated ether being collected in a condenser.

It is a colourless liquid, with an agreeable fruity smell and peculiarly stinging taste; the density is 0·900 at 15·5° C.; the boiling point 17° C. It produces great cold on evaporation, is readily inflammable, sparingly soluble in water, miscible with alcohol and ether, decomposes on storage, liberates nitrogen, turns acid, is gradually decomposed by caustic potash, and furnishes alcohol, water, ammonia, and sulphur when treated with sulphuretted hydrogen.

GLYCERIN.

Glycerin is produced on a large scale as a by-product of candle making, and is put on the market in a very pure state. In this industry, fat is converted into stearin by heating it with either lime or sulphuric acid, or with high-pressure steam, whereby the glycerin is obtained as an aqueous solution, of a strength not exceeding 4–5° Be. In the first-named method of saponification, however, it is contaminated with lime, and in the second with sulphur. After these impurities have been eliminated as completely as possible, the glycerin solution is concentrated to about 24–30° Be., in which condition it is a more or less bluish syrupy liquor, and forms commercial crude glycerin. There are two methods of refining this article: distillation with superheated steam, and decoloration

by means of bone black, the product from the first method being known as distilled glycerin, and that from the second as refined glycerin: the former, however, is the sole means of furnishing a chemically pure product. When chemically pure, glycerin should not give any turbidity either with ammonia and ammonium oxalate, or with nitric acid and silver nitrate, nor should it give any coloration when treated with the last-named reagent alone, or evolve any odour when rubbed in the hand.

Glycerin is miscible with water in all proportions, and therefore is met with in various degrees of dilution in commerce. The percentage of glycerin in aqueous solutions of different densities (at 12-14° C.) is set forth in the subjoined table, compiled by Lang:—

Glycerin %.	Sp. gr.	Glycerin %.	Sp. gr.	Glycerin %.	Sp. gr.
100	1·2641	82	1·2212	64	1·1702
99	1·2664	81	1·2185	63	1·1671
98	1·2637	80	1·2159	62	1·1640
97	1·2610	79	1·2122	61	1·1610
96	1·2584	78	1·2106	60	1·1582
95	1·2557	77	1·2079	55	1·1455
94	1·2531	76	1·2042	50	1·1320
93	1·2504	75	1·2016	45	1·1183
92	1·2478	74	1·1999	40	1·1045
91	1·2451	73	1·1973	35	1·0907
90	1·2426	72	1·1945	30	1·0771
89	1·2398	71	1·1918	25	1·0635
88	1·2372	70	1·1889	20	1·0498
87	1·2343	69	1·1858	15	1·0374
86	1·2318	68	1·1826	10	1·0245
85	1·2292	67	1·1795	5	1·0123
84	1·2265	66	1·1764	1	1·0025
83	1·2238	65	1·1733		

Glycerin readily absorbs perfumes, and is employed, in conjunction with these, for the preparation of toilet articles.

HELIOTROPIN.

Heliotropin is obtained by the oxidation of piperic acid with potassium permanganate. It forms very long, lustrous crystals, melting at 37° C., and boiling at 263° C. The odour is similar to cumarin. Heliotropin is only sparingly soluble in cold water, though more easily so in hot; readily soluble in alcohol and ether; and furnishes piperonylic acid when boiled with alcoholic caustic potash. Artificial heliotropin is prepared from white pepper by extracting with alcohol, —0.833,—treating the extract with caustic potash, and purifying the residue by recrystallisation from alcohol. The piperin obtained in this manner is converted into potassium piperate by heating on the water bath for twenty-four hours, under a reflux condenser, in a flask containing an equal weight of caustic potash and 5 to 6 parts of alcohol. The lustrous yellowish plates of potassium piperate deposited on cooling are washed with cold alcohol, recrystallised from hot water, and, when necessary, decolorised with bone black. To prepare piperonal from this product, 1 part is dissolved in 40–50 parts of hot water, and to the hot solution is added a solution of 2 parts of potassium permanganate, run in slowly with constant stirring. A pulpy mass is formed, which is strained, and is washed repeatedly with hot water until it ceases to disengage the characteristic odour of heliotropin. The united liquids are distilled over a naked flame; the distillates are collected in fractions, because the first runnings are the most productive; and from these the greater part of the piperonal crystallises out on cooling, the remainder being recovered by agitation with ether.

The ready solubility of heliotropin in all ethereal oils and alcohol greatly facilitates its use. The amount taken, how-

ever, must be rather large, since, in the case of soap, a good result will not be obtained unless the quantity of heliotropin attains 500 grams per 100 kilos (about $\frac{1}{2}$ lb. per cwt.); whilst for fine, highly scented toilet soaps double that amount is requisite. The perfume is largely increased and improved by an addition of about 100 grams of cumarin; and very agreeable results can be obtained by adding petit grain oil, bergamot oil, or lemon oil, which freshen up the perfume of the heliotropin.

PYROGALLIC ACID.

Pyrogallie acid is obtained by heating well-dried gallic acid with twice its weight of powdered pumice to 215° C. on an oil bath, whereby the first-named substance is decomposed into carbon dioxide and pyrogallie acid. It forms brilliant white, elongated plates, and, when dissolved in water, especially in presence of alkalis, rapidly absorbs oxygen and turns brown. Gold and silver are reduced from their solutions by this acid; and with salts of iron it furnishes blue-black compounds.

LIVER OF SULPHUR.

Liver of sulphur, or potassium trisulphide, is prepared by fusing together potash and sulphur, and forms a liver-coloured mass, which slowly decomposes on exposure to the air, sulphuretted hydrogen being disengaged; consequently the preparation must be stored in tightly closed vessels. When dissolved in water and brought into contact with salts of lead or silver, liver of sulphur produces a black precipitate of lead—or silver sulphide.

CALCIUM SULPHIDE.

On calcining calcium sulphate—gypsum—with carbon, calcium sulphide is formed as a yellowish white mass, which is transformed into calcium hydrosulphide when treated with

water. The hydrosulphide can also be obtained by passing sulphuretted hydrogen gas through milk of lime, and takes the form of a grey mass, smelling strongly of sulphuretted hydrogen. The proportions employed in the preparation of calcium sulphide are: anhydrous gypsum, 7 parts; wood charcoal, 3 parts; and the calcination is conducted out of access of air. Calcium penta-sulphide may be prepared by heating an intimate mixture of slaked lime to incandescence with $\frac{1}{2}$ —1 part of powdered sulphur, and continuing the heating for half an hour longer.

STARCH.

Starch is prepared from various vegetable substances, such as potatoes, rice, arrowroot, sago, etc., and, in the pure state, takes the form of an insoluble white powder, which, under the microscope, is seen to consist of granules built up of a number of superimposed layers. Considerable differences in price exist between the various commercial forms of starch. Purified potato starch can be used for face and toilet powders.

Wroblewski has proposed the following method for the production of soluble starch:—

One hundred grams of rice starch are carefully trituated with a 2 per cent. solution of caustic potash until a perfectly fluid mixture is obtained, and left to stand for three to four hours, after which a further quantity of potash solution is stirred in, so as to make up the volume to 600–800 c.c. This is heated on the water bath until complete liquefaction ensues, and, after continuing to heat for another twenty to thirty minutes over a naked flame, the liquid is filtered, supersaturated with acetic acid, and finally precipitated with an equal volume of 95 per cent. alcohol. The precipitate is redissolved, reprecipitated, and then taken up with a minimum quantity of water. This solution is poured in a

thin stream into a large volume of absolute alcohol, the precipitate being filtered, washed with alcohol and ether, and finally dried *in vacuo*. The yield amounts to 50–60 per cent.

This soluble starch is a snow-white powder, perfectly non-hygroscopic, insoluble in alcohol of 40 per cent. strength and over, and incapable of dialysation. One hundred parts of water will dissolve 3–4 parts to form a slightly opalescent liquid, which is precipitated by the sulphates of ammonia, magnesia, and soda, but not by sodium chloride. Tannin also produces a precipitate, which, however, is soluble in water after having been washed with alcohol. On the other hand, dilute solutions of soluble starch are not precipitated by phosphotungstic acid, phosphomolybdic acid, potassium-mercury iodide, mercuric chloride, or mercurous nitrate; and lead acetate produces merely a faint turbidity.

VANILLIN.

Vanillin occurs in vanilla pods, and is also obtained from calcium vanillate by dry distillation, from coniferin and coniferyl alcohol. It forms colourless prisms, tastes and smells strongly of vanilla, is readily soluble in hot water, alcohol, and ether. It melts at 80–84° C., sublimes, boils at 285° C. in a current of carbon dioxide, has an acid reaction, and forms crystalline salts.

MENTHOL.

Menthol is the stearoptene of oil of peppermint. It is found in this oil, especially the American and Japanese kinds,—Japanese oil containing 50–55 per cent., American oil 20–25 per cent., English oil 40–45 per cent.,—from which it crystallises out in the cold, more particularly from the less volatile portions. Menthol forms colourless prisms, smells strongly of peppermint, has a burning taste, is sparingly soluble in water, but readily so in alcohol and ether; melts at 42° C., boils at 213° C., is volatile and neutral. It dissolves certain salts,—magnesium

sulphate, for example,—and deflects polarised light towards the left.

The following method is given by Kleber for the rapid approximate determination of menthol in oil of peppermint:—

About 5 grams of peppermint oil are carefully weighed out, along with a measured volume (about 5 c.c.) of anhydrous acetic acid, and boiled for half an hour under a reflux condenser. In the meantime, an equal quantity of anhydrous acetic acid is titrated with normal caustic soda and phenolphthalein. After the boiled liquid has been somewhat cooled down, the condenser is rinsed out with water, the rinsings being added to the acetylated mixture, and the whole titrated with normal caustic soda. The difference in the number of c.c. consumed in the two titrations multiplied by 0.156 gives the amount of menthol in the oil. The whole operation only takes fifty minutes.

OIL OF WINTERGREEN.

Oil of wintergreen, or Gaultheria oil, is obtained from the plant *Gaultheria procumbens*, which grows wild in North America. It has a very agreeable odour, and when pure is quite white, otherwise green.

The best method of preparing this oil by artificial means is that of Thayer—

505.470 grams of salicylic acid and 690.850 grams of methyl alcohol are weighed out, the former being added in small portions to the latter in a wide-necked flask until a saturated solution is obtained. The flask is then connected with a reflux condenser and the contents heated to boiling point, after which the liquid is completely saturated with dry hydrochloric acid gas. The flask is next disconnected from the condenser, and the contents are treated with about 10 grams of salicylic acid, whereupon it is again connected up and the solution saturated afresh with gaseous hydrochloric acid. This procedure is repeated until the whole of the weighed-out quantity of

salicylic acid has been consumed, after which the current of gas is passed through the apparatus for about two hours longer. The contents of the flask will then be found to have separated into two distinct layers, the under one being a dark brown oily liquid. The supernatant liquid is poured off, and the oil washed with water until it ceases to redden litmus paper, after which it is distilled by steam. The excess of water is removed from the oil by means of a separating funnel, and the product is then thoroughly dried with anhydrous calcium chloride. The salicylic acid used for this process has a melting point of $156-157^{\circ}\text{C}$., and is dried at 100°C . The methyl alcohol boils between 58° and 72°C ., and has the density 0.820. Special care must be bestowed on the drying of the hydrochloric acid gas. Thayer found it necessary to first pass the gas over anhydrous calcium chloride, and then through three vessels filled with sulphuric acid. The yield obtained from the foregoing quantities of materials was 500 grams of a colourless or slightly yellow oil, of agreeable odour, a sp. gr. of 1.187, and a boiling point of 221°C .

The natural oil contains, in addition to methylsalicylic ether, a terpene, gaultherilene.

Theoretically, the artificial oil of wintergreen is identical with the natural oil in chemical composition; nevertheless it contains traces of alcohol derived from the component used, and difficult to eliminate. If such an oil be treated with a small quantity of spirit aniline red, the latter dissolves in the artificial oil and imparts a red coloration thereto, whereas it is left undissolved by the natural oil. At the end of about fifteen to thirty minutes the colour of the dyestuff undergoes a partial change to purple red in the artificial oil, but in the natural oil it is cherry red. In mixtures, the colour varies between these limits, according to the proportion of the natural and artificial oil present.

SAFROL.

Safrol distils from ethereal oil of sassafras between 230° and 236° C. It forms readily deliquescent monoclinic crystals, has an odour of sassafras, is insoluble in water, boils at 228° C., and has the sp. gr. 1.114.

According to Schimmel & Co., safrol is the cheapest scent for soaps that can be imagined, no other ethereal oil being similarly capable of removing or masking the disagreeable fatty smell of ordinary soaps. It may also be used in many ways for better classes of soap, advantageous and cheap compositions being practicable with this perfume and the oils of cassia, citronella, lavender, rosemary, spike lavender, thyme, and mint.

IONONE.

The creeping rhizome of the Florentine sword lily, the so-called violet root, exhibits a pungent but agreeable odour of violets. The crushed roots furnish on distillation a pale rose-coloured oil, which, at the ordinary temperature, deposits crystalline scales of nacreous lustre, and must therefore be regarded as violet-root camphor. This product is rarely met with in commerce, but there is now prepared by synthetic means a substance having the same odour as violet root, and the dilute alcoholic solutions of which are sold as ionone.

The odoriferous principle of the violet root cannot be obtained direct by steam distillation, a circumstance undoubtedly due to the high percentage of starch present, which seems to restrain the escape of the volatile constituents. For this reason Tieman and Krueger extract the roots repeatedly with ether, and subject the ethereal extract to distillation by steam. The result of these operations is the recovery of two products, one of which is a muddy residue, which, when treated with alkalis, furnishes myristic acid, or iridic acid, and a brittle crystalline resin. The distillate contains the volatile portions,

consisting of the ethereal oil, together with a larger quantity of myristic acid and its methyl salt, oleic acid and an ester of same, aldehydes, and other substances. The ethereal solution of this mixture is shaken up with a 2 per cent. solution of caustic potash, in order to eliminate a portion of the free organic acids. The residual organic acids and the esters are more easily removed by repeated fractional distillation of the oil by the aid of steam. The neutral oil obtained by this method is then dissolved in alcohol, and treated with an alcoholic solution of potash at the ordinary temperature, the resulting alkaline solution being then poured out into water, the aqueous solution extracted with ether, and the ethereal solution again subjected to distillation. The iris perfume passes over with the first portion of the distillate, and is then boiled with water and Ag_2O . This clarifies the irisketone by the formation of a phenylhydrazine compound, and, after leaving at rest for a day, the hydrazone mixture is distilled by steam. The irisketone remains behind, in combination with the phenylhydrazine, as a brown oil, which is then decomposed by dilute sulphuric acid into phenylhydrazine, and irisketone or ionone. This is next extracted from the aqueous portion by ether, and rectified under slight pressure. Ionone is an oil which is only sparingly soluble in water, but readily so in alcohol, ether, chloroform, benzol, and ligroin (light petroleum spirit), and has the sp. gr. 0.939. The odour of pure ionone is pungent, and in the concentrated condition not at all like that of violets; but when diluted with alcohol and exposed to the air, it immediately resembles the natural scent of the violet blossom.

Artificial oil of violets is prepared in the following manner, according to an American patent:—One kilo of acetone and $\frac{1}{2}$ kilo of citral are dissolved in $1\frac{1}{2}$ kilos of alcohol, and to the solution is added 1 litre of a freshly prepared clear saturated solution of calcium chloride. The mixture is kept stirred, and is heated for several hours under a reflux condenser, the result-

ing oil being afterwards boiled in 5 litres of a dilute solution of ferric chloride, and finally distilled. The citral or citraliferous oils may also be treated, in the acetone-alcohol mixture, with an efficient oxidising agent like barium peroxide, and then further oxidised by boiling with ferric chloride.

THYMOL.

Thymol is found in the ethereal oil from *Thymus vulgaris*, *Monarda punctata*, *Ptycholis Ajowan*, and is obtained by agitation with caustic soda and precipitation with hydrochloric acid. It forms colourless tables, has an odour of thyme, a pungent aromatic flavour, melts at 50° C., and boils at 232° C.

Thymol is only sparingly soluble in water, but readily so in alcohol and ether. Water can only retain 1 part of thymol per 1000 in permanent solution; but strong alcohol will take up its own weight of thymol. On the addition of water to the solution, the greater part of the thymol is deposited, turbidity being produced by water when the alcoholic solution is stronger than 1 per cent. One part of thymol is soluble in 120 parts of glycerin, and the solution still remains clear when diluted with water to 600 parts. Fats and oils, as also vaseline, are excellent solvents for thymol, but require warming. In the preparation of salves the thymol is first dissolved in a few drops of alcohol and then added to the fat.

A stable aqueous solution of thymol can be prepared by dissolving 1 gram each of thymol, tartaric acid, and pure dry caustic soda in a minimum of warm water, and diluting the solution with distilled water to 1 litre. This solution will keep clear for any length of time.

On adding a few drops of caustic potash and a little potassium iodide and iodine solution to a liquid containing thymol, so that the liquid assumes a faint yellowish brown colour, the application of a gentle heat causes the mixture to turn red, the colour

deepening by degrees. On strong heating, or prolonged standing, the coloration disappears and a precipitate is formed. By means of this reaction, as *little* as 0.05 mg. of thymol in 1 c.c. can be detected through the resulting red colour.

BISMUTH WHITE.

Bismuth white, also known as blanc de bismuth, blanc de perles, and pearl white, which is a basic nitrate of bismuth, and at the same time a constituent of several face cosmetics, is prepared by the action of fuming nitric acid on metallic bismuth. During this reaction a large volume of suffocating brown fumes are evolved, and this continues until the bismuth is completely dissolved. When the reaction has terminated, the solution is diluted with a large volume of water, the resulting precipitated powder being then collected on a filter and washed with water until the washings cease to redden blue litmus paper. The bismuth white is dried and stored in tightly closed vessels, since, on exposure to the air, it gradually assumes a yellow tinge from the absorption of sulphuretted hydrogen, for which substance it has a great affinity, and with which it forms a black compound, bismuth sulphide.

Basic bismuth nitrate is insoluble in water, but has a tendency to become altered in composition on prolonged digestion with a large volume of water, especially in the warm, inasmuch as it becomes more basic and the acid washing water absorbs bismuth. It is soluble in moderately dilute sulphuric acid, nitric acid, or hydrochloric acid, the resulting solutions being turned cloudy by the addition of water.

TIN OXIDE.

Tin oxide is obtained by treating metallic tin with fuming nitric acid and then levigating the product. It is a white insoluble powder, and is employed in cosmetics for polishing the finger nails.

CALCIUM CARBONATE.

Calcium carbonate, or levigated chalk, is prepared in the following manner :—One part of calcium chloride is dissolved in 15 parts of distilled water, filtered, and the filtrate treated with a previously filtered solution of soda crystals in distilled water, until a precipitate ceases to form. The resulting fine white deposit is precipitated chalk. When this has completely settled down, the supernatant liquid is poured off, and the precipitate, moistened with a little distilled water, is transferred to a paper filter in a funnel, and washed six or eight times with the wash bottle, after which it is dried by moderate warmth.

Calcium carbonate is almost entirely insoluble in water. According to Bineau, 1 litre of pure water can dissolve 0.016 gram of chalk ; Welzien, however, gives the quantity as 0.036. Hofmann 0.034, and Cruse 0.036 gram. When freshly precipitated, it is more soluble in water containing ammonium chloride, and still more so in water containing carbon dioxide. In the latter case, according to some, calcium bicarbonate is formed ; but, according to others, the monocarbonate is dissolved as a neutral salt, Lassaigne giving the amount as 1 part in 1136 parts of water at 10° C., and Warrington as 1 in 1015 parts of water, saturated with carbonate, at 21° C. Calcium carbonate is decomposed by acids, carbon dioxide being liberated.

ANIMAL SUBSTANCES USED IN THE PREPARATION OF
COSMETICS.

AMBERGRIS.

Ambergris is a greyish brown mass found floating in the tropic seas ; it occasionally contains remains of marine animals, and is regarded as a kind of gall stone or intestinal concretion

of the sperm whale. It is lighter than, and insoluble in, water, but soluble in alcohol, ether, oils, alkalis, or ammonia; softens in the hand, has an agreeable odour when warmed, and burns with a smoky flame. It contains 85 per cent. of fat, and, when distilled with water, gives about 13 percent. of a sweet-smelling oil.

CASTOREUM.

Castoreum is a secretion of the beaver, and is deposited in two pear-shaped sacs situated on the belly in both sexes. The beaver hunters separate these sacs from the carcase, and in this form the castoreum is brought to market. The sacs are about as long as the finger, and of the thickness of a man's thumb in the widest part; they contain a salve-like mass, which is yellow-brown, red-brown, or blackish, according to the food of the animal. It has a strong, disagreeable smell, a bitter balsamic taste, softens when warmed, is combustible, and is for the most part soluble in alcohol. Probably this secretion has some connection with the feeding of the beaver, which preferably consumes resinous plants when any selection is available.

In commerce a distinction is drawn between Canadian and Siberian castoreum; the latter is the more highly prized, and exhibits a peculiar odour of tar and Russia leather, probably due to the presence of certain constituents in the birch bark on which the Siberian beavers feed almost exclusively. American castoreum has an odour more recalling that of pine bark.

MUSK.

Musk is obtained from the musk deer, an animal about the size of a goat, and living after the manner of the chamois in the high peaks of the Himalayas. The male alone secretes musk, which is found stored in a kind of sac, or more properly gland.

Quickly-dried musk loses its odour almost entirely, and the same occurs when it is rubbed down along with *sulphur*

precipitatum, camphor, etc.; but the scent gradually returns when the musk is again moistened.

A good sac should contain 50–60 per cent. of musk, and no artificial seam should be apparent on the surface. Admixtures of blood are detected by a putrescent smell on moistening the musk with water. Added starch is revealed by the iodine test. Additions of metallic fragments, small stones, strips of leather, resin, etc., are often made, in order to increase the weight; the presence of these and similar objects can be detected by the naked eye or by the aid of a magnifying glass. In pure musk, the microscope reveals irregular whitish brown granules, along with cells, drops of oil, and generally also the mycelial threads of fungi.

Pure musk exhibits a highly characteristic behaviour towards caustic alkalies, such as soda, potash, or liquid ammonia, and these substances are therefore employed as tests for purity. On suffusing a sample of musk with a dilute solution of an alkali, a considerable intensification of the odour will become evident within a short time; if the alkali be concentrated or the sample warmed, the smell of musk disappears entirely, and the liquid evolves the pungent odour of pure ammonia. Hot water dissolves about 80 per cent. of the total weight of musk, strong alcohol about one half. When heated in an open porcelain basin, musk burns with liberation of a repulsive, pyreumatic smell, and leaves behind about 10 per cent. of ash.


To get rid of the smell of musk from a vessel in which that substance has been stored, it is sufficient to rub the inner surface over with bitter almonds moistened with water, or camphor moistened with alcohol. The smell may also be entirely removed by means of potassium permanganate.

A method for preparing musk by artificial means was devised by Bauer in 1889. He mixed toluol with the halogen compounds of butane, and boiled the mixture, with an addition of aluminium chloride or bromide, under a reflux condenser.

The product of the reaction is mixed with water and distilled with steam, the fractions passing over at 170-200° C. being collected and treated with fuming nitric and sulphuric acids. The resulting product is washed with water and recrystallised from alcohol, and furnishes yellowish white crystals, smelling strongly of musk; these, when dissolved in alcohol and treated with a trace of ammonia or ammonium carbonate, yield a liquid very similar to musk tincture.

According to the Valentiner process, a musk substitute can be obtained from turpentine. In this method, equivalent quantities of turpentine and isobutyl alcohol are mixed, and then gradually run into a five- to eightfold quantity of concentrated sulphuric acid, arrangements being made for cooling the whole. After one to two hours the mixture is poured into 5-10 times its volume of concentrated fuming nitric acid,—the necessary precautions being observed,—and then poured out into a large quantity of water. The nitro product is thrown down as pale yellow flakes, which are then collected on a filter and washed until neutral. The dried product forms a light yellow powder, with a strong smell of musk. The intermediate product from turpentine, isobutyl alcohol, and sulphuric acid, when thrown down with water, forms a thick whitish mass, which, when treated with steam, furnishes a trace of unattacked turpentine and alcohol, the remainder being non-volatile. It is soluble in all solvents except water; the product is probably of the nature of a dipentene. The final product, the pale yellow, nitrated, musky substance, is probably chiefly a dinitro product mixed with a little mononitro substance; it melts at about 70° C., and, on recooling, forms a brittle, pulverulent mass.

Later on—1892—Valentiner prepared a musk substitute which is soluble in both alcohol and water. This should open a wide field of usefulness to the product. The solubility in water is attained by nitrating a sulpho acid of butylxylol.



COSMETICS

Isobutyl alcohol and acetoxylol (B.P. 137° C.) are mixed in equivalent proportions, and gradually run into 5 parts of concentrated sulphuric acid (66° Be.), the whole being kept cool in order to prevent the evolution of sulphurous acid. After a short rest, the mixture is diluted with about a fourfold quantity of water, to separate the still unattacked condensation product; the latter collects as an oily layer on the surface of the liquid, and is removed when the separation is complete. The underlying clear, aqueous, rose-red liquid is treated with a saturated solution of common salt, in order to recover the para-isobutylxylol sulphonic acid, until the contents of the flask become in a short time interspersed with white crystals. These are collected by filtration, purified by recrystallisation, pressed, and dried at 100° C.

Another artificial musk is prepared by Link and Avenarius. Isodibutylene and halogen acids are heated with toluol, or with m-xylol, ethylbenzol, and similar substances, in presence of ferric chloride or bromide, whereupon tertiary butyltolylmethane, polymeric propylenes, as well as isopropyltoluol and homologues of this substance, are formed. The hydrocarbons are distilled with steam, dried, and fractionated, in order to separate the tertiary butylpropylmethane and the hydrocarbon $C_{15}H_{24}$ or its homologues. These products are then transformed into the corresponding trinitro derivatives, by heating them with nitric acid of sp. gr. 1.525 and fuming sulphuric acid, on the water bath. By removing the free acid and recrystallising from alcohol, these nitro compounds are obtained in the form of pale yellow crystals, exhibiting a musky odour.

CIVET.

Civet is yielded by a civet cat, a carnivorous animal allied to the ordinary cat and the marten, and indigenous to Asia and Africa; also by the musk rat. It is secreted in a double gland, found in both sexes.

Fresh civet is a whitish yellow mass, of the consistence of butter or fat, but turning thick and brown on exposure to the air. Like musk, it is characterised by a very powerful odour, which, as in the case of musk, becomes agreeable when largely diluted, and is used both alone and for the purpose of fixing other perfumes.

CARMINE.

Cochineal or carmine consists of the dried bodies of the female cochineal insect; and the commercial article takes the form of rounded, shrunken granules ranging in size up to that of a pea. The colour of the granules depends on the means employed for killing and drying the insects. If killed by hot water, the cochineal loses its ordinary white encrusting film, and has a reddish brown appearance. Drying in linen bags in a stove furnishes grey cochineal; and, finally, black cochineal is obtained by drying on hot stove plates. The colouring principle in cochineal is carminic acid, which is purple brown, and can easily be rubbed down to a red powder. It is very soluble in water or alcohol, but scarcely so if at all in ether. Caustic alkalis change the colour of the aqueous solution to purple red, and lead acetate produces a purple precipitate.

Cochineal is adulterated in a variety of ways, spent cochineal, according to Prior, being dried and mixed with fresh. In order to restore the original colour to spent cochineal, it is dried in drums containing heavy spar, talc, clay, gypsum, or white lead, whereby the white silky coating is imitated. Suspected samples of this character may be tested by treating them with water, whereupon the mineral matter is readily loosened from the cochineal, and can be separated by decantation for further examination. The presence of these mineral adjuncts can also be detected in the ash. Artificial granules, resembling true cochineal in form and colour, can also be discovered by the water treatment, since they crumble down in water, whereas true cochineal retains its shape. Adulteration with the colour-

ing matter of redwood can easily be revealed by the aid of lime-water, a dilute aqueous extract being thereby entirely decolorised in the case of pure cochineal, but retaining a violet blue tinge when redwood extract is present. The value of cochineal may also be determined volumetrically, by rubbing down the preparation to fine powder, mixing this with a white substance, and comparing the amount of the latter necessary to produce the shade of colour furnished by a standard cochineal treated in the same way with a known quantity of white.

The best method, however, is by a sample dyeing. On the other hand, methods based on the destruction of the colour in an extract, by oxidising agents like chlorine, etc., furnish only dubious results, since these reagents produce similar changes, not only in the colouring matter, but also in other constituents present in the aqueous extract.

Carmine is a red dyestuff, sold in lumps, powder, or dissolved in isinglass or egg albumin, and is usually prepared by precipitating a decoction of cochineal with an acid or an acid salt. By reason of its high price, it is frequently subjected to adulteration; and it has been found to contain admixtures of starch, clay, brickdust, etc. As pure carmine is completely soluble in ammonia, whereas these adulterants are insoluble therein, ammonia forms an excellent reagent for their detection.

To test a liquid for the presence of carmine, it is shaken up with 1 vol. of chloroform and 3 vols. of absolute alcohol; on then adding 2 vols. of distilled water to the mixture, without shaking, the carmine will separate out, almost in its entirety, at the plane of contact between the two layers of liquid. In the case of a fuchsine solution of equal strength, the whole of the colouring matter will be found in the layer of chloroform. Other colouring matters, however, are deposited at the plane of contact in this test, though very few pass completely into the

chloroform. The colouring matters in most general use behave in the following manner:—

Elderberry. The bulk of the colouring matter remains in solution, imparting a rose-red coloration to the upper layer of liquid, and a faint yellow to the chloroform mixture. Both layers are turned green by the addition of ammonia.

Campeachy wood extract. A violet deposit is formed, the upper layer being coloured. On the addition of ammonia the upper layer turns red, the lower one violet.

Red rose. The colouring matter is yellowish red, and gives a complete blue-violet precipitate.

Ratanhia extract. A very small brown deposit.

Red currant berries. The colouring matter is deposited, almost completely, as a rose-red precipitate.

Cochineal. A violet ring is formed, accompanied by partial deposition of the colouring matter.

Red wine. A rose-red ring is formed; an addition of ammonia turns the upper layer dirty yellow.

Raspberry. A rose-red ring on addition of ammonia.

Madder red. A red deposit; the chloroform mixture turns yellow.

Alkanna red. The colouring matter passes into the chloroform layer, and is turned blue by ammonia.

Santal wood. The colouring matter forms a yellowish red solution in the chloroform mixture. Ammonia decolorises the upper layer, whilst the chloroform layer is perfectly colourless.

CHAPTER IV

OILS AND FATS USED IN THE PREPARATION OF COSMETICS.

A JOWAN OIL, from the fruit of the umbellifer *Ptycholis Ajowan*, indigenous to India, is distinguished for its high percentage of thymol, for the preparation of which it is employed. The oil contains 30-40 per cent. of thymol, 30-40 per cent. of thymene, 15-20 per cent. of cymene, and a little carvacrol.

BAY OIL.

Bay oil, from the foliage of the West Indian myrtle, is a thin, yellow-brown, sweet-smelling oil, consisting of a terpene with 30-40 per cent. of eugenol. It is largely used in America for the preparation of bay rum.

BERGAMOT OIL.

Bergamot oil is largely prepared in Italy, and especially Sicily, by pressing the rind of the fruit of *Citrus bergamia*. It is a very fluid ethereal oil, of pale yellow colour, the poorer qualities being often greenish or brownish, but obtained perfectly clear and colourless by distillation with water. The odour is highly agreeable, resembling a mixture of the oils of orange and lemon, and the sp. gr. is 0·87-0·89. On prolonged standing it deposits white crystalline tables of stearoptone, which do not melt below 206° C. It solidifies a few degrees below the freezing point of water.

The most highly prized quality is Messina oil. It differs

from the other oils of the orange tribe by its ready solubility to a clear solution in caustic potash; but it shares with them the property of igniting with iodine, and being incapable of dissolving santaline, the red resinous colouring matter of santal wood.

The purity of bergamot oil can be tested by treating it with alcohol, whereupon it turns pale grey and gives a firmly adherent deposit, which collects in flakes when shaken up. After two days, the solid sediment is inappreciable in quantity, and is difficult to disseminate in flakes in the clear yellow liquid when shaken. The oil being often adulterated with alcohol, the following method for its examination has been recommended:—Fifteen parts of the bergamot oil are mixed with an equal volume of olive oil, or oil of sweet almonds; if alcohol be present, it will separate from the fatty oil like water; if, however, the mixture remain unaltered, one may be certain that the sample has not been adulterated with alcohol. The tannin test also gives reliable results.

BIRCH-TAR OIL.

Birch-tar oil is the liquid portion of the tar furnished by the dry distillation of birch wood, and consists of a mixture of numerous hydrocarbons. It is used as Russia leather oil.

OIL OF BITTER ALMONDS.

The ethereal oil of bitter almonds is obtained by distilling with water the crushed bitter almonds previously freed from their fatty oil by pressure. It chiefly consists of benzaldehyde, but also contains small quantities of benzoic acid, benzoin, etc., and is very poisonous, owing to the presence of 2–5 per cent. of hydrocyanic acid, from which, however, it can be freed by agitation with a solution of ferrous sulphate and slaked lime. The originally yellow oil can be entirely decolorised by distillation over quicklime. It is a thin, highly refractive oil, with a

peculiar, agreeable odour, and a pungent aromatic flavour. The sp. gr. at 15° C. is 1·043, the boiling point is 180° C. ; it is soluble in 30 parts of water, and dissolves readily in alcohol and ether. On exposure to the air it is oxidised into benzoic acid.

The usual adulterants are : nitrobenzol, alcohol, and light ethereal oils. These, however, have already been dealt with.

CAMPHOR.

Camphor is the customary name for a special form of camphor—laurel camphor—which is derived from the tree *Camphora officinarum*, and is the only kind appearing in European markets. It is shipped in the crude state from the producing countries, China and Japan, and is refined in Europe and America. The refined camphor is the only kind met with in the retail trade, though in different degrees of purity. Chinese camphor comes from Formosa, where the camphor trees cover the mountain chain extending from north to south ; the eastern provinces of Central China also furnish camphor. Japan produces about the same quantity as Formosa.

Crude camphor is generally of a greyish colour, the Japanese product, which is usually of higher quality, having a reddish tinge. The impurities amount to 6–10 per cent., though the better Japanese grades have only about 2 per cent. Camphor oil is never absent, and the same applies to a little incom-bustible matter. Bad qualities contain wood splinters. Among the impurities are also gypsum, salt, and sulphur, all of which are easy of detection. All crude camphor naturally also contains water.

Pure camphor melts at 175° C., and boils at 204° C., but also sublimes at the ordinary temperature, as is proved by the crystals deposited on the walls of vessels in which camphor is stored. The crystals are hexagonal, and take the form of tables and prisms when crystallisation proceeds in an unre-

stricted manner. It is readily dissolved by the solvents of resins, and also in concentrated acetic acid, but is almost insoluble in water, one part of camphor requiring over 1000 parts of water for its solution (1300 parts at 20° C.). The alcoholic solution deviates the plane of polarised light strongly to the right. The crystals exhibit double refraction. At 12° C. the sp. gr. is 0.995, but at 6° C. (and under) the density is almost exactly the same as water, so that camphor crystals hardly sink at all in very cold water. A remarkable property of pure camphor is that of describing circular movements on the surface of water, this peculiarity being absent in camphor oil or impure camphor. Unless moistened with alcohol or ether solvent, camphor is only triturable with great difficulty. The flavour of pure camphor is aromatic and rather bitter, warming at first, but cooling afterwards. It is easily ignited, and burns with a smoky flame.

The following method of preparing camphor powder has been patented by B. Smith of New York :—The camphor is dissolved in petroleum ether or benzine, then left to settle, the solution poured off, filtered if necessary, and the solvent expelled by distillation, until the camphor begins to crystallise from the hot solution, whereupon the distillation is suspended and the solution is stirred until cold. The more the crystallisation is interrupted, the finer the powder. The camphor powder prepared in this manner does not ball.

Grated camphor powder is better than that obtained by rubbing down camphor moistened with a solvent, inasmuch as it does not ball together, and is uniformly loose though coarse in texture. It, however, fulfils its purpose, since it can be easily rubbed down to the finest powder when required, as well as being suitable for rapidly preparing solutions of camphor in oil, ether, spirit, etc.

Up to the present no adulterants have been detected in camphor.

LEMON OIL.

Lemon oil is obtained by pressure or distillation from the skins of *Citrus medica* and *Citrus limonum*, and is a thin ethereal oil, colourless or yellowish, with an agreeable odour and taste. The sp. gr. is 0·840–0·870. Old oil has an acid reaction, sharp smell, and sharp, burning flavour. Pressed oil is of a yellowish colour, rarely quite clear, and deposits a thick sediment on standing; the distilled oil is colourless and clear, but of less agreeable taste and smell. The two are mixed together for sale.

Lemon oil is miscible with alcohol in all proportions, and dissolves, to a not perfectly clear solution, in two parts of alcohol of sp. gr. 0·815, or ten parts of alcohol of sp. gr. 0·830. It ignites with iodine. In commerce it is frequently adulterated with turpentine, the presence of which can be detected by the smell when a few drops of adulterated oil are rubbed between the hands. One drop of lemon oil, rubbed with sugar and added to 500 grams of water, should impart to the latter the pure odour of lemons. No alcohol should distil over when the oil is heated to boiling in a retort.

LEMON-GRASS OR CITRONELLA OIL.

Citronella oil, an ethereal oil prepared from lemon grass in Ceylon, is fairly thin, greenish yellow to brown, and has an odour like that of narcissus oil.

Pure unadulterated citronella oil should dissolve to a clear solution in 2 parts of 80 per cent. alcohol, which it will not do if adulterated with turpentine.

EUCALYPTUS OIL.

Eucalyptus oil is an ethereal oil extracted from the leaves of various eucalypti. According to Merck, there are two entirely separate kinds of eucalyptus oil: (1) *Oleum eucalypti australe*,

which is a comparatively cheap oil, used for technical purposes ; (2) *Oleum eucalypti*, from the leaves of *Eucalyptus globulus*, an oil for medicinal purposes, and much dearer than the first-named kind. The means of distinguishing between the two kinds consist—in addition to the smell, which in the case of the oil from *E. globulus* is agreeable and like that of the rose, whereas the cheaper oil smells more like turpentine—in the following characteristic indications :—(1) The *Oleum eucalypti australe* is strongly levo-rotatory, whilst the true oil and its preparations are either slightly dextro-rotatory or optically inactive. (2) Whereas *Oleum eucalypti australe* is only very slightly soluble in 90 per cent. alcohol, and gives merely a turbid solution in the proportions 1 : 15, the oil from *E. globulus* and the preparations from same (eucalyptol) dissolve with this alcohol in all proportions—even 1 : 1—to a perfectly clear solution. The Australian oil is also soluble in absolute alcohol. *Oleum eucalypti australe* ignites when treated with iodine, but the true oil does not. (4) The former assumes a red colour, the latter a yellowish tinge, when exposed to the action of sodium. (5) The sp. gr. of *Oleum eucalyptus australe* is above 0·860–0·870, whereas the density of the true oil varies according as it has been prepared from old or fresh leaves, but is in any case higher than that of the other, namely, 0·900–0·925.

OIL OF LAVENDER.

Oil of lavender is prepared in the south of France from the wild *Lavendula officinalis*, by distillation, in large quantities. It is a thin, colourless to yellow oil, miscible in all proportions with 90 per cent. alcohol, and having a density of 0·876–0·901. The best French oil of lavender, prepared solely from selected flowers, is sold as “Essence de Lavande Mont Blanc,” and is distinguished from the other qualities—in the preparation of which a larger or smaller proportion of stalks and leaves accompany the blossoms—by its exceedingly agreeable odour.

The chief adulterants of oil of lavender are alcohol, turpentine, and oil of spike lavender. The purity of the article is tested by mixing one drop with 10 c.c. of warm water and smelling the mixture, which should exhibit a pure and agreeable odour of lavender. The flavour of the mixture is momentarily bitter and aromatic. One volume of the oil should give a turbid solution with an equal quantity of dilute alcohol (sp. gr. 0.896), but a perfectly clear mixture with 3 vols. of same. On shaking up $\frac{1}{2}$ c.c. of the pure oil with a few granules of rosaniline, the liquid will remain colourless; but if alcohol be present a red coloration is developed.

Oil of spike lavender is distilled in the south of France from the flowers and leaves of *Lavendula spica*. It is colourless or yellow, and resembles oil of rosemary rather than lavender in smell.

Oil of spike lavender should mix to a clear liquid with its own volume of 90 per cent. alcohol, the contrary indicating the presence of turpentine as an adulterant.

PEPPERMINT OIL.

Oil of peppermint is an ethereal oil, obtained by distilling the flowers and other portions of the plant, *Mentha piperita*. The oil from the leaves is of superior odour and flavour to that from the whole plant. The three chief commercial kinds of this oil are the English, German, and American, the first named being the best, and that from Mitcham (Surrey) the best of all. The best marks of American oil are those of H. G. Hotchkiss, L. B. Hotchkiss, and Hale & Parshall.

Pure peppermint oil of good quality is neutral, thin, water-white to pale yellow, occasionally with a greenish sheen; exhibits a pure, penetrating odour of peppermint, and a burning, bitter taste, leaving behind a feeling of coldness on the tongue. It thickens with age, and acquires an acid reaction. The sp. gr. is 0.900-0.920. The oil gives clear

mixtures with equal and larger volumes of 90 per cent. alcohol; also a turbid mixture with its own volume of benzine, but a fairly clear mixture with 9-10 vols. Benzol gives turbid mixtures; both those with ether, chloroform, and amyl alcohol are clear.

The following adulterants are added to peppermint oil: fatty oils, alcohol, turpentine, copaiba oil, oil of mustard, and ginger oil. On heating over the water bath a piece of blotting paper moistened with one drop of the oil, the greasy mark will disappear entirely if the oil be pure, and will cease to be apparent even by transmitted light. When fatty matters are present, the grease mark, though seemingly absent when viewed by reflected light, will always be visible with transmitted light. When the oil contains more than 3 per cent. of alcohol, it is sufficient to let 1-2 drops fall into water, whereupon the drops will either be immediately surrounded by a milky zone, or will soon become whitish or cloudy in contact with the water. The most frequent adulteration is an addition of turpentine, which may then be recognised by the fact that, whilst pure peppermint oil gives a clear mixture with an equal volume of 90 per cent. alcohol, this is not the case when turpentine is present. To test for copaiba oil, 5 drops of the oil are treated with 15-20 drops of fuming nitric acid, shaken up, and set aside for one to two hours. At the end of this time the oily portion must still be liquid and not have solidified at all. To test for traces of mustard oil, 10 drops of the oil are placed in a wide test glass, along with 3-4 c.c. of absolute alcohol, 2-3 drops of silver nitrate solution, and 10-15 drops of ammonia. The mixture should remain clear and colourless, even when heated to boiling point; whereas if mustard oil be present, turbidity and blackening ensue on the application of heat, owing to the formation of liver of sulphur. After boiling, the solution is left to stand for two to three hours; if, at the end of that time, reduction, expressed by a greyish turbidity, has occurred, the sample is adulterated with another ethereal oil.

OIL OF CLOVES.

The volatile oil of cloves is obtained by distillation with superheated steam, or by extraction. It is a clear, colourless, yellowish or yellow-brown oil (reddish brown when aged), somewhat thick, of sp. gr. 1.030–1.065; it exhibits a strong smell of cloves and a similar burning taste. It frequently has a faintly acid reaction, boils at 250° C., and is readily soluble in alcohol or ether.

The adulterants of clove oil are: alcohol, oil of cinnamon, sassafras oil, oil of thyme, and paraffin oil. According to Hager, they may be detected in the following manner:—1. The sp. gr. and alcohol test consists in letting fall a few—5--drops of oil of cloves on to a column (about 10 c.c.) of 5 per cent. brine in a test tube. If the drops sink to the bottom of the glass, on gentle agitation, without becoming turbid or surrounded by a turbid cloud, it may be concluded that no alcohol is present, and that the sample has the right sp. gr. On closing the test glass with the thumb and slowly turning it upside down twice, so that the oil drops are compelled to traverse the column of water again, the drops usually cease to retain their limpidity. 2. Fifteen drops of the oil are mixed with 15 c.c. of water, shaken up, warmed to about 50° C., shaken up again and filtered. The filtrate should hardly exhibit any acid reaction indicative of salicylic acid or oil of cinnamon. 3. The same filtrate is cooled down and passed twice through a paper filter, to make it perfectly clear. One half is then treated with 2–3 drops of ferric chloride solution, whereupon a blue or green coloration indicates carbolic acid or salicylic acid. The milky turbidity produced by ferric chloride in the filtrate is pale violet, bluish, yellowish, or pale greenish, in the case of pure oil, but dark green to blue in presence of carbolic acid or phenol. Should alcohol be also present as an adulterant, a lighter coloration may be expected. The other portion of the

same filtrate should, when treated with half its volume of sulphuric acid, furnish a raspberry-coloured liquid. 4. Half a c.c. of the oil should give a clear mixture with an equal or larger quantity of alcohol, turbidity indicating the presence of paraffin oil. If the clear mixture with 10 vols. of alcohol be divided, and the one half—3-4 c.c.—be treated with several drops of sulphuric acid, the other with 1-2 drops of ferric chloride solution, a blue or violet-blue coloration should be developed in both cases. In the ferric chloride test, old clove oil will generally give a green mixture. 5. Carbon disulphide should give a very turbid mixture. 6. Pure oil of cloves should give a clear mixture with 2-3 vols. of 60 per cent. alcohol. In presence of castor oil and other fatty oils, paraffin oil, sassafras oil, or cinnamon oil, a turbid mixture would ensue. 7. On mixing 50 drops of the oil with 1 drop of water, warming up to 40-50° C. and shaking, a clear mixture will be obtained when the oil contains crystallised phenol. In such event the oil gives a fairly clear mixture with a fivefold volume of benzine. Should anhydrous phenol be present, the mixture will be white, turbid, and non-transparent, whereas pure oil gives an opalescent but still translucent mixture. When, therefore, the oil gives with benzine a nearly clear, or a non-transparent, whitish, turbid mixture, it may also be suspected as adulterated.

In testing ethereal oils, the results obtained by Schimmel & Co. of Leipzig in this connection are worthy of attention. The solubility of these oils in alcohol furnishes the main criterion of their value, and of adulteration. According to the determinations made by the above-named firm, 1 part by volume of—

Aniseed oil	dissolves in	1½-2	parts of	90 per cent. alcohol.
Bergamot oil	„	¼-½	„	90 „
Cajeput oil	„	3-5	„	70 „
Calamus oil	„	all proportions of 90 „		

	dissolves in	1-2	parts of	80 per cent. alcohol.
Cassia oil		1-2		
Citronella oil	"	1-2	"	80
Lemon oil	"	5	"	90
Eucalyptus oil	"	3-5	"	70
Fennel oil	"	5-8	"	80
Geranium oil	"	2-3	"	70
Curly mint oil	"	equal	"	90
Caraway oil	"	3-10	"	80
"	"	equal	"	90
Lavender oil	"	3	"	70
Mace oil	"	3	"	90
Clove oil	"	2	"	70
Orange flower oil	"	1½-2	"	80
Palmarosa oil	"	3	"	70
Patchouli oil	"	equal	"	90
Peppermint oil				
(American)	"	"	"	90
Peppermint oil (Eng- land and Japan)	"	3-5	"	70
Rose oil	"	equal	"	90
" (liquid portion)	"	"	"	70
Rosemary oil	"	½	"	90
Santal wood oil	"	5	"	70
" (East Indian)	"	3	"	90
" "	"	1-2	"	80
Thyme oil	"	15-30	"	70
Wintergreen oil	"	5	"	70
Ylang-ylang oil	"	1-2	"	90

PATCHOULI OIL.

Patchouli oil is an ethereal oil obtained from the leaves and twigs of *Pogostimon patchouli* in the East Indies. It is yellowish or brownish yellow, somewhat thick, and about the same density as water, or a little heavier. It is levo-rotatory, boils at 282-294° C., and on prolonged standing deposits an isomeric patchouli camphor in prismatic crystals, melting at 54-56° C., of sp. gr. 1.045, and with a peculiar, penetrating and persistent odour. The commercial varieties comprise a so-

called natural patchouli, prepared in the East Indies, a rectified oil, and a so-called crystallised patchouli oil, rich in patchouli camphor. The odour is the same in all three, but that of the latter kind is specially persistent.

Patchouli oil is frequently adulterated with cedar-wood oil, which may be detected by the circumstance that the pure oil is readily soluble in its own volume of 90 per cent. alcohol, whereas the adulterated oil is not.

JUNIPER OIL.

Juniper oil is the volatile oil obtained by steam distillation from the fruit of *Juniperus communis*. It is colourless, of sp. gr. 0·840–0·860; that obtained by ordinary distillation being yellowish, and of sp. gr. 0·85–0·9. The odour is that of juniper berries, and the oil has a burning taste. It has a great tendency to resinify, and then becomes thick and acid. The adulterants are juniper wood oil and turpentine.

Juniper oil is soluble in alcohol, and gives a clear mixture with carbon disulphide without thickening. No disagreeable taste should be imparted to water when 3 drops of this oil are rubbed down with sugar and shaken up with 500 grams of water. According to Hager, pure juniper oil mixed with 30 vols. of 90 per cent. alcohol gives a somewhat cloudy but still transparent solution, which, when kept in a warm place, becomes lighter, but still remains turbid. With 3–4 vols. of absolute alcohol, of 2–3 vols. of petroleum ether, the oil gives a turbid mixture, but a clear one with chloroform. Adulteration with alcohol is detected by the addition of a little rosaniline, which is indifferent towards juniper oil, but soluble in alcohol.

YLANG-YLANG OIL.

Ylang-ylang oil is an ethereal oil from *Yona odoratissima*, a plant indigenous to the Philippine Islands, and is shipped from Manilla to the European market as a water-white or faintly

yellowish liquid. It commences to boil at 160° C., but the boiling point extends to 300° C. The sp. gr. is 0.980. Ylang-ylang oil is a mixture of several oils.

CINNAMON OIL.

Cinnamon cassia oil is the oil recovered, in China and Cochin China, by distilling the rind, unripe fruit, calices, and other waste portions of the cinnamon cassia tree. It has a pale yellow colour, which browns with age, is thick, and has a sweet flavour, with burning after taste. The sp. gr. is 1.03–1.09. One part of the pure oil is soluble in 2 parts of 80 per cent. alcohol.

The quality of the oil is revealed by the taste, smell (especially when warmed), and by the high sp. gr., which causes the oil to sink in water. According to Hager, cinnamon cassia oil is frequently adulterated with clove oil; but Deite considers this improbable, since clove oil is usually the dearer of the two. On the other hand, it is frequently adulterated with cheaper thick ethereal oils, especially with cedar-wood oil, in which case the solubility differs from the rate given above.

Cinnamon oil, the ethereal oil obtained from the waste of Ceylon cinnamon by distillation with salt water, is golden yellow when fresh, afterwards turning reddish brown; it has a more agreeable odour and taste of cinnamon than cinnamon cassia oil. The sp. gr. is 1.005–1.03.

This oil is adulterated with the cheaper cassia oil, which is very difficult to detect, except by connoisseurs expert in the flavour and odour of the true oil.

OIL OF ALMONDS.

This is the fatty oil of sweet and bitter almonds. According to Deite, the commercial oil is mostly expressed from small, cheap bitter almonds and apricot kernels. There are two chief kinds, English and French, the latter being the cheaper, but of

poorer quality. The oil is thin, clear, and of faint yellow colour, with a mild, agreeable flavour but hardly any odour. The sp. gr. is 0.915–0.920 at 15° C. It solidifies at –20° C., is miscible in all proportions with ether, benzine, and chloroform, and is soluble in 15 parts of absolute alcohol at boiling heat, but requires 60 parts at the ordinary temperature.

Almond oil is adulterated with poppy oil, sesame oil, earth-nut oil, apricot-kernel oil, beech oil, and walnut oil. In addition to the smell and taste, the following three tests may, according to Deite, be employed for detecting these adulterations:—

1. *The elaidin test.*—Four to 5 c.c. of the oil are placed in a test glass, 1.2 cm. in diameter, along with 2–3 c.c. of nitric acid (sp. gr. 1.185), and well shaken up. A cloudy, whitish or yellowish white oleaginous layer is formed, from which a portion of the nitric acid separates out as a turbid liquid. A couple of copper turnings are then placed in the glass, and the bottom is warmed until bubbles begin to ascend from the copper; the glass is next set in a place kept at a temperature of 17–20° C., and watched at intervals of five minutes during the first half-hour and every fifteen minutes afterwards. The oily layer must always exhibit a cloudy whiteness, and never be yellow, transparent, or almost clear, nor have a yellow, red, brownish, or brown appearance. After four to five hours the oil layer will be set and white—oil of sweet almonds,—or merely thick and cloudy white as at first—oil of bitter almonds, apricot-kernel oil. From eight to twenty-four hours this layer, regarded by transmitted light, shows up as a solidified stratum of white crystalline granules above the blue acid layer, and is in turn surmounted by a thick somewhat turbid oily layer, with a number of white granules deposited on the walls of the glass. The surface of the liquid layer also frequently contains a deposit of the same granules, and occasionally the whole of the oily stratum is interspersed and thickened with the granules. If the entire oil layer is liquid, and free from the granules in question, or if, instead of being

whitish, the layer is almost perfectly clear, or yellow, reddish, or brownish ; or, again, if the layer containing the white granules is covered by a liquid, clear or reddish, yellow or brownish, oil layer, then the oil is not pure almond oil. At the end of thirty hours—the limit of the test—the crystalline granules begin to assume a coloured appearance.

2. *The ammonia test.*—This consists in mixing 1 vol. of 10 per cent. ammonia with 3 vols. of the oil. Pure almond oil gives a white to yellowish white mixture, which becomes and remains liquid after half an hour. If brownish, or coloured a strong yellow, or if nearly solid at the end of half an hour, the sample is an adulterated oil.

3. *The sulphuric ether test.*—This test is performed by mixing 2 c.c. of the oil with 4 c.c. of ether, and then pouring down the side of the tube about 1·5 c.c. of concentrated sulphuric acid, so that it collects at the bottom of the mixture. Meanwhile the acid should not be coloured more than yellow. On then shaking gently, so as to mix the upper portion of the acid layer and the lower part of the oil and ether mixture, no dark coloration should ensue, even when the shaking is continued far enough to saturate the acid. The latter may become somewhat darker in shade, but not dark coloured. In the case of old samples of oil, the mixture of oil and acid, without ether, may become dark in colour, but not when ether is present. Most of the adulterating oils give dark-coloured mixtures under the ether test.

ROSEMARY OIL.

Rosemary oil is the volatile oil obtained from the leaves of *Rosmarinus officinalis* ; it is colourless, or somewhat yellowish, thin, usually with an acid reaction, with a penetrating odour like camphor, an aromatic, bitter, cooling taste, and of sp. gr. 0·880–0·915. On exposure to the air, it resinifies and thickens, a stearoptone being occasionally deposited. Rosemary oil is

soluble in 1-1½ vols. of 90 per cent. alcohol, and gives clear mixtures with ether, carbon disulphide, amyl alcohol, benzol, and benzine. The Italian oil is usually considered the best, the French being next: the German oil is inferior in quality, and generally takes 2 or 3 vols. of 90 per cent. alcohol to dissolve it.

Pure rosemary oil gives a clear mixture with 1-1½ vols. of 90 per cent. alcohol, does not ignite with iodine, and sinks in a mixture of 10 c.c. of 90 per cent. alcohol and 2 c.c. of water. It does not dissolve fuchsine. When heated on a glass slide, it evaporates completely, evolving a pure odour of rosemary.

OLIVE OIL.

Olive oil, the finer grades of which are also known as Provence oil, is pressed out of the fruit of the olive tree. It commences to become cloudy at 4-5° C., attains the consistence of butter at 2° C., and solidifies at a little below zero C. A distinction is drawn between edible oil and oil for manufacturing purposes. The finest edible oil is the so-called virgin oil, which is obtained by cold-pressing selected fruit. The manufacturing oils are divided into burning oil, second runnings, and sulphur oil, the last-named being recovered from the press residues by extraction with carbon disulphide.

Deite states that olive oil is largely adulterated, especially with cotton-seed oil, which is extremely difficult to detect. The hardening of the oil is utilised as a means of detecting this adulterant, the oil being frozen by means of a refrigerating mixture, and, after remaining two hours in this condition, exposed to the pressure of an iron block, which is loaded with weights until it sinks into the solid oil. From the weight required to produce this result, it is possible to tell whether the oil is pure or adulterated with cotton-seed oil, and even whether the latter is present in larger or smaller quantity.

Thus, for example, a load of 1700 grams is required to produce penetration in the case of pure olive oil, less than 1000 grams will suffice with the commoner grades; whilst only 25 grams are required for cotton-seed oil, and with mixtures of the two oils the weight lies between these extremes.

Poppy oil is also used to adulterate olive oil, but may be detected by the rise in temperature ensuing when the oil is treated with sulphuric acid. On mixing 10 c.c. of 66° Be. sulphuric acid with 50 grams of pure olive oil, the temperature of the mixture will rise 42° C. in three to five minutes, whereas in the case of poppy oil the accession of temperature amounts to 86°·4 C. According to the observations of Fehling, the increase of temperature, under these conditions, in mixtures of olive oil and poppy oil, is in direct and uniform proportion to the amount of poppy oil present. Thus, with

10 per cent. of poppy oil, the mean increase of temperature is 40°·5 C.					
20	“	“	“	“	44° “
50	“	“	“	“	55° “
80	“	“	“	“	64° “

Additions of earth-nut oil, when in any considerable amount, are easily detected by the bean-like taste peculiar to this oil, and also by the sandy granules which it deposits at a temperature of 8° C., whereas pure olive first begins to deposit solid particles at 4° C.

In order to impart a green colour to olive oil, it is boiled with fresh spinach until all the moisture has been driven off, whereupon 1 per cent. of sodium carbonate is added in order to partly nullify the influence of light on the chlorophyll. Rancid olive oil is rendered edible by treating it with 3 per cent. of finely sifted calcined magnesia, and mixing the whole, for a quarter of an hour, four or five times daily for a whole week. At the end of this time it is filtered, and, if necessary, washed with warm water.

WAX.

Hager has recommended a very useful method for testing the purity of wax.

A small cylindrical piece of wax, 5-6 mm. thick and 3-4 cm. long, is placed in a test-tube, 8-10 cm. long and 1.2-1.4 cm. wide, where it is suffused with sufficient benzine to cover it to a depth of 1-2 cm. If the wax be pure, small dusty particles will separate from the surface of the cylinder, and at the end of one and a half to two hours the whole will be resolved into a powdery mass with level surface. In the case of yellow wax this may take three hours, the wax being at the same time bleached by contact with the benzine, the latter remaining colourless, or assuming a yellowish coloration. Adulterated yellow wax generally retains its colour, besides staining the benzine yellow. In contrast to the pure wax, adulterated cylinders remain unaltered for half a day, sometimes as much as two to four days, and then do not fall down to powder, but split up into longitudinal rods, which are either straight or bent.

If the adulterants do not amount to more than a small percentage, flakes separate from the wax cylinder; at the end of half a day or so, the sediment consists of flakes interspersed with broken longitudinal columns. Even when the proportion of adulterant is small, indications pointing to the formation of columnar fragments are present. Hager calls this the benzinate test; it is best applied at a medium temperature—14-18 °C. He applied it to all forms of adulteration in wax, and always obtained satisfactory results. Similar behaviour to benzine is exhibited by ethyl ether of sp. gr. 0.720.

SPERMACETI.

Spermaceti is a whitish, rather solid, translucent substance, of nacreous lustre, pulverulent but not brittle, with a slippery

but scarcely greasy feel, and exhibiting a foliaceous crystalline fracture, a somewhat insipid taste, and very faint peculiar odour. The sp. gr. is 0·94–0·95; melting point, 45–50° C. It does not leave any greasy mark on paper; is soluble in 7 parts of hot and 35 parts of cold 98 per cent. alcohol; readily soluble in ether, chloroform, or carbon disulphide; and also soluble in benzol, especially when warmed. On prolonged exposure to the air, spermaceti turns yellowish and rancid; on ignition it burns with a brilliant inodorous flame.

Adulterations of spermaceti with stearin are easily revealed by the altered fracture. In such cases, stearic acid can be extracted by boiling with dilute solutions of sodium carbonate, whereas spermaceti melts and floats on the surface of the liquid. On dissolving the sample in 30 parts of benzol, cooling the solution, and leaving it to stand for three days, pure spermaceti will give a perfectly clear liquid; if, however, small crystals are deposited in the course of an hour, the sample is adulterated with paraffin. If the liquid is turbid to milky, the presence of stearic acid is indicated.

LARD.

Hog's lard consists of olein, palmitin, and stearin, and contains 0·23 per cent. of unsaponifiable matter. On exposure to the air, it quickly becomes yellow and rancid, in which condition it readily takes up copper and lead from vessels containing those metals, and therefore becomes poisonous. To test for these metals, the lard is digested with acetic acid, and a portion of the latter is then treated with potassium ferrocyanide, whereupon a brown coloration indicates copper, whilst lead is revealed by a white precipitate produced on treating another part of the acid extract with dilute sulphuric acid. Lard is soluble in 27 parts of absolute alcohol, or 35 parts of alcohol of sp. gr. 0·9128; also in ether, chloroform, carbon disulphide, hot benzine, and warm petroleum ether.

When fresh, it has a neutral or faintly acid reaction, and furnishes, on saponification and subsequent decomposition of the soap, 8·80 per cent. of glycerin and 95·90 per cent. of stearic acid, palmitic acid, and oleic acid. It is white in colour, granular, and of the consistence of ointment; the odour is faint and peculiar, and the taste agreeable, rather sweet, and fatty. It flows between the fingers; and when melted and recooled exhibits a crystalline structure.

One of the most frequent methods of adulterating lard is by additions of water in order to increase the weight. This can be easily detected by melting a sample and leaving it to stand in a lukewarm spot, whereupon the water will separate out. The weight is also artificially increased by adding solutions of potash, soda, or caustic soda, which at the same time whiten the colour of the lard; also by adding chalk, alum, clay, common salt, or starch. 2 to 5 per cent. of alum, or 1 per cent. of lime, will suffice to mask the presence of 12-15 per cent. of water. Tallow and other fats are also employed to adulterate lard.

To detect the presence of chalk, clay, or alum, a portion of the suspected lard, about 100 grams, is melted with 1-1½ times its own weight of water at 60-70° C., stirred frequently, and set aside for a few hours, whereupon any chalk or clay will settle down. Common salt will be found dissolved in the water, and can be thrown down, as a white precipitate, by an acid solution of silver nitrate. Alum is detected by the copious precipitate furnished with ammonia, or by the white precipitate following the addition of barium sulphide; soda and potash, by the white precipitate with calcium chloride; and potash also, by the yellow coloration given with mercury chloride.

Admixtures of starch or flour give a blue coloration with iodine solution. Tallow is occasionally employed as an adulterant, and is revealed by microscopical examination,

since its crystals (deposited from an ethereal solution) are in the shape of plums or curved like the letter S, whereas those of lard are rhomboidal plates, with sloping angular corners. An approximate estimation of the percentage may be arrived at from the number of the crystals of each kind observed.

The detection of extraneous fats as adulterants in lard, entails a very thorough examination. The chief adulterant of this class is refined cotton-seed oil, mixed with corresponding amounts of beef tallow or stearin. Of course it is not impossible that other oils may be used for the same purpose when the price permits. Simple reactions for detecting such additions have been proposed from time to time, and among them an alcoholic solution of silver nitrate. Though pure lard will stand this test, whereas the commercial qualities do not, one is not justified on this ground alone in characterising the latter as adulterated, since the presence of even a very minute amount of extraneous substances will suffice to colour the reagent brown, and such substances are invariably present in commercial brands of lard. The same applies to the molybdic acid test, which is too delicate for the majority of the commercial grades. It is performed as follows:—One gram of the lard is dissolved in about 5 c.c. of chloroform, and treated with 2–3 c.c. of a solution of sodium phosphomolybdate. After shaking and leaving at rest for a few minutes, the two liquids will be found to have separated again, and the reagent solution above the chloroform will have assumed a more or less intense green to greenish blue coloration, according to the amount of reducing substances present in the fat. The same exactly applies to the Hirschsohn reaction for the detection of cotton-seed oil, which, so it is claimed, will reveal as low as 5 per cent. of that oil. Here 5 grams of the sample are treated with 5–6 drops of a solution of 1 gram of gold chloride in 200 grams of chloroform, then well shaken up, and immersed for twenty minutes in boiling water. When cotton-seed oil is present, the mixture assumes a

coloration, red or rose by reflected light and greenish by transmitted light.

Lead acetate has also been recommended as a reagent, and is said to produce a more or less orange-coloured tinge when shaken up with the melted fat, after the latter has previously been agitated with ammonia. All these reactions, however, are uncertain, and the only way to form a decision is by a thorough examination, to determine—

1. The microscopic appearance of the fat recrystallised after melting.

2. The melting point.

3. The saponification value.

4. The melting and solidification points of the fatty acids.

5. The iodine value.

With regard to the examination of lard, Mansfield says that the difficulty in testing the purity of lard resides not merely in the divergent, and for the most part complex, adulterations, but also in the great lack of concordance in the data published in the literature.

The ordinary examination extends, in addition to the tests for gross admixtures—water, mineral matters—to an investigation of the purity of the sample. The principal tests for this purpose are the determination of the iodine value and the refractometer index. In the case of ordinary commercial lard, the former does not exhibit such fluctuations as are given in the literature of the subject. The fat gives the iodine value 59–63, and the refractometer index 50–51° of the Zeiss butyro-refractometer at 10 C. Divergences from these figures should cause the sample to be suspected. Lower values indicate an admixture of beef tallow, which may be confirmed by the recrystallisation test, benzol being the best solvent for this purpose.

The best means of detecting vegetable oils are the tests of Becchi and Welmanns, which, however, have to be performed

with care, since even pure fats may give slight reactions. It is essential that the fat should be filtered clean. Moreover, since an abnormal iodine value may be given in the case of a complex adulteration, these qualitative reactions should always be performed. In such cases a decision may be obtained by means of a closer examination, which should furnish the following constants:—

Melting point, 38–40° C.

Solidification point, 35–38° C.

Iodine value, 63–64.

Refractometer index at 40° C., 34–36.

Nevertheless, for calculating the percentage of admixture, it is always necessary to isolate the liquid fatty acids. For this purpose the following modification, used in the laboratory of the Paris Bourse, is employed:—

Ten grams of the solid fatty acids are dissolved in 100 c.c. of ether, and shaken up well for some time with 3 grams of zinc oxide. This gives zinc soaps, from which those containing oleic acid can be extracted by ether. The solution is filtered in an aspirator, the ether is expelled by distillation, and the residue decomposed with 20 c.c. of hot dilute hydrochloric acid,—1 : 4,—then washed twice with hot water in a separating funnel, filtered and dried.

Pure lard gives the following values:—

Iodine value of the liquid fatty acids, about 92.

Refractometer index, 44–45.

Both these values are increased to a considerable extent in the case of mixtures containing vegetable oils, since the latter contain linolic acid.

The melting point of the solid fatty acids may be determined by the usual methods. A capillary tube is filled with the melted fat, and, after having cooled down and set, the tube is attached to the tube of a sensitive thermometer. The two are then immersed in a water bath, which is slowly heated until

the fat melts, the temperature at which this occurs being read off on the thermometer scale. To increase the accuracy of the determination, the beaker containing the tubes and water may be placed in an outer vessel, and heat applied to the latter.

The saponification value is determined by the Köttstorfer method: $1\frac{1}{2}$ –2 grams of the fat are treated with 25 c.c. of half-normal alcoholic caustic potash on the water bath for fifteen minutes: as soon as saponification occurs, 1 c.c. of an alcoholic solution of phenolphthalein is added, and the liquid is titrated with half-normal sodium chloride solution. A blank experiment is made by titrating 25 c.c. of the alcoholic caustic potash by itself. The difference between the number of milligrams of caustic potash used and the amount found on back titration is calculated to 1 gram of fat, the result being the saponification value.

For determining the iodine value, Hübl employs an alcoholic mixture of iodine and mercuric chloride: 25 grams of iodine are dissolved in $1\frac{1}{2}$ litres of 95 per cent. alcohol (free from fusel oil), and a similar quantity is used to dissolve 30 grams of mercuric chloride. After standing twelve hours, the solution is ready for use, but its standard must be determined before each test. For the iodine value determination, 0.2–0.4 gram of a liquid fat, or 0.8–1.0 gram of a solid fat, must be weighed out and dissolved in chloroform. It is then treated in succession with 20 c.c. of the iodine solution, and further additions of 5–10 c.c. each, until the solution has acquired a dark brown colour at the end of two hours, whereupon 10–15 c.c. of a 10 per cent. aqueous solution of potassium iodide and 150 c.c. of water are added. The free iodine is then titrated with a solution of sodium thiosulphate containing 24 grams per litre, and the absorbed iodine is referred to percentage units of the fat. The resulting value is known as the iodine value or Hübl value. The figures seem to be constant, or approximately so,

for each oil or class of oils, and are highest in the case of the drying vegetable oils. The Hübl method has proved very useful for the detection of cotton-seed oil in tallow and lard, pure cotton-seed oil having the iodine value 109·1, pure tallow 40·8, tallow with 5 per cent. of cotton-seed oil 44, with 10 per cent. 47·1, with 15 per cent. 49·7, with 20 per cent. 52·9, with 25 per cent. 56·1, with 30 per cent. 59·2, with 40 per cent. 66·2. In the case of lard, the detection is hindered by the presence of beef stearin, pure lard having the iodine value 57-63, whilst that of beef stearin is 23-28.

PARAFFIN.

Paraffin is entirely without taste or smell, bluish white in colour, translucent, with a clear ring and bright vitreous lustre. The melting point varies considerably according to the hardness of the paraffin, the limits being 30-60° C.

The commercial grade known as hard paraffin melts between 55 and 60° C., but softens at a lower temperature, and has a greasy feel even at the ordinary summer temperature. The sp. gr. varies, according to the hardness or melting point, between 0·870 and 0·912. Paraffin is insoluble in water, but is soluble in various proportions of ether, alcohol, chloroform, carbon disulphide, turpentine, and especially in light hydrocarbons, from which it crystallises out in plates. It is not attacked by concentrated acids and alkalis at the ordinary temperature. It boils above 300° C., and decomposes on distillation. It is soluble in nearly all volatile and fatty oils, and is miscible in all proportions with stearic acid, wax, fats, and resins. Solid paraffin melting at 56° C. expands by about one-seventh of its bulk, and exhibits the property, discovered by Albrecht, of expanding to a similar extent when dissolved in oils, and thus converted into the liquid condition, at the ordinary temperature, the solution being lighter than either of its components.

STYRAX.

Styrax belongs to the balsams, and is obtained from a small tree, *Liquidambar orientale*, by subjecting the bark to pressure. It forms a thick turpentine-like mass, grey in colour, of a sharp burning taste, agreeable odour, and soluble in strong alcohol: the solution exhibits a very agreeable aroma when highly diluted, but not before. Styrax possesses the valuable property of fixing a number of very delicate perfumes and reducing their volatility.

A substance not to be confounded with Oriental styrax is that from American sources, put on the market as white Peruvian balsam or liquidambar, and derived from *Liquidambar styraciflua*. This is quite transparent, pale yellow in colour, and more fluid than the Asiatic styrax, which it is sometimes used to replace, though of a less delicate odour.

PERUVIAN BALSAM.

Peruvian balsam is brought to market from Central and South America. It is derived from *Myroxylon peruiferum*, and is obtained by making incisions in the bark and stem, from which the balsam then exudes. It is of syrupy consistence, viscous, and the colour is brownish red when viewed in thin layers, or blackish brown in thick layers. The taste is bitter and burning, afterwards irritating; the odour is similar to vanilla, but less delicate. Peruvian balsam is completely soluble in alcohol, chloroform, and amyl alcohol; partly so in ether, benzol, petroleum ether, turpentine, fatty oils, and caustic alkalis.

Peruvian balsam is frequently adulterated with alcohol, copaiba balsam, castor oil, and styrax. The detection of the adulterants is attended with difficulty; but, on the other hand, the characteristics of Peruvian balsam are so decided that one can easily determine whether a sample is adulterated or not.

The chief criteria are the density and solubility. The former—1·14–1·16—is usually checked by means of a solution of common salt, containing 1 part of dry sodium chloride to 5 parts of water, and having the sp. gr. 1·125. If a drop of Peruvian balsam, introduced into this solution, sinks, the sample may be considered pure; in the contrary event, it is certainly adulterated, all the adulterants of Peruvian balsam having a relatively lower density, namely, alcohol, copaiba balsam (0·95), castor oil (0·96), turpentine (0·87).

NUTMEGS.

Nutmegs are nuts, almost globular in shape, about as large as a walnut, greyish brown in colour externally, and covered with a reticulated whitish grey coating, the inside being reddish brown with white marbled patches.

Good fresh nutmegs should be close, heavy, and at the same time so rich in oil that the latter exudes on the nutmeg being pierced with a needle. Light nutmegs, which are hollow and worm-eaten, besides being poor in aroma, are unsuitable.

BENZOIN.

Benzoin, a soft resin, is derived from a tree of the Styracinee family, indigenous to tropical Asia. As a rule, fine benzoin forms nodules of extremely divergent size, light grey in colour, and interspersed by white almond-like lumps—almond benzoin. This kind of benzoin, which, from its origin, is also known as Siam benzoin, is mostly in small lumps—tears—which are translucent, pale yellow to brown externally, but milkwhite in fracture, and exhibiting a very powerful odour of vanilla. Less fine, but still highly appreciated, is the variety known as Siam benzoin, in masses consisting of large red-brown lumps, enclosing white portions—almonds. Inferior in quality, but highly prized, is the true Sumatra almond benzoin, which comes to market in the form of lumps as large as the

fist. Calcutta benzoin is in the form of large friable blocks of a dirty reddish brown colour. Penang or Sumatra benzoin invariably contains cinnamic acid as well as benzoic acid; this kind usually comes on the market as chocolate-brown masses, containing numerous rather large crystals; it is packed in drums. The inferior grades of benzoin consist of brownish fragments without any white patches, but frequently contaminated to a large extent with splinters of wood, bast fibres, remains of leaves, etc., and suitable only for cheap preparations.

In addition to the foregoing properties, fine benzoin must also exhibit a sweetish, sharp burning taste, be readily pulverulent, and when heated in a porcelain basin, liberate, in addition to the irritating vapours of benzoic acid, a very agreeable perfume. It should also be almost completely soluble in strong alcohol.

Benzoin melts at 65–80° C., is readily soluble in alcohol and ether, and partly so in chloroform, amyl alcohol, benzol, petroleum ether, sodium carbonate solution, and caustic alkalies. It is insoluble in fatty oils, but soluble in cold acetic acid.

VASELINE.

Vaseline is the final product of the distillation of petroleum residue. It is yellowish, yellow, or pure white in colour, of the consistence of ointment or lard, is devoid of taste or smell, translucent in thin layers, and possesses the property of not becoming rancid. Thanks to this property, vaseline enjoys a fairly extensive employment in connection with the care of the skin; although, on the other hand, its incapacity of absorption by the skin—in which respect it differs from the animal and vegetable fats—would not seem to render it specially suitable for that purpose.

The consistence of various kinds of vaseline is not always the same, but ranges from that of goose fat to the firmness of butter; accordingly, the melting point also varies, ranging

between 30° and 45° C. Vaseline is frequently mixed with spermaceti, paraffin, or ceresin, in order to make it firmer; and artificial vaselines are also known.

Vaseline is insoluble in water and glycerin, but miscible with the latter when the same is free from water; it is only very slightly soluble in 90 per cent. alcohol, but is soluble in 150 parts of absolute alcohol. It is more or less soluble in ether and chloroform, according to circumstances, more readily so in benzol, and most easily of all in carbon disulphide. With ethereal and fatty oils it is miscible in all proportions. It dissolves bromine and iodine at the ordinary temperature, but sulphur not below 120 – 130° C. Furthermore, it exhibits the peculiar property that when heated for about fifteen hours to 110° C., it will absorb oxygen, whereby it acquires an acid reaction and smell.

In order to test vaseline for the frequently observed addition of vegetable and animal fats, La Reole and Dupin propose to rub down a given quantity—5 grams—of vaseline with 5 drops of a solution of potassium permanganate, in a mortar. In the case of pure vaseline, the rose-red coloration of the reagent will be retained, but will be instantly converted into a chestnut brown when extraneous fats are present; and the amount of fat thus added as an adulterant can be determined from the depth of the colour.

Michle has proposed the following test for the purity of vaseline:—Two grams of vaseline are dissolved in 5 grams of chloroform, and shaken up energetically with 10 c.c. of water, a single drop of phenolphthalein solution being then added. In the case of pure vaseline no colour change is produced, but a rose-red is formed on adding a single drop of decinormal caustic potash, thus indicating freedom from acid and alkali. † Another test by the same author is to heat 10 grams of vaseline along with 10 grams of water on the water bath for a quarter of an hour, with stirring. After cooling, the water is poured off, and

should be perfectly neutral, giving no sulphuric acid reaction with barium chloride, and leaving merely an imponderable residue when evaporated to dryness on a watch-glass.

Vaseline is browned when heated on the water bath with 98 per cent. sulphuric acid, though weaker—73 per cent.—acid has no action, except when an imperfectly refined sample—technical vaseline—is present. The best method of performing this test is to melt 10 grams of the vaseline in the water bath, and add 50 drops of 73 per cent. sulphuric acid, the whole being then heated, with stirring, for another quarter of an hour. If the vaseline be pure, the acid will exhibit scarcely any alteration, no dark ring being formed at the plane of contact, nor is the acid coloured brown. Again, if 5 grams of vaseline be heated with 5 grams of sodium carbonate and 25 c.c. of water, for half an hour on the water bath and stirred, the aqueous layer being poured off when cold, and then treated with sufficient hydrochloric acid to produce supersaturation, the liquid will remain clear, provided neither resins nor fatty acids are present.

Vaselone, a substitute for vaseline, is a solution of stearone and margarone in a neutral mineral oil. The stearone is obtained by distilling 75 parts of stearic acid with 25 parts of dry powdered lime, in an iron cylinder or retort, at a temperature of 400° C. The product melts at 86° C. The margarone is prepared in the same manner, by distilling beef fat with lime. The vaselone is prepared by dissolving 15 parts of margarone and 5 parts of stearone in 100 parts of pure, inodorous mineral oil. The finished product resembles vaseline, except that it is not transparent. It is white, inodorous, neutral, and is not attacked by acids or other chemical reagents.

LANOLIN.

Lanolin is purified wool fat. It melts at 40° C., and can absorb up to 105 per cent. of water without losing its salve-like

consistence. It is employed as a basis for ointments, an adjunct to cosmetic soaps, etc. etc., being difficult to saponify, and therefore not liable to become rancid. A. von Rod removes the colouring matter and smell of crude wool fat by treating it with oxidising agents, such as bleaching powder, or permanganate, in alkaline solution. The wool fat is extracted by strong alcohol, benzine, sulphuric ether, benzol, ether, and chloroform.

The following method for testing lanolin has been recommended by Astolfi, on the basis of comparative colometric determinations, and can also be extended to form a quantitative examination. He treats 0.1 gram of lanolin with 10 c.c. of concentrated sulphuric acid, the result being an intense red liquid with greenish fluorescence. The colour decreases in intensity in proportion as the wool fat is adulterated with substances other than cholesterin. This solution is then treated with 10 c.c. of chloroform, and shaken up energetically. On the separation of the liquid into two layers, the chloroform will be found coloured a brilliant red if the wool fat be pure, but will be less red if the sample contain adulterations of water, extraneous fats, and the like. Furthermore, when the lanolin is pure, a black ring will be discernible at the plane of contact, but is less decided when the sample is adulterated. By comparing these results with those furnished by a sample of known purity under the same treatment, the extent of the adulteration, if any, can be estimated.

CHAPTER V.

GENERAL COSMETIC PREPARATIONS.

GENERAL cosmetic preparations cover a rather wide field, inasmuch as they comprise not only such as are concerned with the care of the skin, but also the hair, teeth, buccal cavity, and callosities.

Many cosmetics have been developed into specialties, or are capable of being so developed with profit; these are dealt with in a separate section.

In this case, as in the case of special preparations, the rule should be followed of using only such materials as are perfectly innocuous to the health and well-being of the consumer. Apart from general considerations of humanity and morality, the manufacturer should not forget his liability to prosecution in the event of injury arising from the presence of noxious ingredients in his preparations. The position is further complicated by the circumstance that vanity, and the desire to conceal personal imperfections, are so dominant in some individuals, as to render them careless whether the articles they use are dangerous to health. This is a factor that should be prominently borne in mind, especially in recommending preparations for special purposes; and the straightforward manufacturer of cosmetics will discard any ingredients that are either useless for the purpose in view, or likely to prove injurious under prolonged use.

True, as regards the efficacy of many preparations, the position of the manufacturer is often a very unpleasant one. Even

though he may personally feel a conviction that preparations for promoting the growth of the hair and beard, for example, are not able to do what is claimed for them, it would be acting contrary to the interests of his business were he to condemn such preparations in general. Purchasers of these articles are not dissuaded by such a course, and if they do not obtain the wished for preparation at one seller's they resort to another. This necessity for supplying an existing want leads to the production of preparations which, even if harmless, are nearly or entirely valueless.

This explains why space has been accorded in the present work to certain preparations of an inferior character, the object of the book being to describe the manufacture of cosmetics on a technical scale, and not to serve as a guide to the laity.

Before, however, proceeding to details of preparation, the author will devote a little space to the general considerations underlying the proper application of cosmetics.

The care of the skin has been greatly facilitated by nature, the numerous glands present in the cuticle all serving some useful purpose. The human skin is composed of three separate parts: the under layer or future skin, the derma or true skin, and the epidermis or upper skin, which is very thin, and consists for the most part of dead or dying dermal cells; it is continually casting off scales and being recruited from the true skin.

The skin contains various extraneous structures, such as the sweat glands, which eject their well-known secretion; the fat glands, whose function it is to keep the skin covered with a layer of fat, and thereby soft, shiny, and supple; and, finally, the so-called hair bulbs forming the seat of the hairs.

The chief task in the rational care of the skin is to keep the said glands constantly in good health and working order. This end is attained by the employment of various preparations,

which, in addition to maintaining the general well-being of the body, also impart bloom to the skin itself.

Personal beauty is really nothing more than the perfectly normal development of the face and body during complete health, and therefore the product of regular structure and normal function of the body. However, just as the health may be disturbed by the most divergent causes, so also can the skin, the seat of external beauty, be affected by alterations and diseases, which may exert such an injurious influence thereon as to unfavourably change the whole external appearance.

This is where the task of cosmetics comes into operation, to supply preparations suitable for the rational care and beautifying of the skin.

Pure glycerin is a substance having an extremely protective action on the skin, which it renders white, smooth, supple, and lustrous; in fact, sunburnt skin cannot be restored to its original light colour so quickly by any other means—except hydrogen peroxide—as by glycerin. The best method of using glycerin is as a wash, equal parts of viscous glycerin and orange-flower water—or some other aromatic water with a fine perfume—being mixed together and coloured rose-red by a very small quantity of fuchsine. Concentrated glycerin is unsuitable for use as a wash, because it absorbs moisture from the skin, and produces a sensation of burning or heat.

In such cases where the preparation is not necessarily liquid, but may be of the consistence of ointment, the glycerin may be advantageously replaced by vaseline or lanolin.

In addition to ordinary soap, the so-called emulsions, pastes, vegetable milks, and crèmes form excellent means for improving the skin; in fact, they are, in some respects, superior to soaps, since they contain substances which have not only a cleansing action, like soaps of all kinds, but also render the skin lighter, clearer, and softer.

It is a frequent practice to draw a distinction between two groups of emulsions, namely, emulsions consisting of masses which, when brought into contact with water, exhibit the property of emulsifying in that liquid; and true emulsions, which already contain the necessary water, and therefore exhibit the characteristic milky appearance. The difference between the two, however, is very slight, consisting merely in the larger or smaller content of water; so that they may all be dealt with under one heading.

The milkiness of emulsions on being brought into contact with water is due to the circumstance that they contain, in addition to fat, substances possessing the property of converting the fat into exceedingly fine drops, as a result of which the entire liquid acquires the peculiar milky appearance. As a glance through the microscope will show, cow's milk (and that of other animals) consists of a perfectly clear liquid containing the fat drops—butter globules—in suspension, the white colour of the milk being the result of the difference between the refractive property of these globules and that of the clear liquid.

Whereas soaps always contain a certain quantity of free alkali, the emulsions are almost if not quite destitute of this class of substance, and therefore, whilst sharing the cleansing properties of soap, are without its drawbacks. Consequently they are excellently adapted for the care of the skin, to which, when applied regularly, they impart a warm, fresh tint, as well as smoothness and delicacy.

Glycerin is a material of special value in the preparation of emulsions, not merely because of its property of softening the skin, but also on account of its capacity for dissolving colouring matters, so that it acts as a beautifier in the true sense of the word; skin that has been sunburnt ever so badly is turned quite white again in a surprisingly short space of time by the use of glycerin. In addition, glycerin retards the decomposi-

tion of preparations, and helps to retain them unchanged during prolonged storage.

The value of this latter property should on no account be underestimated, since all emulsions are particularly liable to decompose and turn rancid, owing to their content of finely divided fat. Complete exclusion of air, and storage at a low temperature, form the sole means, under ordinary circumstances, of retarding rancidity, which is accompanied by the disengagement of an unpleasant smell that cannot be masked by any perfume; but an addition of glycerin to any emulsion will at the same time render it more stable, thanks to the antiseptic action of the glycerin itself.

Another preservative which has long been known, and which is exceedingly powerful in this respect,—barely $\frac{1}{2}$ per cent. being sufficient for the purpose,—is salicylic acid; this substance preventing decomposition of any kind, rancidity among others. This acid, being harmless, and without either taste or smell, can be used wherever it is necessary to oppose the destructive influence exerted by air, ferments, etc., on different substances. Whereas it was formerly the practice to prepare emulsions in only small quantities at a time,—sufficient for a few weeks,—it is now possible, thanks to the preservative properties of salicylic acid, to make the emulsions on a large scale and store them for a long period, without fear of deterioration.

The only fats used for the preparation of emulsions are fatty oil of almonds, olive oil (both obtained by pressure), and lard. The almond oil can be advantageously prepared by pressing crushed almonds; the fresh almond shreds also find employment in cosmetics. The olive oil and lard must be purified in a most careful manner, by warming for an hour with about a tenfold volume of water, containing soap in the proportion of 1 part per 100 parts of the fat under treatment. The fat is afterwards washed 5 or 6 times with fresh quantities of clean warm

water, until the washings have a neutral reaction, and do not colour either red litmus paper blue, or blue litmus paper red—the one reaction indicating the presence of free alkali (soap), and the other that of free fatty acids, and therefore the existence of rancidity in the fat. Both of these would injure the quality of the product, more especially rancidity; and it is therefore essential that the fat should be perfectly neutral to test paper.

The so-called meals and pastes consist of the meal of vegetable substances containing fat, and exhibiting the property of forming an emulsion with water. They are frequently used as washes, and, being free from the smallest trace of alkali, form the most delicate of all such preparations, and are specially adapted for washing the face and delicate hands.

The various kinds of vegetable milk are nothing more than emulsions, already containing sufficient water to give them a milky appearance. They are employed direct, as washes. Owing to their larger proportion of water, they are more liable to decomposition than the preparations already mentioned, the fats being extremely susceptible to fermentation,—rancidity,—which condition is further favoured by the extreme fineness of division.

To improve the keeping properties of these preparations, an addition may be made of 5–10 per cent. of glycerin, which at the same time increases their cosmetic value; equally to be recommended is an addition of $\frac{1}{2}$ per cent. of salicylic acid, which will prolong the stability indefinitely. Each of the vegetable milks is compounded on the basis of a mixture of soap, wax, spermaceti, and some aromatic water, on which latter the specific name of the finished preparation usually depends. This composition is for the purpose of maintaining in suspension the fatty vegetable substance—almond meal, pistachio meal, etc.—which is the cause of the milky appearance of the liquid.

The crèmes, whilst similar in constitution to the emulsions and vegetable milks, differ from these latter in their thicker consistence, which renders them suitable for rubbing into the skin. The crèmes are really salves, scented with some well-known perfume after which they are called. Fat is the chief component of these mixtures, and, being capable of imparting fulness and softness to the skin, is the cause of their hygienic influence. A properly prepared crème should have the consistence of freshly solidified wax, but give way, like pomatum, when pressed by the finger. The efficacy and beautiful transparent appearance of a crème can be improved by an addition of very thick glycerin; but this must be made with care, otherwise the mass will lack the proper consistence.

The preparation of cold cream entails the mixing of wax, spermaceti, and almond oil with an aromatic water and a perfume. The first stage is easy enough, the wax and spermaceti being simply melted together at the lowest possible temperature, and then incorporated with the almond oil by stirring. It is, however, more difficult to combine this basis with the rest of the components; the aromatic water is poured in, as a thin stream, with continued stirring, and it is only when the contents of the basin have become homogeneous that the perfume is added, this being done quickly, and the still fluid mass poured at once into the vessel in which it is to be kept. Since the introduction of Chinese gelatine, however, the preparation of cold cream has been greatly facilitated, wax, spermaceti, or paraffin being entirely dispensed with and replaced by Chinese gelatine, made up into a solution that will give the required consistence when it sets, and containing a small quantity of salicylic acid to prevent rancidity. This gives a cold cream resembling marble or alabaster. These preparations are usually packed in decorative urns or porcelain vases, closed with ground stoppers or tinfoil-covered corks to prevent rancidity. The perfume must always be added last of all,

and not until the mass has already been cooled down to solidification point, since, if added earlier, there is certain to be a good deal lost by evaporation.

The hair requires a certain care and preservation. It may be laid down as a general rule that the use of sharp combs and brushes for the hair should be entirely avoided, elastic combs and soft brushes alone being suitable for the purpose. The use of ordinary water for damping the hair is also to be deprecated, more particularly because hair moistened in this way soon disengages an objectionable smell. Furthermore, the immoderate plastering of the hair with fats and oils should be avoided, the natural fat in the hair itself being usually sufficient to keep it smooth; and even when this is not the case, as frequently happens, all that is necessary is a corresponding supplementary addition of oily matter. In such event the hair should be frequently cleansed with a lukewarm wash to remove the fat, which has a tendency to clog thereon along with dust and dirt. It is impossible to warn too strongly against a too frequent use of cold water,—such as washing the hair daily with cold water,—since this practice will inevitably result in baldness. In very many cases the employment of fat may be entirely superseded by moistening the hair and roots with glycerin twice a week. A suitable application of glycerin will also remove the troublesome scurf on the scalp. Beyond this, all meddling with the hair should be avoided by all who are desirous of preserving this growth. Dyeing the hair, even with the most harmless materials, and singeing the hair, are both highly injurious. The growth and well-being of the hair are furthered by the use of a pervious head covering, and by leaving the head bare in the hot weather.

The therapeutics of baldness—which appears to be continually on the increase—is not yet complete, notwithstanding the extensive and comprehensive experiments that have been

made of late years. In many cases of baldness among females, it is not improbable that the cause may be traced to a neurotic condition, whilst in other instances everything points to a parasitic origin of the disease. To this class belongs undoubtedly the typical Alopecia areata, which has been definitely proved to be transmittable from one individual to another; consequently, the best treatment of this disease is of an anti-parasitic character. The important treatment introduced by Lassar must be regarded as of this class: daily washing the hair for several minutes with coal-tar soap—or with sodium carbonate, 15·0; potassium carbonate, 15·0; medicinal soap, 70·0; rose water up to 200 parts,—followed by rinsing and drying. Following this treatment, the hair is rubbed with a 0·2 per cent. solution of corrosive sublimate, then with absolute alcohol containing $\frac{1}{2}$ –1 per cent. of naphthol, and finally with 2 per cent. salicylol. In addition, Lassar recommends first rubbing the roots of the hair with a paste composed of β -naphthol, 10·0; flowers of sulphur, 40·0; yellow vaseline, 25·0; green soap, 25·0 parts. Then, after five to thirty minutes, 25 per cent. chrysarobin-lanolin is rubbed in with a pointed brush, the skin of the scalp being afterwards impregnated with tar oil, which is washed off with coal-tar soap. The head is then washed with the above solution of sublimate and naphthol-alcohol, and oiled with 2 per cent. salicylol or 2 per cent. salicyl-vaseline. A pomade composed of lanolin or horse fat, containing 2 per cent. of turpentine, is applied at night. Lassar does not hold a very high opinion on the utility of galvanic treatment, iodine tincture, or formalin, though he does not deny they may occasionally be of service. Perhaps, also, the Finsen light treatment may have a favourable effect on this complaint.

Mention should also be made of Jackson's recommendations for the care of the hair, these being particularly *à propos* in

cases where the hair has a tendency to fall out, as it often does after fevers. The scalp should be rubbed, once or twice a week, with a pomatum containing 1 gram of precipitated sulphur and 30 of cold cream, the hair being washed every two to three weeks, and oiled as soon as dry. Washing the hair too often is injurious. The best soaps to use for this purpose are the liquid kinds, *e.g.* an alcoholic solution of Sapo viridis. It is important that the hair should be oiled after every washing, in order to prevent it getting too dry. When the hair is naturally poor in fat, it should be treated, at least once a week, with a hair oil that does not easily turn rancid. The hair may be cut as often as desired, but should not on any account be singed.

The basis of every hair pomade is some fat or other, which is scented with an agreeable perfume, and occasionally coloured. The fats most in use for this purpose are lard, beef marrow, tallow, olive oil, and almond oil, to which is added—of course only in the case of solid fats—a certain quantity of paraffin, spermaceti, or wax, or better still Chinese gelatine, in order to improve the consistence of the pomade. Complete purity of the fat is essential, only absolutely neutral—acid-free—fat being permissible; whereas any that contains even a trace of rancidity—free fatty acid—is entirely precluded, on account of the penetrating smell appertaining to many of these acids.

Bersch gives the following recipe for preparing pure fats for this purpose. The fat is melted, with about a threefold quantity of water, in a bright iron, or better still, enamelled vessel, the water containing an addition of 1 per cent. of alum and 1 per cent. of common salt, both calculated on the weight of the fat. The mixture of fat and water is worked about with a broad, flat paddle, though better results are obtained from a stirring apparatus attached to the melting pan, and, after the mass has been left alone for some time, the fatty coagulum is skimmed off the surface. When a small

pump is available, the operation can be considerably accelerated by pumping the fat and water out of the pan, and allowing the mixture to return in the form of a fine spray.

When fat that is already partly affected with rancidity has to be purified for cosmetic purposes, the alum mentioned above should be replaced by a $\frac{1}{2}$ per cent. solution of caustic soda.

After the fat has been treated sufficiently with water, it must be washed, in order to completely remove the substances employed in its purification. Formerly this washing was performed in a manner resembling the grinding of oil colours, the fat being spread on a flat plate and rubbed with a muller, in a stream of water flowing from a vessel above the plate, until purified. At present this expensive process has been replaced by a mechanical operation, the fat being treated with water in a vertical mill.

It sometimes happens that, even when the greatest care has been bestowed on the purifying of the fat, the pomades prepared therefrom become rancid on prolonged storage, a circumstance that may seriously damage the reputation of the manufacturer. Fortunately, two substances are known to possess the property of retarding the rancidity of fats, namely, salicylic acid and benzoin. Consequently, both these substances are largely used as preservatives in the preparation of cosmetics, especially for pomades. An addition of 1-5 parts of solid salicylic acid per mil. will suffice to attain the desired object, while in the case of benzoin, about 20 grams are taken per litre of fat. This resin is almost completely soluble in fat, and imparts to the latter an odour of vanilla. For the finest pomades, use is made of sublimed benzoin in the proportion of 10-15 grams per litre of fat.

The care of the buccal cavity and teeth is a matter of enormous importance, and the preparations employed in this connection also belong to the domain of cosmetic technology.

It is essential that the mouth and teeth should be kept

perfectly clean, a result impossible of accomplishment without the use of suitable mouth-washes and tooth-powders. In order to effectually combat the numerous bacteria peopling the buccal cavity, as well as to prevent the decomposition of residual organic matters in the mouth and to destroy same, antiseptic mouth-washes must be employed. At the first glance, salicylic acid would appear to be most suitable for this purpose, both on account of its general harmlessness and its predominant antiseptic properties. Nevertheless, it cannot be recommended at all, its decalcifying action on the teeth precluding its use. An experience of many years has convinced the author that the best of all mouth-washes is an alcoholic solution of thymol, dropped into water. One gram of thymol is dissolved in 30 grams of rectified spirit and kept as a stock solution. For use, 10-15 drops are placed in a glassful of water, and the mixture is held in the mouth for a short time, and the mouth finally rinsed out with the same. It is highly advisable to repeat this treatment both in the morning, after each meal, and again at night before retiring to rest. Inveterate smokers desiring to free their breath from the taint of tobacco smoke, can cleanse the buccal cavity with a very dilute solution of potassium permanganate, a couple of crystals of this salt being dissolved in a small glass of water—to which it imparts a rose-coloured tint. With this solution, which does not brown the teeth, the mouth is rinsed out several times.

In connection with the care of the teeth, the greatest errors are met with in the choice of tooth powders. Such powders as contain any attritive substances, like pumice, and even charcoal, should be avoided. In their normal condition, human teeth are covered with a hard lustrous mass of vitreous character, the so-called enamel, which offers great resistance to the action of acids or putrefying substances. On the other hand, this enamel is very sensitive to sudden changes of

temperature, and is easily cracked, thus admitting acid or putrefactive substances to the interior of the teeth, and effecting their destruction. The true internal substance of the teeth consists of a porous mass of bone, which is easily destroyed; and when this takes place the whole of the teeth are completely spoilt. Tooth powder containing pumice acts on the teeth like a file, quickly wearing away the enamel and exposing the perishable internal substance. Charcoal, it is true, possesses valuable disinfectant properties, but this advantage soon disappears when it is remembered that charcoal always contains certain quantities of silica, *i.e.* a substance capable of scratching glass, and one that must inevitably wear away the enamel of the teeth if used continuously for cleaning them.

According to the author's long experience, the best tooth powder is one composed exclusively of levigated chalk, perfumed with a few drops of oil of peppermint. It is, however, necessary to see that the brush used is not too hard. This chalk powder, applied to the inside and outside of the teeth with a soft brush, will clean them thoroughly, and the peppermint oil is refreshing and enlivening.

Experiments on the value of mouth washes have been performed by Dr. Karl Roese. The subject of the experiments was fed in the morning on coffee and bread, the mouth being then rinsed for one minute with a sterile solution of common salt and peptone, at blood temperature. One tenth of a c.c. of this solution was then examined for the presence of micro-organisms, the result being that a single dose of the rinsing liquid was found to contain from 10 to 800 millions of bacteria. This first rinsing was followed by a rinsing with the mouth wash under examination. It was finally ascertained by the experimenter that the more healthy the teeth and the stronger the muscles of the mouth, the larger the proportion of bacteria gaining access to the stomach.

According to these experiments, the strongest antiseptic is a

mouth wash consisting of water containing benzoic acid and corrosive sublimate; this, however, being poisonous, is not suitable for daily use. Formaldehyde has no effect, a result to which its ready decomposibility largely contributes. Consequently, the preparation known as kosmin, the chief ingredient of which is formaldehyde, does not possess the value claimed for it.

Eau de Botot does neither good nor harm. Salicylic acid is injurious, owing to its decalcifying action on the teeth. A better preparation is the one known as Odol, on account of its content of salol.

An excellent, disinfecting, simple, and perfectly harmless mouth wash is afforded by 40 per cent. alcohol. This considerably widens the small blood vessels of the mucous membrane, and in this manner increases metabolism. This artificial circulation of the blood dissipates congestion in diseased gums, and restores the same to health.

The class of skin beautifying preparations comprises rouge, preparations for concealing discoloured portions of the skin (freckles, moles), that would otherwise spoil the complexion, and the toilet powders used to impart whiteness and smoothness to the skin.

To take the rouges first, the use of these preparations can only be recommended to a limited extent. For theatrical purposes, rouge is indispensable. In the first place, the rouge should not contain any ingredient injurious to the skin; and, secondly, any preparation of this kind must be removed as quickly as possible, because it chokes the pores of the skin.

The preparations employed to remove or conceal freckles and moles seem to be of a very dubious character. Here also it should again be emphasised that the preparations ought not to contain any ingredients of a dangerous character. With regard to others, the manufacturer of these cosmetics has to face the point already referred to, namely, the use of materials he feels to

be worthless ; nevertheless, he must comply with the demand, if his business is to prosper.

Toilet powders are mostly of an innocuous character, and are applied to the skin by means of a powder-puff or a hare's foot. For the most part they consist of some kind of starch and powdered steatite, suitably perfumed, and often coloured as well. The kind of starch used is immaterial : rice, wheat, and potato starch all serve the same purpose, provided they are sufficiently white and finely powdered. The better the powder is rubbed into the skin, the whiter does the latter become, and the less is the powder noticeable.

According to the author's experience, the best face powder for keeping the skin clean, especially in the case of persons who perspire freely, is salicylic acid dusting powder.

As an appendix, consideration may be devoted to hair dyes, depilatories, and preparations designed to impart strength and gloss to the hair.

So far as hair dyes are concerned, it should be remarked that the production of any given hair dye is often only a question of fashion. The same fundamental rule must be followed in this case as in those already discussed, namely, that no injurious materials should be employed. In using hair dyes, it is essential, especially in the case of such as consist of two portions, to first free the hair from grease, since the chemicals employed do not stick on an oily surface. In dyeing the hair, it is advisable to use a dilute solution first, and if this does not produce the desired shade, the operation should be repeated ; whereas it may happen that, by using the dye in a concentrated condition, the colour produced will not resemble that of any shade of natural hair. The complete removal of grease from the hair, previous to dyeing, may be effected either by washing the hair once or twice with soap, or with a dilute solution of soda. The hair must then be allowed to become thoroughly dry before applying the dye.

Depilatories are easily prepared, and are certain in their action, the substance of the hair being quickly destroyed. Of course, no injurious ingredients must be used. At the same time, it should be remembered that the effect of these preparations is, and can only be, the same as that of shaving, namely, to promote the aftergrowth of the hair treated; and a person who once commences to use depilatories is obliged to continue the practice at diminishing intervals. The only means yet known of effectually removing hair permanently is by electrical treatment, and the operation is too complicated and difficult to be successfully attempted except by a skilled medical man.

Wax pomades, stick pomades, or bandolines, serve to stiffen the hair. The active ingredients are wax, spermaceti, tragacanth, and gum arabic.

Brilliantines, or preparations for imparting gloss to the hair, consist chiefly of glycerin, suitably scented, and coloured light red or violet in order to improve the appearance. Since the aniline dyes are generally harmless, and at the same time soluble in glycerin, they are used for this purpose.

CHAPTER VI.

MOUTH WASHES AND TOOTH PASTES.

FORMALIN MOUTH WASH AND TOOTH PASTE. As a substitute for kosmin, the former is prepared in the following manner:—

Fifty grams of formalin—40 per cent. formaldehyde—are dissolved in 1000 grams of pure alcohol, then triturated with 200 grams of tincture of benzoin, 50 grams of myrrh tincture, 3 grams of peppermint oil, 2 grams of aniseed oil, 1 gram of oil of cinnamon, 15 drops of Ceylon cinnamon oil, and 2 grams of cochineal, the whole being then intimately mixed and filtered.

Formalin tooth paste is prepared by triturating 30 grams of formalin with 1000 grams of purest levigated chalk, then adding 200 grams of powdered violet root, 50 grams of magnesium carbonate, 100 grams of soap powder, 10 grams of peppermint oil, 2 grams of bergamot oil, and 1 gram of citronella oil, the whole being then mixed up to form a paste with about 700 grams of chemically pure glycerin.

Instead of 10 grams of peppermint oil, half that quantity may be taken, in conjunction with 3 grams of menthol.

AN ANTISEPTIC MOUTH WASH which is said to retain its properties indefinitely, has been described by Huchard. It consists of a series of antiseptic substances, any of which taken separately would suffice, namely—

Crystallised boric acid	.	.	.	4.0 grams.
Eucalyptol	.	.	.	1.0 gram.

Salol.	.	.	.	.	.	2·0 grams.
Menthol	.	.	.	.	.	0·25 gram.
Thymol	.	.	.	.	.	0·1 „
Alcohol	.	.	.	.	.	100·0 grams.

This can be scented according to taste, and coloured red with cochineal.

The same author has also given the following recipe for a
SALOL TOOTH POWDER:—

Salol	.	.	.	.	.	4·0 grams.
Calcium phosphate	.	.	.	.	.	20·0 „
Calcium carbonate	.	.	.	.	.	20·0 „
Magnesium carbonate.	.	.	.	.	.	20·0 „
Sodium bicarbonate	.	.	.	.	.	15·0 „

This is scented with peppermint oil.

Frohmann (a Berlin dentist) recommends the following soap for brushing the teeth:—Thymol, 0·25; ratanhia extract, 1·0 in 6·0 of hot glycerin; calcined magnesia, 0·5; sodium biborate, 4·0; medicinal soap sufficient to make the whole up to 30·0; peppermint oil, 1·0.

This soap should be used after dining, and again before retiring for the night. By altering the proportions of the sodium biborate and soap, the same recipe may be also made up as a tooth powder.

A stable aqueous solution of thymol is prepared by Hermite by adding caustic soda and tartaric acid to a 0·05 per cent. solution of thymol in water, this strength being quite sufficient for antiseptic purposes. One gram each of thymol, tartaric acid, and pure, dry caustic soda is dissolved in a minimum of warm water, and the solution diluted with distilled water to 2 litres. It will remain clear for any length of time.

A EUCALYPTUS TOOTH PASTE may be prepared from 160 grams of precipitated chalk,—levigated chalk,—45 grams of soap powder, 45 grams of wheat starch, 1 gram of carmine, 30 drops of peppermint oil, 30 drops of geranium oil, 60 drops of

eucalyptus oil, 12 drops of clove oil, and 12 drops of aniseed oil, mixed together and made into a paste with a mixture of glycerin and alcohol in equal parts.

EAU ANATHÉRINE.

Guaiacum wood	100 grams.
Myrrh	250 "
Cloves	150 "
Santal wood	150 "
Cinnamon	50 "
Alcohol	4 litres.
Rose water	2 "
Mace oil	5 grams.
Rose oil	5 "
Cinnamon oil	5 "

The solids are digested with the alcohol, the ethereal oils are dissolved in the filtered liquid, and the rose water is added last.

EAU DE BOTOT.

This tooth tincture is prepared in different ways, the most highly appreciated being those from the French and English recipes. An ingredient often used in their preparation is ratanhia root, from *Krameria triandra*, a plant indigenous to South America. The root is long, about as thick as a quill, rounded in shape, and with a strongly astringent taste. The alcoholic extract is coloured red.

(a) French recipe—

Aniseed	280 grams.
Cochineal	20 "
Mace	10 "
Cloves	10 "
Cinnamon	80 "
Peppermint oil	20 "
Alcohol	3 litres.

(b) English recipe—

Cedar essence	4 litres.
Myrrh essence	1 litre.
Ratanhia essence	1 „
Lavender oil	20 grams.
Peppermint oil	30 „
Rose oil	10 „

EAU DE BORATE DE POTASSE (BORAX TOOTH TINCTURE).

Borax	150 grams.
Myrrh	150 „
Red santal wood	150 „
Sugar	150 „
Eau de Cologne	1 litre.
Alcohol	3 litres.
Water	1½ „

The solids are digested with the alcohol, the eau de Cologne being then added, followed by the water in which the sugar and borax have previously dissolved. Finally, the whole is filtered.

EAU DE PERMANGANATE DE POTASSE (POTASSIUM
PERMANGANATE WATER).

Potassium permanganate	100 grams.
Distilled water	5 litres.

The potassium permanganate dissolves readily in the distilled water, and gives a beautiful violet solution, a few drops of which are added to a glass of water at the time of using. After the mouth has been cleansed with this solution, a little peppermint water is preferably used for brushing the teeth. On linen and other stuffs, as well as on the skin itself, the above mouth wash leaves brown stains, which can only be removed by the aid of hydrochloric acid.

EAU DE SALVIA (SAGE WATER).

Lemon oil	20 grams.
Sage oil	50 „
Alcohol	1 litre.
Water	4 litres.

The ethereal oils are dissolved in the alcohol, and the solution is united with the water.

EAU DE VIOLETTES (VIOLET WATER).

Violet root essence	1 litre.
Eau de roses, triple	1 „
Alcohol	1 „
Oil of bitter almonds	5 grams.
Neroli oil	2 „

THE PREPARATION OF FACE POWDERS AND ROUGES.

The machinery for manufacturing these preparations consists of a series of apparatus disposed in such a manner that the ingredients can be conveyed from one part to another by means of worm conveyors. The motive power is supplied by a 4.h.-p. electro-motor.

In making LANOLIN or GREASE POWDER, the *modus operandi* is as follows :—

The lanolin or fat is placed in a copper vessel, situated at one side and connected with the main apparatus, and is therein heated to melting point—30 40° C.—by means of hot water, which surrounds the entire apparatus. When this point is reached, the fat is drawn up into a cylinder in the top of the apparatus, by lifting a plunger, a tap in the bottom being then closed in order to prevent the fat flowing back into the heating pan. On allowing the piston to descend, by its own weight and that of the loaded piston-rod to which it is attached, the fat is forced through a perforated false bottom or sieve,

situated between the cylinder and the lower part of the apparatus, and is delivered into the latter in the form of fine spray. To insure the intimate admixture of this spray with a portion of the dry powder, the latter is carried from a vertical storage cylinder by means of a worm conveyor, from which it is blown into the apparatus by the aid of a fan, as a fine dust, through two orifices in the mixing chamber.

The progress of this operation can be followed by looking through a small window in the lower part of the apparatus. The mixture of fat and powder falling to the bottom is set in motion, by stirrers driven from below, to break up any balls of powder that may have formed. At the end of three to four hours the mixture is withdrawn and transferred to another mixing chamber behind the fan, where it is treated with ether, to enable the fat to take up a further quantity of powder. When the ether has been evaporated, there remains a fine powder, forming the basis of the cosmetic in question.

This fatty powder is next mixed with the numerous ingredients by mechanical means, and transported, on a worm conveyor fed by a hopper, to a horizontal drum on the right of the fan, where the different ingredients are united to a homogeneous powder by means of two large rollers, one of which is closely set with short bristles, the other being of fluted wood. The rollers run in opposite directions. On the inner walls of the drum are arranged fan-shaped shovels, which return the materials to the rollers several hundred times in succession. As soon as this treatment is completed, the contents of the drum are automatically transferred to a centrifugal sifting machine, situated directly underneath and covered with fine gauze.

The powder passing through the meshes of this machine falls into an adjacent worm conveyor, and is delivered, in an upward direction, on to plush rollers, which impart softness and tractability to the powder. Thence it falls through a funnel-shaped

tube, and, after being scented, is packed in tins or boxes for sale. By means of this machine about 5 cwt. of powder can be manufactured in a day (ten hours).

The machine for preparing DRY ROUGE PASTE is fed with the rouge powder through a funnel, which delivers the powder into a hollow iron body, from which a sufficient quantity of rouge is discharged at each stroke of the machine to fill a hollow chamber corresponding in size and shape with the block of rouge paste to be manufactured. In this chamber the powder is subjected to the simultaneous pressure of two pistons, acting from above and below respectively, which compress it into a solid lump, and at the same time impress it with the desired design or marks on the upper side. During the return stroke of the machine, the compressed block falls out of the mould, and is then removed and packed into boxes, etc.

STARCH is used for toilet purposes under various names, either alone or mixed with perfumes or colouring matters. All manner of fancy names are bestowed on these preparations: *Blanc végétal*; *Poudre du Sérail*, *Poudre de Riz*, etc., all meaning the same thing, namely, well-washed potato-starch powder. This latter, in fact, is the only form of starch used for toilet preparations, the so-called rice powder being merely potato starch that has been well washed by levigation. Wheat starch is unsuitable for the purpose, owing to the cohesion of the individual granules. Starch may be scented with any desired perfume, but preference is given to such as are most lasting, violet root being in special favour.

Starch may be scented direct by mixing it with powdered violet root. It is, however, difficult to reduce the violet root to such a fine state of division as is required for the purpose in view, to render the granules indistinguishable from those of starch. For finer goods it is therefore preferable to perfume the starch with the scent of the violet roots, without bringing it in contact with the cellular matter of the root. This object

is attained by extracting the perfume from the root by treating it with alcohol in a closed vessel for several weeks, the solvent then acquiring the agreeable odour of the root. On employing this alcoholic solution to sprinkle a quantity of starch powder, and leaving the latter to dry, the solvent evaporates, leaving behind the volatile oil. The perfumed starch must, however, be packed at once into tins or tightly closed boxes, since the scented oil of violet root soon loses its aroma when exposed to the air, and is converted into an inodorous substance.

The same process is applied to the scenting of starch with other perfumes, the odorous vegetable matter being extracted with alcohol, and the solution employed to moisten the starch. Many perfumes are obtainable in a pure state, such as caraway-seed oil, lemon oil, bergamot oil, etc., and these when converted into an alcoholic solution are used to sprinkle the starch spread out in a thin layer.

Toilet starch is also frequently coloured, usually red, in order to impart a delicate red tint to the skin. Only very fine red colouring matters are suitable for this purpose; and pure carmine and finest madder lake are most esteemed. In order to secure the most intimate possible admixture with the starch, it is necessary to add the finely divided colouring matter by degrees, each portion being well triturated with the starch, and the whole finally united and shaken up in a large wooden box. Where large quantities are to be manufactured, it is advisable to effect the mixing in a rotary drum until the starch appears to be of a uniform colour throughout.

Starch that has been coloured red is frequently scented as well, the perfume in this case being imparted by the oil of geranium rose, which has a very delicate odour, resembling true oil of roses. Such scented coloured starches are sold under names like *Rose Powder*, *Poudre de Riz à la Rose*, *Poudre des Dames Rosée*, etc.

The preparation of good powders entails the use of snow-

white steatite powder, very light calcium carbonate, starch, and zinc white. The best flesh tints for blondes is obtained with carmine alone, whilst for brunettes the powder is coloured with burnt umber or sienna. An addition of finely powdered violet root is recommended, since it gives the preparation a fine perfume.

A very good powder is prepared from the following recipe:—

Zinc white, 200 grams; precipitated chalk, 3000; steatite powder, 500; wheat starch, 1000; essence of white rose, 30; jasmine extract, 30; orange-flower extract, 30; cassia extract, 30; musk tincture, $7\frac{1}{2}$ grams.

The following American recipes for FACE POWDER are also worthy of attention:—

1. Zinc oxide	76 grams.
Precipitated chalk	96 „
Powdered talc (steatite)	16 „
Starch powder	32 „

This mixture is scented with a sufficient quantity of extract of roses, jasmine, orange flowers, cassia, or musk.

2. Magnesium carbonate	8 grams.
Powdered talc	16 „

Scented with the necessary amount of rose oil.

3. Starch	7 grams.
Rice meal	1 „
Powdered talc	1 „
Orris root	1 „

Scented the same as No. 2.

4. Zinc oxide	4 grams.
Rice meal	14 „
Precipitated chalk	4 „
Powdered talc	2 „
Orris root	2 „

Scented like No. 1.

5. Zinc oxide	2 grams.
Orris root	2 „
Rice meal	16 „

Scented with the necessary quantity of rose oil, geranium oil, ylang-ylang, cumarin, or acetic ether.

FACE POWDER, WHITE.

Finest levigated zinc white	50 grams.
Venetian talc	50 „
Magnesium carbonate	50 „
Rose oil	20 drops.
Orris oil	30 „

The whole being intimately mixed.

FACE POWDER, ROSE.

Above white powder	150 grams.
Carmine	0.50 gram.

POUDRE DE PISTACHES (PISTACHIO POWDER).

Pistachio meal	5 kilos.
Steatite	5 „
Lavender oil	20 grams.
Rose oil	15 „
Cinnamon oil	5 „

The pistachio meal must be entirely free from oil, and ground as fine as possible.

ROSE POWDER.

Starch	10 kilos.
Carmine	20 grams.
Rose oil	15 „
Santal oil	15 „
Betiver oil	10 „

VIOLET POWDER.

Starch	5 kilos.
Violet root	5 "
Bergamot oil	20 grams.
Lemon oil	20 "
Clove oil	10 "
Neroli oil	10 "

SALICYLIC ACID POWDER.

Starch	500 grams.
Salicylic acid	10 "

SKIN GLOSS.

Sodium carbonate	50 grams.
Powdered spermaceti	50 "
Starch	500 "
Bitter almond oil	10 "
Benzoin	20 "

The ingredients are intimately mixed, and then stored in tightly closed packages. For use, the mixture is stirred in with a little water.

CALODERM.

Wheat starch	2 kilos.
Powdered almonds	500 grams.
Violet root	500 "
Rose essence	500 c.c.
Glycerin	200 "

The whole is mixed to a paste, which is diluted with water before applying it to the skin.

PÂTE AU MUSC, FOR WASHING THE HANDS.

Powered white soap	1000 grams.
Violet root	250 "
Starch	250 "
Lemon oil	20 "
Neroli oil	10 "

Musk essence	5 c.c.
Glycerin	40 „

The mass is boiled until it forms a homogeneous paste, and this is then packed in flat porcelain pots.

ROUGE.

The following preparations can be recommended as fulfilling their purpose, without injuring the skin in any way :—

FRENCH WHITE.

Steatite	2 kilos.
Lemon oil	5 grams.
Bergamot oil	5 „

The streatite is reduced to the finest possible state of division, then levigated, dried, and scented. On account of its greasy character it easily sticks to the skin, and forms the best of all cosmetics of this class, being unalterable in colour and without any injurious action on the epidermis.

LIQUID PEARL WHITE.

Rose water	1 litre.
Orange flower water	1 „
Basic bismuth nitrate	500 grams.

When allowed to stand, the bismuth nitrate settles down completely, leaving the supernatant liquid quite clear. It is therefore necessary to shake up the preparation before use. If left on the skin for any length of time, this cosmetic turns yellow, in consequence of the formation of black bismuth sulphide.

LIQUID VEGETABLE ROUGE.

Ammonia	60 grams.
Carmines	35 „
Esprit de roses, triple	70 „
Rose water	2 litres.

This cosmetic, which is chiefly used for colouring the lips, is

prepared by digesting the finely powdered carmine with the ammonia for several days in a 4 litre flask, then adding the other ingredients, and leaving the whole to stand another week, with frequent shaking. After the sediment has completely settled down, the clear supernatant liquid is decanted into bottles for sale.

In order to obtain this preparation as it should be, neither trouble nor expense must be spared in obtaining the very finest carmine procurable, this being the only way of producing a brilliant vivid red preparation of the proper shade for the lips.

SOLID ROUGE (ROUGE EN PÂTE).

Carmine	.	.	.	.	.	.	30	grams.
Talc (steatite)	.	.	.	.	.	.	600	"
Gum arabic	.	.	.	.	.	.	50	"

The ingredients are powdered as finely as possible, then intimately mixed by trituration in a porcelain mortar, and formed into a paste by the careful addition of water. This paste is packed in very shallow porcelain pots, about 1½ inches in internal diameter. A darker rouge, such as is required for the stage or for persons with brunette complexions, is obtained by increasing the proportion of carmine.

LEAF ROUGE (ROUGE EN FEUILLES).

Circular discs, about 2 inches in diameter, are cut from thick, very smooth, highly satinised paper, and covered with a layer of carmine, the adhesion of which to the paper is secured by an addition of gum arabic. For use, all that is necessary is to breathe on the leaf, rub it over with a ball of fine cotton-wool, and apply the latter to the face.

VEGETABLE BLUE FOR VEINS (BLEU VÉGÉTAL POUR LES VEINES).

Venetian chalk	.	.	.	.	.	500	grams.
Berlin blue	.	.	.	.	.	50	"
Gum arabic	.	.	.	.	.	30	"

The powdered ingredients are converted into paste with a little water, and then shaped into sticks. For use, the stick must be breathed upon, then rubbed with the rough side of a piece of white glove leather, and the blue veins traced on the skin, which has previously been coated with pearl white. That a certain amount of skill is necessary to produce the proper effect, goes without saying.

VEIN BLUE FOR THEATRICAL MAKE-UP.

White bole	.	.	.	.	.	80	grams.
Zinc white.	.	.	.	.	.	10	"
Berlin blue	.	.	.	.	.	10	"
Gum arabic	.	.	.	.	.	3	"

These ingredients are worked up to a plastic mass with orange-flower water, then made into sticks, about $\frac{1}{4}$ inch thick, and dried.

HYDROGEN PEROXIDE FOR THE SKIN.

For delicate skins, hydrogen peroxide may be used as a wash, if diluted with its own volume of water, a little ammonia—1–2 drops to a dessert-spoonful—being added to the wash at the time of use. Where the hydrogen peroxide comes in contact with the skin, small bubbles of oxygen will be observed to form, whilst at the same time the dead rough surface of the skin will be converted into a white, soap-like mass. Since the peroxide destroys merely the dead portions, the smooth skin makes its appearance, and, being left unattacked in any way, soon becomes strong and able to resist external influences.

BIRCH LEAF AND ST. JOHN'S WORT OINTMENT.

A patented preparation from the above ingredients is used for burns and roughness of the skin. The half unfolded young leaves of the birch tree are extracted with alcohol for a week, the dark green resinous extract, then obtained by

pressure, exhibiting the aromatic odour of the birch, and containing all the efficacious constituents of the leaves.

The freshly cut *St. John's wort*, gathered while in bloom, is digested for three weeks with a mixture of wool fat and oil at about 40° C. The viscous reddish brown extract recovered by pressure contains all the active ingredients of the sap, in consequence of the faculty of wool fat for taking up aqueous liquids.

The two extracts are mixed in equal proportions, and, on the alcohol being driven off by evaporation, furnish a dark green ointment.

CHAPTER VII.

HAIR DYES, DEPILATORIES, AND HAIR RESTORERS.

Preparations for Darkening the Hair.

BLACK DYES.

1. **T**HIS dye consists of a 2-5 per cent. solution of silver nitrate, supersaturated with ammonia. It is applied to the hair after the latter has been well washed with soap. When the foregoing solution has dried on the hair, a second solution is applied, consisting of dilute alcohol containing 3-4 per cent. of pyrogallic acid, or a 6-8 per cent. solution of potassium sulphide.

2. The pericarp of the cashew apple contains, in addition to gallic acid, two special organic substances, one of which, cardol, is of oily consistence, and has a highly stimulative action on the skin; whilst the other, anacardic acid, which is for the most part present in the free state in the husk, does not share the injurious properties of cardol. To extract it from the pericarp, the latter is cut up and treated with ether, the solvent being driven off and the residue extracted with water until the washings cease to give the tannic acid reaction. The residue is then dissolved in 15-20 parts of alcohol, and shaken up with freshly precipitated lead hydroxide, quickly filtered, and washed with alcohol. It is advisable to exclude air as far as possible during this process. The precipitate consists of lead anacardate, whilst the cardol, being left behind dissolved in the alcohol, is thus separated from the anacardic

acid. On treating the precipitate with ammonium sulphide and sulphuretted hydrogen, and filtering the liquid, the filtrate will contain, in addition to the excess of ammonium sulphide, the anacardic acid in the form of an ammonium salt. The filtrate is then concentrated to a small volume, and treated with sulphuric acid, which causes the deposition of the anacardic acid in the form of white masses which quickly solidify. These masses are pressed between filter paper and then dissolved in ammonia, the solution being soluble in water, and forming a hair dye. In application, the hair is rubbed over with this solution, and afterwards combed with a comb dipped in ferrous sulphate, whereupon it soon assumes a more or less black colour, which is fairly lasting. Anacardic acid can also be made up into a pomade with vaseline or hair oil; and ferrous sulphate may be replaced by iron oleate as the developer. This last-named compound is obtained by the interaction of ferrous sulphate and medicinal soap. It is advisable to keep both the anacardia pomade and developer in tin tubes.

3. *Tannin Hair Dyes*.—(a) In a clear glass bottle—

Powdered gall nuts	.	.	.	.	400 grams.
Water	.	.	.	.	500 "
Rose water	.	.	.	.	500 "

The gall nuts are boiled with the water, the boiling liquid being strained through a close cloth into the rose water, and filled, whilst still hot, into the bottles, which should be closed at once. Unless filled while hot, the contents of the bottles are liable to develop mould.

(b) In a dark bottle—

Silver nitrate	.	.	.	.	150 grams.
Water	.	.	.	.	1 litre,

together with sufficient ammonia to redissolve the precipitate formed at first.

4. *Melanogene*.—(a) In a dark bottle—

Silver nitrate	10 grams.
Distilled water	80 „
Ammonia	30 „

(b) In a clear glass bottle—

Pyrogallic acid	1 gram.
40 per cent. alcohol	500 grams.

5. *Eau d'Afrique*.—(a) In dark bottles—

Silver nitrate	3 grams.
Distilled water	100 „

(b) In clear glass bottles—

Sodium sulphide	8 grams.
Distilled water	100 „

6. *Krinochrome*.—(a) In clear glass bottles—

Pyrogallic acid	10 grams.
Distilled water	180 „
Alcohol	100 „

(b) In dark bottles—

Silver nitrate	12 grams.
Ammonia	60 „
Distilled water	300 „

7. Potassium permanganate, 150 grams; distilled water, 2 litres. The hair is washed, and the solution is applied with a soft brush; the dye takes effect immediately, and the hair is coloured, according to the degree of dilution, all shades, from blonde to the deepest brown.

8. One part of powdered peat, free from sand, is left in contact with 10 parts of ammonia and 5 parts of water for two days in a glass flask, then slowly heated to boiling, strained through a cloth, and concentrated to the consistence of syrup, on the water bath. The resulting brown extract is dissolved in 10 parts of distilled water containing 2 parts of alcohol and a little eau de Cologne. After the hair has been washed with

soap, spirit and water, to remove grease, it is damped with the brown liquid.

9. *Bismuth Hair Dye*. — One hundred grams of metallic bismuth are dissolved in a sufficiency of ordinary nitric acid (about 280 grams), the solution being treated with a concentrated solution of 97 grams of tartaric acid, and precipitated by a copious addition of water. The precipitate is then washed with water until the washings no longer give a strong reaction with blue litmus paper, after which it is carefully separated from the filter paper and dissolved in ammonia. This solution is employed to dissolve 75 grams of purified sodium hyposulphite, filtered, treated with an addition of 2-5 per cent. of glycerin, and filled into bottles. It contains about 5 per cent. of bismuth, and should be applied once daily to the hair or beard to be dyed. The colour develops slowly to a deep brown, and when this result is produced further applications are made only as required.

10. Two liquids are used: (a) 10.15 glycerin, 6.42 sal ammoniac, 76.18 water, and 7.25 bismuth nitrate; (b) 1.05 pyrogallie acid, 13.68 sodium hyposulphite, and 85.27 gaultheria water.

BLONDE DYE.

1. A dilute solution of hydrogen peroxide is applied with the brush, after freeing the hair from grease.

2. (a) 5.0 potassium permanganate, 95.0 distilled water. This solution is kept in a brown bottle, with ground stopper.

(b) 1.0 sodium hyposulphite, 25.0 distilled water. The hair is washed with a solution of soda, rinsed with warm water, and dried. Solution (a) is then applied with a brush and comb (which must not be greasy), and evenly distributed through the hair. To remove stains on the skin, a small rag is moistened with water, followed by a little soap and solution (b), and used to rub out the stains.

3. The hair, freed from grease, is moistened with a mixture of 1 part of iron acetate, 1 part of silver nitrate, and 2 parts of bismuth nitrate dissolved in 10 parts of water, the colour being developed, after the lapse of an hour, by means of a solution of potassium bisulphide in its own weight of water.

4. A yellow hair dye may be obtained by the aid of Orleans, boiled in water that has been rendered slightly alkaline with soda. The hair must first be washed with a solution of alum. A Nanking yellow colour can be imparted to hair by treating it with a dissolved neutral salt of iron, *e.g.* sulphate, acetate, or chloride, followed by a solution of an alkali—sodium carbonate, potassium carbonate, or lime water. This dye will modify the colour of fiery red hair.

Preparation for darkening red hair. A solution is prepared of 3.5 grams of calcium saccharate, 15 grams of glycerin, 7 grams of jockey club, 15 grams of rectified spirit, and sufficient water to make 300 c.c. With this liquid the hair is sprayed every morning, and then well brushed.

Depilatories.

CALCIUM SULPHIDE.

This preparation is made ready in the following manner:—Quicklime, calcined to whiteness, is suffused with water,—an operation requiring some care, owing to the heat disengaged,—the resulting powder being shaken loosely into a high glass, containing a wide glass tube reaching to the bottom and connected with an apparatus charged with water and ferrous sulphide. On treating the latter with sulphuric acid, sulphuretted hydrogen is disengaged, and, passing through the aforesaid tube, is immediately absorbed by the lime. When the absorption is complete and the whole of the lime has been converted into calcium sulphide, as may be recognised by the escape of sulphuretted hydrogen from the lime vessel, the generator is disconnected,

and the calcium sulphide is worked up into a paste, as follows:—

Calcium sulphide	.	.	.	.	2 kilos.
Sugar	.	.	.	.	1 kilo.
Water	.	.	.	.	1 "
Starch	.	.	.	.	1 "
Lemon oil	.	.	.	.	30 grams.
Peppermint oil	.	.	.	.	10 "

This paste must be packed immediately into air-tight pots, since the calcium sulphide is decomposed on exposure to air. For use, a little of the mass is moistened with sufficient water to make it easy of application to the skin, and is laid on the hairs to be removed. At the end of a half to three-quarters of an hour, the whole must be washed off with water.

BARIUM SULPHIDE.

This preparation can be obtained by passing sulphuretted hydrogen over solid barium hydroxide and mixing the product with starch paste. It is similar in action to calcium sulphide, but has the disadvantage of being more expensive and less stable.

BARIUM SULPHIDE PASTE.

This must always be prepared immediately before use by stirring up barium sulphate, carbon, and linseed oil to a paste, and calcining the mixture thoroughly, a dark blue product, barium sulphide, being obtained. Ten parts of this, rubbed down fine, are mixed with 5 parts of zinc oxide and 5 parts of starch, the powder being incorporated with sufficient water to form a soft paste, which is then spread fairly thick, by the aid of a wooden spatula, on the hair to be removed. When perfectly dry, *i.e.* in about ten minutes, the paste is washed off with water, and leaves the skin clear and free from hair.

According to the patent specification of G. Huettemann and

J. Zrzawy, of Brüx (Bohemia), a stable depilatory, which does not affect the skin, is obtained by quenching quicklime with a solution of sugar, and passing sulphuretted hydrogen through the resulting solution of calcium saccharate. The process can also be modified by first preparing a solution of sugar saturated with sulphuretted hydrogen, and then stirring this up with the lime. This furnishes a pale green mass, which is then mixed with such a quantity of lime that the mixture contains about 5 per cent. of the fundamental substance.

For use, this mixture is stirred up to a pap with water, and applied to the surface to be freed from hair, the whole being washed off again after about ten minutes.

Hair Restorers.

After a long period of neglect, the beneficial effect of bay rum on the scalp is beginning to be more and more widely recognised on the Continent. This solution of bay oil in alcohol is renowned for its refreshing action in cases of nervous headache, languor, and other nervous disturbances.

Even superior to bay rum as a hair wash, is the following preparation:—

A solution is made of 11 grams of bay oil and 5 drops of clove oil, in 500 grams of alcohol, with which are incorporated 1500 grams of tincture of Quillaya bark and 10 grams of Tinct. Capsici annui. The whole is mixed with 0.5 gram of mace and 15 grams of ammonium carbonate, with frequent shaking, and is filtered at the end of three days. Even repeated filtration, however, will scarcely furnish a clear solution, and therefore the liquid should be stored in blue or opaque bottles. The Quillaya bark tincture is prepared by digesting 20 parts of Quillaya bark with 100 of alcohol, and mixing with 120 parts of rose water, 20 of eau de Cologne, and 30 of glycerin.

The following recipes have been given for preparing bay rum :—

Bay oil	.	.	.	.	.	16 parts.
Clove oil	.	.	.	.	.	1 part.
Pimento oil.	.	.	.	.	.	1 „
Jamaica rum essence	.	.	.	.	.	75 parts.
Alcohol	.	.	.	.	.	2650 „
Water.	.	.	.	.	.	1850 „

mixed together and filtered.

Seventy parts of pure bay oil and 200 parts of Cachaca essence are dissolved in 10,000 parts of 95 per cent. alcohol, and mixed, by frequent shaking, with 3000 parts of triple concentrated rose water and 3000 parts of water. When intimately mixed, the whole is left at rest for one to two days, and then filtered. An inferior quality may be prepared by replacing the rose water with water.

One kilo of 95 per cent. spirit is mixed with 14 grams of bay oil, the mixture being left for fourteen days, and then incorporated with 2 kilos of Jamaica rum.

A mixture is compounded of 200 grams of 95 per cent. spirit, 200 grams of water, 1 gram of bay oil, 10 drops of pimento oil, and 1 drop of clove oil, and is filtered after standing several days.

One kilo 100 grams of 95 per cent. spirit, 4 grams of warm water, 30 grams of Venetian soft soap dissolved in 460 grams of warm water, 60 grams of Jamaica rum essence, 6 grams of bay oil, 4 grams of clove oil, 30 grams of absolute alcohol, and 1 gram of cinnamon oil, form another mixture.

According to Fr. Toellner, a fine glycerin preparation can be compounded by dissolving 0·4 of heliotropin, 0·5 of ionone solution, and 3 drops of rose oil in 25·0 of 95 per cent. alcohol, filtering, and mixing the filtrate with 1000·0 of chemically pure glycerin.

SOAP ESSENCE.

The most suitable soap for preparing soap essence is found to be white olive oil soap, the potash in which plays a useful part in the preparation of these essences for cosmetic purposes. The principal formula is—

White olive oil soap, 200 parts ; 80 per cent. alcohol, 1000 parts ; pure potassium carbonate, 12 parts. The soap is dissolved in the alcohol by agitation in a wide-mouthed flask over a water bath, and the potassium carbonate is added to the solution, which is then slightly coloured—with safranin or rosaniline—and scented. The whole is left to settle for a few days, and afterwards filtered. A very fine scent is prepared by mixing 10 parts of vanilla tincture, 20 parts of violet root tincture, 20 of rose extract, and 50 of orange-flower extract.

Rodiguet recommends for the preparation of these essences—

White soap, 1 part ; 85 per cent. alcohol, 3 ; distilled water, 1 part. And for hygienic purposes, the following mixture, which lathers copiously with water :—Olive oil soap, 20 parts ; distilled water, 30 parts ; 60 per cent. alcohol, 60 ; potash, 1 part ; and scent, 1 per cent.

To this class belong also the so-called scent essences. For these, spirit of wine is recommended ; corn spirit, if used at all, requires to be free from fusel oil, and to be filtered through a mixture of equal parts of washed bone black and levigated chalk. The finished mixture only acquires its full aroma after storage for some time, and will be improved by distillation at the lowest possible temperature over the water bath.

A preparation closely resembling true eau de Cologne can be compounded from the following recipe :—

Siam benzoin, 66 grams ; lavender oil, 133 grams ; rosemary oil, 66 grams ; neroli oil, 350 grams ; petit grain oil, 350 grams ; cedrate oil, 350 grams ; Portugal oil, 700 grams ; bergamot oil, 700 grams ; French oil of geranium rose, 60 grams ; 95 per cent.

alcohol, 140 litres. The ethereal oils are dissolved in the spirit in the order given above, the finely powdered Siam benzoin being added last. After the mixture has been left to stand for at least four weeks, with repeated shaking, it is placed in a still along with a corresponding quantity of water, and 135 litres are collected, after rejecting about 2 litres of first runnings. The distillate is left to stand another month in dark-coloured glass vessels in a dark place, and is then ready for filling into bottles for use.

The preparation of soap scents, for supplying small soap makers, ought to be a very profitable branch of the manufacture of cosmetics. A few recipes for this purpose are given below, the figures quoted referring to grams, and the total quantities sufficing for 1 cwt. of soap:—

For Marshmallow soap.—Lavender oil, 50; clove oil, 30; cassia oil, 30; ginger-grass oil, 40; Peruvian balsam, 120; tolu tincture, 70.

For Herb soap.—Marjoram oil, 20; Curomoji oil, 25; curly-mint oil, 20; sage oil, 20; rosemary oil, 30; calmus oil, 15; caraway oil, 80; lavender oil, 80; clove oil, 20.

For Clove soap.—Lemon oil, 25; cassia oil, 25; santal-wood oil, 25; cedar-wood oil, 25; clove oil, 200.

For Cinnamon soap.—Lavender oil, 50; clove oil, 30; lemon-grass oil, 40; cassia oil, 250.

For Vanilla soap.—Peruvian balsam, 80; clove oil, 40; lavender oil, 40; palmarosa oil, 30; vanilla tincture, 250; benzoin tincture, 150.

For Geranium soap.—Palmarosa oil, 100; African geranium oil, 100; lavender oil, 30; cedar-wood oil, 30; turpentine, 40.

For Eau de Cologne soap.—Lavender oil, 50; rosemary oil, 40; Portugal oil, 160; lemon oil, 50; lemon-grass oil, 40.

For Balsam soap.—Nutmeg oil, 10; palmarosa oil, 40; clove oil, 25; cassia oil, 20; Peruvian balsam, 200; benzoin tincture, 50.

For Windsor soap. — Lavender oil, 100 ; caraway oil, 75 ; rosemary oil, 50 ; oil of thyme, 25 ; lemon-grass oil, 40.

Mention should also be here made of the preparation of Quillaya bark tincture—from 1 part of Quillaya bark cut into strips, and 5 parts of dilute alcohol. Dieterich, however, states that the extraction of the bark is incomplete except when an excess of water is employed, and he therefore recommends the following improved recipe :—

Quillaya bark	.	.	.	.	.	1 part.
Water	.	.	.	.	.	4 parts.
90 per cent. alcohol	.	.	.	.	.	1 part.

COSMETIC POMADES.

The care of the skin leads to a good demand for cosmetic preparations, especially in the cold season. Cold, damp winds, low temperature, and the widespread neglect to dry the hands thoroughly after washing, cause the skin to take cold in many ways, the effects being recognised by brittleness and chapping.

In addition to vaseline and glycerin, lanolin is a substance that has made its way on to nearly every toilet table ; and as this preparation is now obtainable ready packed by makers of cosmetics, many of them have abandoned the production of lanolin crème, although this is a more profitable venture, and the sale can be largely increased by employing perfumes to suit the current taste.

A lanolin toilet cream, prepared according to the following formula, will rapidly gain favour under all circumstances :—

Anhydrous lanolin (Norddeutscher Wollkaem- merei)	.	.	.	.	.	650 grams.
Peach kernel oil	.	.	.	.	.	200 „
Water	.	.	.	.	.	150 „

This may be scented with about 15 drops of ionone, or 20 drops of synthetic ylang-ylang (Schimmel).

The so-called lanolin milk, prepared in the following manner, is also highly appreciated :—

Anhydrous lanolin 100 grams.
melted and mixed with

Glycerin 100 „
Water 750 „

the whole placed in a wide-necked vessel, and to it added, with continued energetic shaking—

Benzoin tincture 50 grams.
Gum mucilage 30 „

finally scenting the preparation as in the case of the cream.

The glycerin gélées have the advantage over lanolin of being more economical; but, on the other hand, are attended with the drawback that they are less readily absorbed by the skin. These preparations are packed either in glasses or tubes, the latter being preferable, and more and more popular, on account of their greater handiness.

The following is a good recipe for such glycerin gélées :—

Finest white tragacanth powder 50 grams.
mixed with

Alcohol 100 „
Glycerin 200 „

shaken up with a suitable amount of perfume, and then quickly incorporated, by shaking, with

Water 650 grams.

This forms a transparent mucilage, which can be filled into bottles, etc., at once.

The reduced consumption of egg yolks, owing to the introduction of the chrome tanning process on the one hand, and the largely heightened demand for egg albumin for the preparation of albuminised paper on the other, induced Bernegau to seek new methods of utilising the former article, egg yolk. He

succeeded in applying this material to the manufacture of cosmetics, and in this connection remarks that he ascertained the capacity of yolk of egg for being rapidly absorbed by the skin, by massage experiments on his own hands, and found beneficial results from the operation. After numerous trials, he recommends the following as the best method of preparing a toilet cream from the aforesaid ingredient, and it undoubtedly forms a profitable article:—Preserved yolk of egg, 1 part; benzoinated olive oil, 1 part; alapurin, 1 part, scented with rose oil or with cumarin and violets, are worked up to a toilet cream to which the inventor has given the name Viteline Crème. This cream is specially adapted to the preparation of excellent toilet soaps, which it renders extremely soft and tractable. It has a beautiful yellow colour, and, apart from its good properties as an emollient for the skin, it is also calculated to achieve a good reputation as a cosmetic for the hair. Alapurin may be replaced by any other preparation of wool fat, or by vaseline; and boroglycerin may be substituted for the benzoinated olive oil.

Camphorated vaseline ice, an appreciated toilet article, is prepared from—

Coarsely powdered camphor	.	.	.	.	2 grams.
Paraffin	.	.	.	.	2 "
Glycerin	.	.	.	.	2 "
White wax	.	.	.	.	5 "
White vaseline	.	.	.	.	8 "

The paraffin, vaseline, and wax are melted together, the camphor being dissolved in this mixture (by warming, if necessary), and the glycerin incorporated by stirring until cold.

A brilliant transparent pomade may be prepared from the following recipe:—

270 grams of perfectly pure tallow, free from water, are weighed out and melted along with 120 grams of pale resin, and heated to 87·5 C. Then 160 grams of 38° caustic

soda lye are heated to 75° C., mixed with 300 grams of 92-94 per cent. alcohol, and the alcoholic soda mixture stirred into the fat, a large vessel being used in order to prevent the mass overflowing. A kind of transparent soap is produced. Another method of attaining the same end is to weigh out 8000 grams of East Indian castor-oil, 3000 grams of Cochin cocoanut oil, 5000 grams of olive oil, and heat them to 75° C. The above soap is heated along with 4000 grams of 96 per cent. alcohol until bubbles cease to rise, whereupon the two—oil and soap solution—are well stirred up together, coloured with gamboge dissolved in alcohol, and the pomade scented with any desired perfume.

A new recipe for preparing cold cream without stirring has been proposed by Idelson, and is said to furnish a particularly white very homogeneous, and stable preparation. To this end 135 grams of white wax, 75 grams of spermaceti, and 540 parts of white vaseline are melted over a gentle fire, then strained into a warm, wide-necked bottle, and into the mixture is slowly poured a hot solution of 12 grams of borax in 180 grams of rose water and 20 drops of geranium oil, the whole being well shaken up and stored in a cool place.

A lip salve that never turns rancid, and always retains its handsome red tinge, is prepared from—

Paraffin	80.0 grams.
Vaseline	80.0 "
Alkannin	0.5 gram.
Bergamot oil	1.0 "
Lemon oil	1.0 "

Scurf pomade.—The following recipe is recommended :—

Benzoinated lard	120 parts.
Precipitated sulphur	4.9 "
Lanolin	20 "
90 per cent. alcohol	20 "
Salicylic acid	1 part.
Geranium oil	1 "
Rose water	60 parts.

In connection with the care of the hair, mention may here be made of the recipes for preparing China pomade and cantharides pomade.

China pomade—

Beef marrow	335 grams.
Lard	225 „
Oil of sweet almonds	55 „
Peruvian balsam	3 „
Quinine sulphate	3 „
Clove oil	7 „
Rose essence	30 drops.

Cantharides pomade—

Beef marrow	900 grams.
White wax	100 „
Mace oil	3 „
Clove oil	3 „
Rose essence, or geranium oil	30 drops.
Cantharides tincture	28 „

As an appendix to these preparations come the wax pomades, beard pomades, bandolines, and brilliantines :—

POMADE FIXATEUR À LA ROSE

is prepared in the following manner :—

Gum arabic	1 kilo.
Soap	1 „
Wax	2 kilos.
Rose water	2 litres.
Bergamot oil	20 grams.
Lemon oil	100 „
Oil of thyme	10 „

The gum and soap are dissolved in the rose water, the ethereal oils being then added, followed by the wax, and the liquid is stirred until set. This pomade is also coloured, three kinds being prepared, namely, blonde (uncoloured), brown,

and black. The brown is produced by an admixture of umber or terra di Siena, and black by ivory black. The colours must be ground to an extremely fine state of division with oil, stirred up with wax, and added to the other ingredients. The resulting wax pomades are fairly hard; when the weather is cold, they may be softened to the requisite degree by holding the vessel in the hand for a short time.

STICK POMADE.

This is a wax pomade which differs from the foregoing mainly in being merely rather more solid. It is usually prepared in the form of sticks, of circular or oval section, wrapped in tinfoil, and coloured and scented to taste. The usual kinds are white (uncoloured) pomade for light blonde hair; rose-red, coloured with carmine; brown, coloured with umber; and black, prepared with ivory black. The colours are always ground in oil. The rose-red pomade is prepared with alkanna root, which is digested for some time with the melted fat. The basis of these pomades consists of—

Lard	.	.	.	.	.	.	2 kilos.
Talc	.	.	.	.	.	.	6 "
Wax	.	.	.	.	.	.	3 "

and the consistence can be varied by changing the proportion of wax. They are usually scented with bergamot, lemon, clove, or thyme oil.

HUNGARIAN POMADE (POMADE HONGROISE).

This is prepared from—

Gum arabic	.	.	.	.	.	1 kilo.
Soap	.	.	.	.	.	1 "
Wax	.	.	.	.	.	1 "
Rose water	.	.	.	.	.	1 "
Bergamot oil	.	.	.	.	.	40 grams.
Santal-wood oil	.	.	.	.	.	20 "

For brown and black, the pomade is suitably coloured. It is made up into sticks about the thickness of a lead pencil.

WAX FOR THE MOUSTACHE AND BEARD.

Turpentine	1 kilo.
Almond oil	1 „
Wax	3 kilos.
Violet pomade	1 kilo.
Peruvian balsam	500 grams.
Clove oil	30 „
Santal-wood oil	20 „
Cinnamon oil	20 „

In preparing bandoline, 400 grams of tragacanth are powdered, placed in 8 litres of rose water, and left in a warm place, with frequent stirring, until the tragacanth has swollen up into a mucilaginous mass. This is then pressed through a coarse cloth, afterwards through a finer one, and finally receives an addition of a little liquid carmine and 20 grams of oil of bitter almonds.

A useful brilliantine is prepared from 4 kilos of glycerin and 21 grams of jasmine—or other extract; also from—

Glycerin	2 kilos.
Castor-oil	2 „
Bergamot oil	20 grams.
Lemon oil	20 „
Neroli oil	10 „

We will now consider the manufacture of preparations for use against freckles, moles, warts, and sweaty hands :—

FRECKLE WASH (EAU DE HEBE).

Distilled vinegar	660 grams.
Lemons, chopped in fine pieces	135 „
85 per cent. alcohol	88 „
Lavender oil	23 „
Water	88 „
Cedra oil	6 „

This mixture is left to stand in the sunlight for three to four days, and filtered. It is applied to the freckled skin with a soft sponge before retiring to rest, and is allowed to dry on.

Another wash for freckles is compounded from—

Angelica root	50	grams.
Orris root	50	„
Styrax	20	„
Bergamot oil	10	„
Lemon oil	10	„
Alcohol	2	litres.

EAU LENTICULEUSE.

Potassium carbonate	200	grams.
Sugar	20	„
Orange-flower water	2	litres.
Alcohol	200	grams.

LILIONESE I.

Potassium carbonate	400	grams.
Water	2000	„
Rose water	400	„
Alcohol	200	„
Rose oil	10	„
Cinnamon oil	5	„

LILIONESE II.

Rose water	2	litres.
Orange-flower water	1	litre.
Glycerin	500	grams.
Potassium carbonate	100	„
Benzoin tincture	20-50	„

The quantity of the alcoholic benzoin tincture used is just so much as is required to make the liquid opalescent or slightly milky.

The following recipe is said to furnish a reliable wash for removing freckles:—

A solution is prepared of dilute nitric acid, 7 grams; eau de

Cologne, 115 grams ; and neroli oil, 10 drops : to which is added, hydrogen peroxide, 60 grams ; glycerin, 100 grams ; cochineal extract, 3·5 grams ; and enough distilled water to make 325 grams. It is then left for a fortnight before filtering. The preparation is applied with a linen rag after the face has been washed, and is allowed to dry on.

MOLES.

The best remedy for these is hydrogen peroxide, a piece of cotton-wool, soaked in this liquid, being laid on the mole for five minutes, morning and evening ; the resulting irritation and scaling of the skin are counteracted by rubbing with lanolin or by coating with a solution of borax and glycerin. Consequently, the manufacturer of cosmetics has to make two preparations for curing moles, namely, hydrogen peroxide and a lanolin pomade. The method of application should be printed on the wrappers of both. The hydrogen peroxide must be filled into opaque bottles and labelled somewhat as follows : "Moisten a pad of cotton-wool with this liquid, and apply the same on the mole, morning and evening, five minutes at a time." The lanolin pomade should be labelled : "After the mole has been moistened with the liquid for five minutes, a little of this salve should be rubbed on the spot."

WARTS may be eliminated in a simple and reliable manner by the use of formalin,—40 per cent. formaldehyde,—a piece of stick being steeped in this liquid and rubbed well into the wart. The operation is said to be painless, except where cracks occur in the flesh, when a temporary smarting is produced. After two or three applications, the wart will begin to shrink, and will fall off without leaving any scars. The cosmetic manufacturer could therefore also sell formaldehyde solution for warts, with printed directions on the bottle, and perhaps supply a suitable wooden instrument as well.

Another remedy for warts, also exhibiting the advantage of not leaving any scar behind in addition to being painless, has been proposed by Couvel-Dubregyré. This preparation can also be made up by the cosmetic manufacturer in fancy packages, with directions for use. It consists of a solution of potassium bichromate in hot water; on cooling, this deposits crystals, and the supernatant liquid is applied to the warts with a brush, once or twice a day.

A remedy for SWEATY HANDS—a much-sought preparation—is furnished by the following mixture, and finds a ready sale:—

Borax, 4·0; salicylic acid, 4·0; boric acid, 1·0; glycerin, 16; alcohol, 16·0. The solids are dissolved in the spirit, the glycerin being added afterwards. The preparation is used for rubbing into the hands three to four times a day.

A proved remedy for the smell of perspiration, capable of easy preparation and direct use, is the following:—A 10 per cent. solution of lead acetate is treated with 50 grams of calcined magnesia, and applied externally.

For protecting the hands against the action of caustic or corrosive antiseptics, the following precautions may be adopted:—

Carbolic acid and corrosive sublimate make the skin brittle and cracked when frequently brought in contact therewith, even in the case of dilute solutions. Washing with water is not sufficient to cleanse the hands thoroughly. After carbolic acid, the best thing to wash with is alcohol, a solution of common salt being the best after corrosive sublimate; a thoroughly good soaping is advisable in either case, and a spoonful of borax should be rubbed in with the soap lather. After the hands are dry they should be rubbed over with a little lanolin.

Disagreeable smells, such as those from iodoform, creosote, or guaiacol, can best be removed by washing with linseed meal or mustard meal and water. The smell of iodoform, in particular, can be easily removed by rubbing the hands with ergot powder

and water. Ground coffee, oil of turpentine, tar water, and ethereal oils are also recommended for this purpose. When the hands are dry, they should be dusted over with powdered talc, renewed every hour, in order to prevent eczema, which frequently results. Chapping of the hands, so often occurring in winter, may be prevented by rubbing them with a mixture of glycerin and rosewater, or glycerin, boric acid, and vaseline. For healing cracks and chaps, the following salve is recommended:—Menthol, 0·75; salol, 1·5; olive oil, 15·0; lanolin, 45·0.

Particulars have already been given for preparing a rightly appreciated preparation of crushed almonds, but it differs from the one now subjoined, namely—Crushed almonds, 5 parts; siliceous earth, 6 parts; potato starch, 5 parts; glycerin, 2 parts; violet root, 1 part; borax, 2 parts; talc, 1 part. Oil of bitter almonds or nirtobenzol is employed as the scent. Another, but less useful, preparation for the same purpose is the following:—Calcined soda, 50 grams; powdered borax, 100; finest violet root powder, 250; and 2000 of finest wheat flour or rice meal, are mixed, and with the mixture are incorporated 120 of fatty almond oil. Finally, 100 grams of alcohol and about 50 drops of oil of bitter almonds are added, and the whole is rubbed through a fine hair sieve.

PLAISTERS.

The paper plaisters, first described by Redoin, are also highly suitable for cosmetic purposes. This author has given the following recipes:—

Salicylic acid plaister.—The paper is dipped in a mixture of salicylic acid, 1 part; paraffin, 50; and liquid paraffin, 50 parts.

Sublimate plaister.—Blotting paper—not satin paper—is steeped in a solution of 2·5 parts of corrosive sublimate in 1000 of spirit, or saturated with a hot solution of 2 parts of sublimate in 1000 of water and 50 of glycerin, and then hung up to dry.

Styptic plaister.—Satin paper is steeped with a solution of

1 gram of alum in 18 grams of ferric chloride solution. When blotting paper is used, a hot solution of aluminium sulphate, 2.0 parts; burnt alum, 1; benzoic acid, 1; ferric chloride solution, 6; and distilled water, 4 parts, is employed.

Antiseptic sticking plaister. — A substitute for diachylon plaister can be prepared from satin paper steeped in the following solution:—Salicylic acid, 1 part; gum arabic, 45; distilled water, 55 parts.

The paper used for the foregoing is of the kind generally known as cigarette paper.

The author, being interested in the preparation of an antiseptic sticking plaister as a substitute for diachylon plaister, has worked out a process for making this on the large scale. The solution used for this purpose is that just mentioned, consisting of salicylic acid, gum arabic, and distilled water. The salicylic acid is dissolved by warming the water, and when this is accomplished the gum arabic is brought into solution. This solution is very viscous on account of the high percentage of gum, but it is not clear, nor is it necessary that it should be so.

Redoin, in his directions, speaks of steeping and impregnating the cigarette paper in the solution. This, however, is not feasible in the general acceptance of the term, the solution being too concentrated to penetrate the paper sufficiently. Moreover, paper plaister prepared in this way would be as sticky on one side as the other, which would render the application of the plaister well-nigh impossible, and certainly would not fit it to replace diachylon plaister. In addition, Redoin's directions are attended with a further important drawback, namely, that in consequence of the large proportion of gum present—which, however, is essential—the paper, on drying, becomes brittle and liable to crack, so that it is difficult to cut it into strips or small pieces for application to wounds, etc., and therefore the plaister is practically unsaleable.

However, the author has found a way to obviate all these inconveniences, and has succeeded, by a very simple means, in producing an antiseptic sticking plaster of faultless outward appearance, with a brilliant gloss on the side applied to the skin. It can be cut into pieces of any size, has been pronounced by all who have seen it a perfectly saleable article, and has proved, in use, capable of practically replacing the more expensive diachylon plaster.

In making this article, the above-named solution is mixed thoroughly with about 2-3 grams of concentrated, water-white glycerin, which renders the paper soft, supple, and very glossy. The glycerin, being hygroscopic, produces a welcome softness, and makes the paper handle well; the sheets can be bent in any direction, folded, cut, torn; in fact, one can do anything with it, and it never gets glassy, brittle, or cracked.

Care, however, is necessary not to overdo a good thing, since, if too large an amount of glycerin be taken, its hygroscopic properties become unpleasantly apparent, the paper refusing to dry, with the result that, when two sheets are laid together, they cannot be separated without tearing.

So far as the steeping of the paper is concerned, the method adopted, as the result of practical experiments, is as follows:—The solution is placed in a porcelain basin, the sheet of cigarette paper is spread out on a board, and the solution is applied to the one side with a broad, flat brush. In winter, the uniformly coated sheets are placed to dry near a stove, but in the summer are merely exposed to the air. Drying ensues in a very short space of time, and as soon as this is accomplished the sheets can be cut to the proper size for packing.

This paper when moistened possesses great adhesive power, and when applied to the wound, etc., to be protected, never stretches or tears in drying. Being glossy only on the sticky side, it is easy to tell which side should be moistened for use.

FUMIGATING PAPERS

are strips of paper impregnated with substances that give off an agreeable aroma when heated, so that by merely placing a strip of such paper on a hot stoveplate, or holding it over a flame, the entire atmosphere of a room can be scented.

These preparations are divided into two classes: such as are burned, and such as are used more than once. In making the former, the paper is impregnated with a solution of saltpetre before the perfume, whilst the other papers are rendered combustible by steeping with a hot solution of alum, in consequence of which they merely carbonise and do not burn, even when strongly heated.

The following recipes will be found suitable for making these preparations, which have only a slight relationship with cosmetics properly so called:—

The paper is treated with a solution of 100–150 grams of saltpetre in water, and when dry is repeatedly immersed in a strong solution of benzoin or incense, being dried after each immersion.

An excellent fumigating paper may be prepared from the following recipe:—

Benzoin	150	grams.
Santal wood	100	"
Incense	100	"
Grass oil	10	"
Essence of betiver	50	"
Alcohol	1	litre.

For use, the paper is placed in contact with a glowing—not burning—substance, whereupon it immediately ignites without flame, but throwing off brilliant sparks, and diffuses an agreeable perfume.

Another recipe is: Cinnamon oil, 4·0; powdered nutmeg, 4·0; liquid styrax, 4·0; gum benzoin, 4·0; ambergris, 0·3; musk, 0·3. Sufficient alcohol is added, by drops, to form a somewhat fluid mass, which is applied with a large soft brush to one side

of a piece of fine paper, the latter then laid between oil or wax paper. For use, the impregnated paper is held over a flame, so that it merely smokes without taking fire; thus treated, it disseminates an agreeable perfume, and can be set aside for use several times over.

Incombustible fumigating paper is prepared by dipping the paper in a hot solution of 100 grams of alum per litre of water, and then impregnating it with the following mixture:—

Benzoin	200 grams.
Tolu balsam	200 „
Tonka bean essence	200 „
Betiver essence	200 „
Alcohol	600 c.c.

When warmed, this paper diffuses a very agreeable perfume, and can be used a number of times over. It will not ignite, but merely carbonises, even when strongly heated. Some makers produce inferior qualities of these articles by simply dipping the paper, prepared with alum, into melted benzoin or incense.

SUNDRY PREPARATIONS.

Mention may here be made of a number of cosmetic preparations that are in common use in this country:—

Rose Bandoline.—Tragacanth, 52 parts, and rosewater, 950 parts, are digested in a closed vessel for a couple of days at a gentle heat. The mixture is strained, and scented with rose oil; or 1 part of quince seeds is macerated with 40 parts of rose water for several hours, with frequent stirring, and scented with rose oil or other ethereal oil after straining.

Crème de Mauve.—One part of glycerin is mixed with 1 part of alcoholic extract of jasmine, and coloured with a little fuchsin. This preparation is used to impart gloss to the hair.

Oleolisse Tonique de Piver—Five parts of castor-oil are dissolved in 15 of alcohol, and scented with bergamot oil or Portugal oil.

Laurel Tincture, artificial Bay Rum.—Tincture of laurel leaves—1 : 10—186 parts ; laurel oil, 4 ; borax, 30 ; ammonium carbonate, 30 ; rose water, 950 parts.

Extrait Végétal.—Vanilla essence—1 part of vanilla, 30 parts alcohol ; 2 parts of orange-flower extract ; 1 part each of jasmine extract, rose extract, and tuberose extract ; alcohol, 16 ; rose water, 20 parts.

Amandine.—A syrup is prepared from 2 parts of sugar and 1 of water : 16 parts of this syrup are mixed with 4 parts of almond-soap crème until homogeneous, and into this is stirred 400 parts of almond oil scented with 4 parts of oil of bitter almonds, 4 of bergamot oil and clove oil. The mixing of the almond oil with the syrupy mass entails a certain amount of skill ; and some force is required, especially towards the end, as the mass gets stiffer.

Almond-Soap Crème.—This is prepared by melting 500 parts of pure fat and mixing the same with 200 parts of a 25 per cent. solution of caustic potash, the latter being added very slowly and with continued stirring, which is continued until the mass is cold, whereupon it is triturated in a mortar with 12 parts of alcohol and 1 of oil of bitter almonds until frothy and homogeneous.

Almond Paste.—Thirty-two parts of blanched bitter almonds are triturated to a fine paste, to which is gradually added an intimate mixture of honey, 60 parts ; yolk of egg, 30 ; almond oil, 60 ; bergamot oil, 1 ; and clove oil, 1.

Milk of Roses.—Castile soap is dissolved in 4 parts of rose water by the aid of gentle heat, whereupon 1 part of white wax and 1 of spermaceti are added to the solution ; when these are melted, a strained mixture, compounded of 16 parts of blanched sweet almonds and 1000 parts of water, is added by degrees and with constant stirring. Finally, 150 parts of alcohol and 8 of rose oil are incorporated with the mixture.

Frangipani Sachet Powder.—Violet root powder, 1500 ; patchouli leaves, 120 ; santal wood, 120 ; neroli oil, rose oil, santal-wood oil, 2 parts of each ; powdered musk sac, 10 ; and civet, 5 parts.

CHAPTER VIII.

COSMETIC ADJUNCTS AND SPECIALTIES.

IN this section we will deal with certain artificial essences which are frequently used as adjuncts to cosmetic preparations.

The preparation of ionone will be considered first. The odoriferous principle of the violet root cannot be obtained direct by steam distillation, owing to the large percentage of starch present, which apparently retards the escape of the volatile constituents. For this reason Tiemann and Krueger extract the root repeatedly with ether, and then subject the ethereal extract to distillation with steam. This furnishes two products: a muddy residue, which, when treated with alkalis, yields myristic or iridic acid, and a brittle crystalline resin; whilst the distillate contains the ethereal oil together with a larger quantity of myristic acid and its methyl salt, oleic acid and an ester of same, aldehydes, and other substances. The ethereal solution of this mixture is shaken up with a 2 per cent. solution of caustic potash, in order to remove part of the free organic acids. The residual organic acids and esters are more speedily eliminated by repeated fractional distillation of the oil with steam. The neutral oil obtained by this process of distillation is then dissolved in alcohol, and treated with an alcoholic solution of caustic potash at the ordinary temperature. This alcoholic solution is poured into water, the aqueous solution is extracted with ether, and the ethereal solution again distilled with steam, the perfume thereupon passing over with

the first portions of the distillate. The resulting oil is then boiled with an aqueous solution of silver nitrate. The iriske-tone is next clarified by the formation of a phenylhydrazine compound, left to stand for a day, and the hydrazone mixture then subjected to steam distillation, which leaves the iriske-tone behind, in combination with phenylhydrazine, as a brown oil, which can then be decomposed into phenylhydrazine and ionone by dilute sulphuric acid, the ionone being extracted with ether and rectified under a low pressure. Ionone is an oil which dissolves badly in water, but readily in alcohol, ether, chloroform, benzol, or ligroin, and has the sp. gr. 0.939. The aroma of the pure substance is sharp, and, when highly concentrated, quite different to the scent of violets. However, on dilution with alcohol and exposure to the air, it at once exhibits the odour of the natural blossoms.

Artificial essence of violets is obtained by the following method:—

Ten kilos of acetone and 30 grams of citral are dissolved in 100 kilos of benzol, and to the solution is added 5 kilos of a 10 per cent. solution of sodium ethyl. The mixture heats spontaneously and turns red. After twenty-four hours the mixture is shaken up with water, whereupon a portion separates out on the surface of the latter. This portion is then collected and distilled with steam until the smell has disappeared. The residue is boiled with 15 parts of 7 per cent. sulphuric acid, poured out when cold, neutralised, and finally distilled. The result is a pale yellow liquid which in a concentrated state smells like santal, but when considerably diluted assumes a pure odour of violets.

Artificial vanilla essence is prepared as follows:—

3.75 parts (grams) of Peru balsam, 1.75 c.c. of orange oil, 125 grams of violet root essence, 62 grams of powdered Tonka bean, 30 drops of musk tincture, 250 c.c. of rectified spirit, 10 grams of magnesia, 125 c.c. of water. The balsam and oil are

dissolved in 60 c.c. of spirit and triturated with the magnesia, this mixture being then added to the rest of the ingredients, and the whole kept in a warm place for four weeks, to be finally coloured with caramel, and filtered.

Musk Tincture for cosmetic purposes.—Hauer gives the following recipe for this purpose:—

2.5 parts of musk with the musk bag, or 2.0 parts without the bag, and 2 parts of powdered sugar, are placed in a flask and suffused with 10 parts of boiling water containing 0.3 part of sodium carbonate, and are left to stand therein for a day. Hereupon 35 parts of spirit are added, macerated for fourteen days, and then filtered. This process gives a strong tincture of musk, in which the whole of the musk, except the skin, is dissolved.

An excellent eau de Cologne can be obtained by the recipe given by Fehr.

The following ingredients are dissolved in 23 kilos of finest spirit of wine:—

Lemon oil	350 grams.
Bergamot oil	270 „
Finest French lavender oil	20 „
Peppermint oil	12 „
Acetic ether	12 „
Neroli oil	120 drops.
Best French oil of thyme	100 „
Best French rosemary oil	100 „
Rose oil	20 „

After a few days are added—

Triple orange-flower water	1100 grams.
Triple rose water	200 „

This preparation furnishes a successful imitation of the renowned product of J. M. Farina, “opposite the Julichsplatz,” Cologne. If the predominant odour of rosemary oil be objected to, the amount of this oil may be diminished. After keeping the preparation for three months, it will need

an addition of 5-7½ kilos of spirit, after which it must be stored for another three months before being sold.

Eau de Cologne, in any case, should be kept for a year before it is sold; and under these conditions inferior recipes even will furnish really good qualities of this scent.

COLOURING COSMETIC PREPARATIONS.

For this purpose the employment of betulin has recently been proposed.

If the roots of *Betula alba* be boiled with an alkali, and the filtered, dark-red decoction be treated with hydrochloric acid, a precipitate—up to 20 per cent.—is obtained, which, when well washed and carefully dried, furnishes a reddish brown powder, with an odour of leather and a faintly bitter taste. This product is known as betulin, and, when dissolved, forms an excellent colouring matter for cosmetic preparations.

The best method of procedure is to triturate 0.05 gram of betulin in a porcelain mortar, under gradual addition of 5 grams of 90 per cent. alcohol, containing a little glycerin, until complete solution is obtained. The product is made up to 20 grams with distilled water, and thus gives a handsome red-brown liquid, which may be further diluted with water, though the larger the proportion of alcohol the better the colour. A slight addition of alcohol increases the tinctorial power, whereas acid has the opposite effect, and, if added to a solution already extensively diluted with water, will throw down the colouring matter. Nitric acid precipitates the colouring matter as a violet-red deposit, the same colour being also assumed by the alcoholic solution. The colouring matter is not precipitated by organic acids, but it is thrown down by aqueous solutions of sulphate of quinine and lead acetate, whilst ferric chloride gives a greenish black coloration.

With regard to other colouring matters, the following remarks may be made:—

For all alcoholic cosmetic preparations and such as contain glycerin, the aniline dyes are the best colouring matters; on the other hand, they are unsuitable for such cosmetic articles as contain fat, whether animal or vegetable fat is immaterial. Of course the aniline dyes used must be guaranteed pure, and especially free from arsenic; a guarantee which the large firms of colour makers will certainly be able to give.

Yellow colouring matters for this purpose are saffron, turmeric, and palm oil.

Saffron gives a dark yellow or orange colouring material, which is readily soluble in alcohol, and also in petroleum ether, and is absorbed by fats. The finely comminuted saffron is preferably digested with petroleum ether, the solvent being then removed by distillation, and the final traces allowed to escape spontaneously; this method furnishes the pure colouring matter, which is easily miscible with fat. The colouring matter of saffron can also be prepared by digestion with melted lard or almond oil.

Turmeric, the yellow colouring matter of curcuma (turmeric) root, is recovered by extracting the root with alcohol or petroleum ether in the same manner as saffron. It is unsuitable for preparations containing alkalis, since these substances turn the colour brownish red.

Palm oil imparts its yellow colour to the soaps prepared from this fat; but the colouring matter is completely bleached on exposing the soap in a damp state to the air for some time.

The chief red colouring matter is carmine, and this is best prepared by treating finely powdered cochineal with ammonia, adding a little dissolved alum, and exposing to the air and direct sunlight, whereupon the colouring matter will be deposited in the form of red flakes, which are collected and dried.

Carthamine red is contained in the flowers of the marigold.

There are two colouring matters, one yellow and one red. The former is extracted from the dried flowers by means of water, the red colouring matter being then recovered from the residue by treating the latter with a weak solution of soda. On carefully diluting this solution with acetic acid, a precipitate of safflower red is obtained, and, when dried, forms a mass exhibiting a metallic greenish lustre, and furnishing a red powder, which is used for Rouge en Feuilles or Rouge en Tasses.

Alkanna is obtained from the root of the same name, the red colouring matter being extracted with petroleum ether; it is also soluble in melted lard or warm oils.

Ratanhia root yields a reddish brown colouring matter which is soluble in alcohol, and can be recovered by extraction with that solvent. In the preparation of tooth tinctures and mouth washes, for which ratanhia is a favourite colouring agent, use is often made of finely grated red santal wood or Pernambuco wood, both of which, in addition to their content of astringent matters,—tannins,—also yield a handsome colouring matter to alcohol.

Green colours are produced by the aid of chlorophyll, which can be easily prepared by treating triturated leaves with alcohol, and evaporating the solvent. In preparing green powders, use is often made of dried and pulverised leaves of a brilliant green colour, among those most suitable being spinach, laurel, celery, and parsley leaves. Soaps are coloured green by a mixture of yellow and blue, a yellow soap being melted and treated with an addition of finest smalt and ultramarine until the desired shade of green is produced. Indigo-carminé must not be used for this purpose, as it would stain the skin blue when the soap came to be used.

Smalt or ultramarine is used for producing blue in many cosmetic preparations, but both are insoluble. The only soluble blues are aniline blue and indigo-carminé, but the latter

can only be used for colouring pomades. It is prepared by suffusing perfectly dry, finely powdered indigo with fuming sulphuric acid, and, at the end of twenty-four hours, adding finely powdered chalk until effervescence ceases, after which the preparation is filtered.

Violet is produced by suitably mixing a red and a blue colouring matter.

Brown is imparted to cosmetic preparations by means of caramel, produced by heating sugar in an iron pan until converted into a deep black mass, that has a brown appearance only when viewed in thin layers. Caramel is soluble in water, but not in alcohol; it may also be employed for colouring soap.

Black is produced almost exclusively by finely pulverised vine charcoal or ivory black. Liquids are coloured black by the aid of Indian ink.

COSMETIC SPECIALTIES.

The manufacture of special preparations seems to be a very profitable business in all branches where it can be carried on, provided that the products themselves are perfectly unobjectionable in every sense, and that they are judiciously advertised—an absolutely indispensable condition under the circumstances of modern commerce. And in the case of cosmetic specialties—which in most instances seem to have developed into hygienic preparations—the prospects of a good reception and ready sale appear to be particularly bright.

As in the case of general cosmetics, the main question in connection with specialties is whether the preparations themselves are capable of fulfilling the objects for which they are produced; and if so, to what extent. It is therefore necessary to discover, in addition to ingredients that serve to improve either the external appearance or the mechanical perfection of the product, such substances as may lay claim to specific action.

If the preparation be deficient in ingredients of this latter class, or if their proportion be incorrectly selected, then the specialty in question will never meet with the desired reception, and will remain for the most part entirely inert.

Another point on which great care is necessary, is to see that none of the ingredients used are likely to have any action injurious to the health of the consumer; and in this respect, unfortunately, some cosmetics, especially those of a secret character, often leave much to be desired. In this connection, the manufacturer must possess a thorough knowledge of the properties of his ingredients, since the lay consumer is not in a position to judge whether the addition of this or that substance is injurious or not. The consumer, relying on the knowledge, skill, and experience of the manufacturer, makes use of the preparation in the manner and to the extent indicated in the printed instructions accompanying the package.

In considering the description and preparation of the specialties mentioned in the following pages, it will frequently be necessary to adopt a critical attitude in order to obtain a clear insight into the value and import of some among them. Such methods of preparation as are simple, and conducted in a matter-of-course manner, will be passed over briefly, fuller consideration being reserved for special methods, wherein any doubt might arise as to the right course of procedure, and where any lack of skill might endanger the success of the preparation.

Finally, it is perhaps almost superfluous to mention that, in the present age of luxury, very peculiar care should be bestowed on the manufacture of cosmetic specialties. Happily, the high state of development attained by the paper, glass, porcelain, and metal industries facilitates our task, so that, owing to the large available selection of tasty materials for packing, the only limitations imposed on the maker are those of personal taste,

the appreciation of the article by the public, and—by no means least—the price at which the preparation can be sold.

First consideration will be devoted to

SPECIALTIES IN HAIR WASHES.

The premier place in this class must be accorded to hair dyes, which are largely prepared as specialties. Here the main point is to be sure that no ingredients capable of producing injury to health are present.

Broers has made experiments with various innocuous dyes for the hair. The employment of sodium sulphide and bismuth subnitrate imparts a hateful grey appearance to the hair. Potassium permanganate, in the condition of a 1 per cent. aqueous solution, forms a fairly powerful brownish blonde dye.

A 5–10 per cent. aqueous solution of walnut shells or walnut leaves has no effect, neither is Indian ink of any use, the colour being removed by washing with water.

On the other hand, better results are furnished by henna, the leaves of *Lawsonia inermis*. The powdered leaves are stirred up to a pulp with water, applied to the hair with a brush, and afterwards washed off again. The longer the henna acts on the hair, the darker is the dye; the colour varies from light brown to orange red. After an exposure to the dye for forty-five minutes, the hair becomes pale brown; after four hours it is red-brown to golden blonde,—a shade greatly in favour among the ladies,—and after twenty-four hours is orange red and unnatural looking. This possibility of regulating the colour by the time of the exposure, renders henna specially adapted for use as a hair dye.

Notwithstanding the unfavourable results mentioned above as having been obtained by the use of the harmless ingredient, walnut shells, a few recipes in which these appear are now given.

HAIR DYE FROM WALNUT SHELLS.

Green walnut shells	450 grams.
Powdered alum	30 "
Rose water	120 "

The ingredients are triturated together in a mortar, pressed, and treated with 90 per cent. alcohol in the proportion of 30 parts per 100 of liquid, the mixture being then left for four days in a closed vessel, and finally filtered and scented to taste.

WALNUT HAIR OIL.

Green walnut shells	120 grams.
Powdered alum	15 "
Olive oil	600 "

trituated in a mortar, then warmed on a sand bath, with continued stirring, until all the moisture has been dissipated, and afterwards pressed, strained, and perfumed to taste.

WALNUT POMADE.

Green walnut shells	450 grams.
Powdered alum	60 "
Olive oil	720 "
Palm oil	120 "
White wax	90 "

warmed on a sand bath, with stirring, until all moisture has been driven off, then strained, and, when nearly cold, mixed with 300 parts of pomade, previously melted on the water bath.

EXTRACT OF WALNUT SHELLS.

The freshly peeled shells from green walnuts are lixiviated with ammoniacal water,—2 : 1,—the menstruum being concentrated to the consistence of syrup, then diluted with suitably scented spirit,—2 : 1,—and, if necessary, treated with an addition of 2 per cent. of pyrogallic acid, the latter increasing the

rapidity of the dye. Before applying this dye, the hair must have been freed from grease and thoroughly dried.

Another recipe runs as follows:—The green outer shells, which can be obtained cheap at the time the walnuts ripen, are pounded with a pestle and then suffused with water containing 1 per cent. of salt. After three days the whole is poured into a large pan, on which a mark is then made to show the height of the liquid, it being necessary to replace the water lost by evaporation. The liquid is heated to near boiling point for four to six hours, then left to cool down, and well pressed. In the absence of a press, this may be done with the aid of a fine linen cloth, or preferably a sack of flax canvas, about 40 inches long and 20 inches wide, which is half filled with the mass from the pan, held over an open vessel, tied up at the mouth, and twisted by means of a couple of sticks, two men being required for the operation. Nevertheless, the bag must not be twisted so tightly as to cause it to burst. The liquid thus obtained is returned to the pan, and is concentrated to one-fourth its initial volume, which is measured for that purpose, the best plan being to place a quantity of water equal to one quarter of the liquid in the pan, marking the level, pouring out the water again, and then allowing the nut liquor to evaporate until the level of the mark is reached. The finished extract then receives an addition of 16 per cent. of 95 per cent. alcohol, and is either stored in tightly closed vessels, for stock, or finished off, ready for use, by the addition of some perfume.

LANOLIN ROSE CREAM.

This preparation for the hair is compounded from lanolin, 25; almond oil, 60; potash soap, $7\frac{3}{4}$, melted together, stirred up with 150 parts of water and 30 of rose water whilst cooling, and finally scented with essence of roses.

A HARMLESS HAIR DYE.

An innocuous dye, to be applied with the brush morning and evening after using an alkaline wash, is composed of—

Pyrogallic acid	.	.	.	.	.	4	grams.
Citric acid	.	.	.	.	.	0.3	„
Boroglycerin	.	.	.	.	.	10	„
Distilled water	.	.	.	.	.	90	„

AUREOL.

Aureol for dyeing the hair has proved itself a thoroughly harmless preparation, is very stable, and the colour is not altered, even by soap. It consists of 1 per cent. of methol, 0.3 per cent. of amidophenol chlorate, and 0.6 per cent. of monoamidophenylamine, all dissolved in 50 per cent. of alcohol containing 0.5 per cent. of sodium sulphite.

The hair must be first washed with soap or soda, to remove all grease. The aureol is then mixed with an equal quantity of hydrogen peroxide, and applied, by means of a fine comb, to the still damp hair. At the end of two to three hours the hair assumes a handsome dark brown colour, which can be made still darker by repeating the application as desired.

The so-called Turkish Water for dyeing the hair is prepared by kneading finely powdered gall nuts with a small quantity of oil to form a homogeneous paste, which is then roasted in an iron drum until the evolution of oil vapours has ceased. The residue is taken up with a little water and kept over a fire until perfectly dry. The mass, very slightly moistened, is then carefully mixed with a metallic powder and kept in a damp place, where its dyeing properties are gradually developed. The mixture is sometimes perfumed with ambergris or some essential oil.

To employ this preparation for dyeing the hair, eyebrows, eyelashes, or whiskers, a small quantity is crushed in the palm

of the hand, or between the fingers, and carefully rubbed on the hair, etc. At the end of a few days a fine black colour will be developed, the hair remaining soft, glossy, and retaining its colour for a considerable time, even after a single application.

The tinctorial properties of this preparation are evidently largely due to the pyrogalllic acid formed during the roasting of the gall nuts.

AMERICAN HAIR WASHES.

The popular American hair washes—shampoos—may be prepared from the following recipes:—

1. The white of two eggs is beaten to a froth, mixed with 150 grams of water and 90 grams of ammonia, and finally shaken up well with 10 grams of eau de Cologne and 120 grams of alcohol.

2. Finely powdered borax 60 grams and 40 grams of glycerin are triturated, mixed with 400 grams of rum and 300 grams of bay rum by stirring or shaking, and then incorporated with the whites of two eggs previously beaten up to a froth.

3. Ammonia, 15 c.c. ; glycerin, 30 c.c. ; soap spirit, 30 c.c. ; 95 per cent. spirit, 60 c.c. ; bergamot oil, 20 drops, and sufficient water to make a total volume of 500 c.c.

4. Ammonia, 30 c.c. ; cantharides tincture, 15 c.c. ; Paprica tincture, 15 c.c. ; 95 per cent. spirit, 250 c.c. ; water, 250 c.c. ; bay rum, 500 c.c.

5. Ammonia, 15 c.c. ; 95 per cent. spirit, 125 c.c. ; a little scent, and enough water to make 250 c.c.

6. Alcohol, 50 c.c. ; sulphate of quinine, 1 gram ; cantharides tincture, 8 grams ; borax, $3\frac{1}{2}$ grams ; ammonia, $3\frac{1}{2}$ grams ; glycerin, 100 c.c. ; caramel solution, $3\frac{1}{2}$ grams, and sufficient bay rum to make 500 c.c. This preparation is popular under the name "Compound Quinine and Glycerine Hair Tonic."

7. Castor-oil, 30 c.c. ; bergamot oil, 40 drops ; glycerin, 100 c.c. ;

cantharides tincture, 12 c.c.; ammonia, 12 c.c., and sufficient water to make 500 c.c. The oil is dissolved in the alcohol, mixed with the tincture, and the mixture of ammonia and glycerin is gradually run in. The liquid is called "Eclectic Hair Tonic."

8. Sulphate of quinine, 1.75 c.c.; alcohol, 55 c.c.; cantharides tincture, $3\frac{1}{2}$ grams; capsicum tincture, $1\frac{3}{4}$ grams; ammonia, 15 c.c.; glycerin, 30 c.c.; with sufficient bay rum to make 175 c.c. This preparation is known as "Quinine Hair Tonic."

Shampoo powder is a mixture of ammonium carbonate, 1; borax, 1; and Quillaya bark, 2 parts, with bay-leaf oil or some other perfume. It is made up in packets, with the following directions for use: "Put the powder into a covered jug along with a cupful of boiling water, close the lid, and leave the liquid to stand awhile until cool enough to use."

A popular specialty is Hair Curling Essence, though its efficacy is doubtful. The preparations mainly rely on the adhesive properties of gum, or on the removal of grease from the hair, followed by curling. The following recipes are recommended:—

HAIR CURLING WASHES.

1. Purest potash, 7 grams; ammonia, $3\frac{1}{2}$; glycerin, 15; alcohol, 12; rose water, 550, with a little scent. The loosely tied or loose hair is moistened with this liquid, and will curl on drying.

2. Borax, 100 grams; gum arabic, $3\frac{1}{2}$ grams; hot water, 1 litre; spirit of camphor, 45 c.c. The hair is moistened with this preparation at night.

3. Potash, 7 grams; ammonia, $3\frac{1}{2}$; glycerin, 15 grams; rose water, $\frac{1}{2}$ litre.

4. Gum arabic, $3\frac{1}{2}$ grams; sugar, $3\frac{1}{2}$ grams; rose water, 50 c.c. The hair is moistened with this liquid before retiring to rest, and is then done up in curl-papers.

5. 175 grams of tragacanth are digested, with frequent stirring, for forty-eight hours in $4\frac{1}{2}$ litres of rose water, then

strained through a linen cloth, left for a few days to clarify, and scented with rose oil.

6. Dry potash, 7 grams; cochineal, $1\frac{3}{4}$ grams; ammonia, $3\frac{1}{2}$ grams; rose spirit, $3\frac{1}{2}$ grams; glycerin, 7 grams; rectified spirit, 75 c.c.; and distilled water, 500 c.c., are digested, with stirring, for a week, then decanted and filtered. The hair is moistened with this liquid, and then loosely tied up; the effect appears after drying.

7. Potassium carbonate	12.0 grams.
Ammonia	7.5 „
Glycerin	7.5 „
Rectified spirit	50.0 „
Water sufficient to make	600.0 „

The hair is moistened with the liquid, and then dried, whereupon it curls. All grease must have been removed from the hair beforehand by washing with soap.

M. Joseph recommends, as a harmless and simple hair dye, the use of a 1–20 per cent. argente vaseline, *i.e.* an ointment prepared from yellow vaseline with an addition of 1–20 per cent. of silver nitrate, according to the shade to be produced,—the silver salt having been dissolved in a minimum of water. This preparation is applied, by means of a comb, every eight to fourteen days.

A very profitable branch of specialties is that of—

PREPARATIONS FOR PROMOTING THE GROWTH OF THE HAIR.

A preparation of this class which effectually counteracts the tendency of the hair to fall out, and can be confidently recommended, has been given by Dieterich, the recipe being as follows:—

Chloride of quinine, 0.4; tannin, 1.0; alcohol, 80.0; cantharides tincture, 1.0; glycerin, 60.0; eau de Cologne, 4.0; vanillin, 0.01; and santal-wood powder, 0.005 gram. This liquid is left to stand for five days and then filtered. It should be employed to wash the hair every other day.

The santal-wood powder may very well be omitted, in which case the long standing and the filtration are rendered unnecessary.

A hair pomade for the same purpose, and one that has proved efficient in preventing the hair falling out in cases of nervous headache, has been recommended by Leistikow. The composition is as follows:—Cantharides tincture, 3·0; chloral hydrate, 2·0; lanolin, 5·0; vaseline, 10·0; cherry-laurel water, 10·0; and limewater, 10·0 grams. The chloral hydrate is dissolved first, the lanolin and vaseline liquefied by warmth, and the liquid ingredients added by degrees, and incorporated by constant stirring and mixing.

Another recommended specialty for the hair is

SCURF SOAP.

One hundred grams of *Sapo viridis* are dissolved, by the aid of warmth, in a mixture of 15 grams of glycerin, 50 grams of eau de Cologne, and 50 of rectified spirit, 3 grams of B-naphthol being added to the whole. The head is washed with this soap every second or third day.

SPECIALTIES FOR THE BEARD AND WHISKERS.

Pomade dye.—Twenty parts of stock soap are carefully melted on the water bath so as to prevent lathering, and then mixed with 3 parts of mucilage—1 part of gum arabic to 2 of water. To this is added the colouring matter, consisting, for brown whiskers, of mahogany-brown dye with a little vine black, or the latter alone for black beards, the whole being afterwards scented with a mixture of 20 parts of geranium oil, 10 of bergamot oil, 15 of clove oil, and 2 of cedar-wood oil. The preparation is cast into a mould, then cut up into small squares and packed in boxes lined with tinfoil; each box contains also a small brush for applying the pomade, and a looking-glass in the lid.

Tube pomade.—This is prepared in the following manner :—1000 of white wax, 500 of soap powder, 500 gum arabic, 625 of water, 20 of bergamot oil, 10 of sweet-orange oil, 2 drops of rose oil—to $2\frac{1}{2}$ kilos.—The gum arabic is dissolved in the water, strained, and the mucilage added to the melted wax, the mixture being warmed up and mixed with the soap powder, the scent being added last of all.

A noteworthy cosmetic, suitable for the production of specialties, is—

GLYCERIN-GELATINE.

To prepare this article, $8\frac{1}{2}$ grams of gelatine are left to soak in 180 grams of rose water, then dissolved on the water bath, and mixed with 20 grams of egg albumin when cold. On heating the preparation up again, the solution clarifies, owing to the coagulation of the albumin; and it then receives an addition of 0.75 gram of salicylic acid dissolved in 180 grams of glycerin, to serve as a preservative. The whole is filtered through a hot-water filter, and packed, while warm, into wide-necked flasks, labelled "Glycerin Cream."

BRILLIANTINE.

This name is applied to cosmetics for glossing and stiffening the hair of the beard and forehead-fringe. The following recipes are suitable :—

1. Castor-oil, 6 grams ; Venetian soap, 2 ; benzoin, 2 ; alcohol, 200, scented with a little rose oil or neroli oil.
2. Glycerin, 10 ; alcohol, 100 ; rose water, 100.
3. Castor-oil, 6 ; glycerin, 6 ; benzoin, 2 ; alcohol, 200, scented with a little rose oil or neroli oil.
4. (Solid Brilliantine), finest olive oil, 1 kilo ; spermaceti, 300 grams ; bergamot oil, 25 ; clove oil, 29 ; French geranium oil, 10 grams.
5. Glycerin, 6 ; castor-oil, 6 ; benzoin, 2, dissolved in 200 strong spirit. A little acetic ether is added, with suitable scent.

6. Melted white paraffin, 90; castor-oil, 10; Essence triple de Reseda—or another—30; 93 per cent. alcohol, 60; sufficient rose water or neroli water is then added to make the two liquids about equal.

7. Lanolin, 120 grams; spermaceti, 60; castor-oil, 350, are melted together, and, when cold, mixed with bitter-almond oil, 0.06 gram; clove oil, 0.6; and bergamot oil, 1.29 grams.

8. Spermaceti, 250 grams; almond oil, 1250 grams; lemon oil, 100 grams. The spermaceti is dissolved in the almond oil and mixed with the lemon oil, the whole being poured into hot vessels, which are allowed to cool very slowly in order to produce crystallisation.

9. Honey, 30 c.c.; glycerin, 15 c.c.; alcohol, 50 c.c.

Mouth washes are also a suitable article for specialties, a particularly profitable preparation being a

FROTHING MOUTH WASH

compounded from powdered Quillaya bark, 50 grams; glycerin, 45 grams; sodium salicylate, 7 grams; bergamot oil, $1\frac{3}{4}$ grams; oil of wintergreen, $1\frac{3}{4}$ grams; clove oil, 10 drops; alcohol, 30 c.c., with sufficient carmine solution to colour the preparation, and enough dilute alcohol to make it up to 500 c.c. The powdered Quillaya bark is first macerated with dilute alcohol and glycerin and strained, the filtrate being added to the ethereal oils dissolved in the alcohol. The mixture is next coloured with the carmine solution, shaken up with a little powdered talc, filtered until clear, and made up to the prescribed volume (500 c.c.) with dilute alcohol.

Equally to be recommended is the manufacture of

MOUTH WASH TABLETS,

which meet with a ready sale, and seem worthy of attention. Bernegau gives the following recipe for this article:—Heliotropin, 0.01 gram; saccharin, 0.01 gram; salicylic acid, 0.10 gram;

menthol, 1 gram ; milk sugar, 5 grams ; and sufficient alcohol (scented with rose oil) to make 100 tablets. The mass may be coloured red with eosine dye, green with chlorophyll, or blue with indigo-carmin.

Another specialty which is profitable, though not very closely related to cosmetics properly so called, is

NERVE PENCIL.

Mayet gives the following recipe :—Chloral hydrate, 1 part ; menthol, 1 part ; cocoa butter, 2 parts ; spermaceti, 4 parts. The pencil is applied to the affected part, or, in case of headache, rubbed on the forehead, and generally effects a complete cure, or at least considerably reduces the pain. The method of preparation is to melt the cocoa butter and spermaceti, then add the menthol and chloral hydrate, and cast the whole in moulds.

Among the class of rouge preparations, which has already been treated under general cosmetics, there are two which are suitable for specialties, namely—

COSMETIC FOR THE HANDS AND VEINS.

That for the hands is a concentrated liquid preparation, obtained by intimately mixing 100 grams of finest pure zinc white with fine talc, 100 grams ; glycerin, 70 ; benzoin tincture, 20 ; heliotropin extract, 10 ; and rose water, 270 grams. However, unless coloured with a little carmine, the preparation imparts merely a rather cadaverous appearance to the hands, when viewed by artificial light particularly. A very good modification is that obtained from carmine, 0·4 ; talc, 30 ; precipitated chalk (which may be partly replaced by zinc white), 70 ; benzoin tincture, 40 ; and distilled water, 500–700, according to the price.

The vein cosmetic, which is chiefly used as a theatrical make-up, is compounded of white bole, 80 ; zinc white, 10 ; Berlin blue, 10 ; and gum arabic, 3 parts, mixed to a plastic mass with

orange-flower water, and made into sticks about $\frac{1}{4}$ inch in diameter.

Another class of special preparations in large demand, and which, if properly made, should be saleable to great advantage, besides being capable of producing the desired effect, is that of

DEPILATORIES.

In the first place, it must be premised that depilatories cannot prevent the re-growth of hair on places whence it has been removed by their action; on the contrary, according to the author's experience, depilatories produce exactly the same result as shaving, the aftergrowth of the hair being accelerated. Nevertheless, depilatories are in large demand, ladies having a great objection to hairy growths on the face and arms. Numerous recipes for these preparations exist, but they are by no means all of equal value; furthermore, it should be borne prominently in mind that no substances capable of corroding or otherwise attacking or injuring the skin are permissible in depilatories. This point is the more important in view of the fact that, as already mentioned, depilatories require to be constantly employed when once their use is begun. The following recipes will be found best suited for the purpose in view, and also financially advantageous to the maker:—

1. Quicklime slaked with a little water, so as to furnish a friable powder, is stirred up to a thin pulp with a further quantity of water, and the mixture is saturated with a current of sulphuretted hydrogen gas. Ten grams of the resulting calcium sulphide are mixed with 5 grams each of starch and starch sugar and a little oil of lemon, to form a thick pulp, which is applied to the seat of the hairs and left in contact for twenty minutes, after which it is scraped off with a paper knife.

2. A practical and cheap depilatory can be prepared in the following manner:—Powdered barium sulphide, 1 part; precipitated chalk, 1 part. It is essential that both these

ingredients should be perfectly dry and in a finely powdered condition, in order, on the one hand, to prevent the mixture becoming prematurely exhausted, and, on the other, to obviate irritation that might arise from the granular or sandy condition of the depilatory. If these precautions be observed, the product will consist of a very fine, light grey powder, which, on being made into a pulp with a little water, will act on the hair in three to four minutes, without any injury to the skin. It is also very cheap to produce. It is erroneous to suppose that the barium sulphide cannot be procured ready made, since the same can be obtained in any quantity from such firms as Schering, J. D. Riedel, T. Teichgraeber, J. G. Braumueller & Sohn, of Berlin, and others.

3. Barium sulphide, 10 parts ; zinc oxide, 5 ; and wheat starch, 5 parts, are triturated to a very fine powder, which is made up into a thick pulp with water before use, and is applied for about ten minutes.

The preparation of toilet powders has already been described, and, for the purposes of the present section, namely, specialties, it will be sufficient to deal with

FACE POWDER

alone, in connection with which article the author's experience tends to show that the best of all face powders, and one calculated to prevent or remedy all impurities of the skin, is compounded solely of starch, talc, and salicylic acid.

In the subjoined recipes it is highly probable that the violet root therein mentioned could be advantageously replaced by some other ingredient, since this powder can never be obtained in such a finely divided state as the other constituents.

Violet powder.—Zinc white, 100 grams ; talc, 200 ; violet root powder, 200 ; rice meal, 500 ; extrait de Jasmin, 15 ; bergamot oil, 0·5 ; rose oil, 0·59 ; ylang-ylang oil, 1 ; musk tincture,

5 (the latter prepared from musk, 20 grams; rose water, 250 grams; alcohol, 2 litres).

Rose powder.—Rice meal, 3 kilos; carmine, 4 grams; rose oil, 8 grams; santal, 8 grams.

White powder.—Rice meal, 2 kilos; bismuth white, 250 grams; citronella oil, 4 grams; bergamot oil, 4 grams; geranium oil, 2 grams.

Ordinary powder.—Rice meal, $2\frac{1}{2}$ kilos; talc, $2\frac{1}{2}$ kilos; magnesia, 80 grams; geranium oil, 13 grams.

Poudre de Riz.—Magnesia, 500 grams; talc, $4\frac{1}{2}$ kilos; rice meal, 10 kilos; geranium oil, 200 grams; carmine, 20 grams; the whole scented with bergamot oil, 30 grams; ylang-ylang oil, 10 grams; neroli oil, 10 grams; and eau de Cologne, 200 grams.

A not unprofitable specialty, that can easily be produced as a subsidiary article on a small scale, is

SOAP POWDER.

The following method is perhaps the best, since no mill is required:—

1. Silver soap, or grain soft soap	25 parts.
Linseed oil soap	20 "
Calcined soda	65-70 "

crutched together, whereupon the mass becomes warm; it is then turned over with the shovel at short intervals, until it falls into small pieces. When quite cold it is passed through a fine sieve, and the powder is finished.

2. Olein	35 parts.
40° Be. soda lye	14 "
Calcined soda	7 "
Water	28 "

The mixture is left to saponify, and forms a clear paste, which is boiled once. The fire is then drawn, and 35 parts of calcined soda are crutched in, until a granular mass is formed,

and this is stirred till cold. If the powder is to be scented, or, as has latterly been practised, is to be endowed with an odour of ammonia or turpentine, the scent—generally essence of mirbane—or either ammonia or turpentine is added; when cold the product will be obtained in the state of a very solid mass, which has to be reduced to powder by grinding and sifting.

A cheaper powder can be prepared, in exactly the same way, from—

Palm kernel oil	.	.	.	.	.	13·5 parts.
Palm oil	.	.	.	.	.	2 „
40° Be. soda lye	.	.	.	.	.	10 „
Water	.	.	.	.	.	80 „

to which are added 105 parts of calcined soda, for producing the paste.

A very profitable article, and one sure of a good sale, if only well advertised, is a

WATER SOFTENING PREPARATION,

composed of equal parts of anhydrous soda and slaked lime. For use, a small teaspoonful is added to 10 litres ($2\frac{1}{4}$ galls.) of the water to be softened, and the whole left to settle a short time.

Although the ratio between the equivalent proportions of lime and soda is about as 1 : 2, trials made with this preparation show that, at the ordinary temperature, the water is obtained free from lime, the reaction proceeding slowly and there being always still a little soda present; but when used in the warm, the water always remains somewhat calcareous.

An article in considerable demand is

SHAVING CREAM.

Although soap creams have been on the market for some years, complaints are constantly arising that the cream has a

rancid smell and irritates the skin ; but at length, after numerous attempts, a cream has been prepared free from these defects, and fulfilling all requirements. For the manufacture of this article a water bath is indispensable. The following ingredients are saponified in an enamelled pan, which is set in a larger vessel containing hot water :—Perfectly fresh lard, 15 kilos, and sesame oil, 3 kilos. The fat is raised to a temperature of $37^{\circ}\cdot5$ C., and must be maintained thereat by suitably heating the water bath. Seven kilos of 40° Be. potash lye, $1\frac{1}{2}$ kilos of 40° Be. soda lye, and $\frac{1}{2}$ kilo of water are then weighed out separately, the soda lye being added first to the fat in a very thin stream, the potash afterwards, and both well stirred in until the mass is so thick that the stirrer will stand upright therein. This may take as much as a day to accomplish, but is the only way in which proper saponification can be secured. The cream must not be allowed to cool during the operation, but has to be kept warm by means of the water bath. The finished cream must be filled into stone jars and stored in a cellar. A fine, but expensive, perfume for this preparation consists of pure bitter-almond oil, 250 grams ; pure Turkish-rose oil, 50 grams ; cumarin, 2 grams. This scent must be added to the cold soap cream in a mortar shortly before sending out, or before use, and is incorporated by an industrious mixing and pounding.

A glycerin cold-cream soap is prepared as follows :—

Cocoonut oil, 30 kilos ; tallow, 20 kilos ; crude palm oil, 2 kilos, are warmed to 75° C., and into the mixture is stirred 30 kilos of 36° Be. soda lye, 2 kilos of glycerin, and $\frac{1}{2}$ kilo of spermaceti (the latter warmed to the same temperature as the fat). The soap is then covered up and left to get hot spontaneously, which it will do in about an hour and a half. The product is a clear, solid, yellow soap, greatly resembling a settled curd soap. It is put into the frames, and scented with cassia oil, 50 grams ; bergamot oil, 50 ; lavender oil, 120 ; musk

tincture, 60 ; benzoin tincture, 80 ; oil of wintergreen, 20 ; clove oil, 80 ; and cinnamon oil, 10 grams.

WASHING CREAMS.

1. Prime white soft soap, 320 parts ; pulverised olive oil soap, 80 ; sesame oil, 20 parts, are well purified, and scented with 5 parts of oil of lemon rind.

2. Potash soft soap, 250 parts ; best soda soap, 120 ; olive oil, 60 ; water, 60 ; potash, 7 parts.

3. Oil soap, 60 parts ; dry soap powder, 30 ; honey, 15 parts ; with sufficient rose water to produce a fine lathering cream.

4. Eight parts of lard and 2 of cocoanut oil are saponified with $4\frac{1}{2}$ parts of 40° Be. potash lye on the water bath, coloured rose-red, and scented with rosewood oil and bergamot oil.

5. 30 parts of best lard and 6 parts of sesame oil are melted, and into the mixture, at a temperature of 37°·5 C., is crutched a mixture of 3 parts of 40° Be. caustic soda lye and 1 part of water, run in in a thin stream, this being followed in the same way by 14 parts of 40° Be. caustic potash lye. The soap mass is maintained at the above temperature by the aid of a water bath, and is stirred until complete saponification is secured.

HYGIENIC SOAP.

is a specialty that should not be underestimated. The first preparation of this class to be mentioned is—

Marble-dust soap, which is prepared by liquefying household soap and mixing with a threefold volume of sifted marble dust. An addition of 4 per cent. of lysol is also recommended. Even without this addition, the soap in question fulfils its purpose of cleansing the hands by mechanical means, dispensing with the use of brushes, which are objectionable from a hygienic point of view.

Nicotine soap.—Mentzel of Bremen has prepared a nicotine soap for use in cases of skin diseases. It contains 10 per cent.

of tobacco extract, which is equal to about 0·7 per cent. of nicotine. The use of this soap, which is brownish black in colour, and is scented with bergamot oil, is principally indicated in cases of parasitic skin diseases, especially those produced by animal parasites. Taenzer has successfully used nicotine soap, both in cases of scabies and several instances of pityriasis and parasitic eczema. It is unsuitable for moist eczemas and pustulous affections; but is recommended for the alleviation of itching. No unfavourable subsidiary action has been observed.

SOAP ESSENCES.

These are coloured or uncoloured scented alcoholic solutions of soap. The alcohol must be strong, otherwise the solutions will not be clear; 80 per cent. alcohol seems to be the best. Tallow and lard soaps are not suitable for the preparation of these articles, since they tend to remain opaque on cooling; and white olive oil soaps are best for the purpose. Potash plays a favourable part in the production of these essences; nevertheless an excess of the alkali must be carefully avoided. The most usual formula is—White olive oil soap, 200 parts; 80 per cent. alcohol, 1000 parts; pure potassium carbonate, 12 parts. The soap is cut into small strips, and introduced into a closed vessel, placed in a water bath. The alcohol and potash are then added, and the bath is slowly heated, stirring being practised without interruption. When complete solution has been effected—in about forty-five to sixty minutes—the vessel is removed from the water bath, and the scent is added. The soap may be coloured yellow or red, by saffron or rosaniline respectively, and is then left to settle, being afterwards decanted, or preferably filtered with care. Rabiquet recommends the following recipe:—White soap, 1 part; 85 per cent. alcohol, 3 parts; distilled water, 1 part. A special detergent preparation may be obtained by mixing—White soap, 2 parts; 80 per cent. alcohol, 4 parts; soda crystals, 1 part.

SMELLING BOTTLES.

English smelling salts are prepared by mixing 2 parts of coarsely powdered sal-ammoniac with 1 part of pure potassium carbonate in a mortar or on paper, the smelling bottle being nearly filled with this mixture, which is covered by a plug of cotton-wool or asbestos. A little scent, according to the taste of the purchaser, is dropped on this, and the bottle is closed with a stopper.

French smelling salts.—The stoppered smelling bottle is half filled with strong, suitably scented glacial acetic acid, and sufficient loose asbestos, or glass wool, is added to prevent the contents running out when the bottle is reversed. The scent used may be lavender oil, bergamot oil, cinnamon oil, clove oil, or thymol and eucalyptol.

Spray perfume, for spraying in the air of rooms, is prepared from the subjoined recipe :—

Five kilos of 92 per cent. alcohol, 4 kilos of warm water, 1 kilo of eau de Cologne, 1.3 kilos of jasmine extract, 700 grams of rose extract, 300 grams of acacia extract, 750 grams of violet root tincture, 1 gram of musk tincture, $\frac{1}{2}$ gram of crystallised nerolin.

SPANISH LEATHER, AND SPANISH PASTE.

The first of these articles is prepared by immersing a square strip of wash-leather for three to four days in a solution of—Benzoin, 250 grams; bergamot oil, 20 grams; lemon oil, 20 grams; grass oil, 20 grams; lavender oil, 20 grams; nutmeg oil, 10 grams; clove oil, 10 grams; neroli oil, 40 grams; rose oil, 40 grams; santal oil, 40 grams; essence of Tonka bean, 20 grams; cinnamon oil, 10 grams; alcohol, 1 litre. The leather is then taken out, drained, spread out on a glass plate to dry, and afterwards coated on the rough side, by means of a brush, with a mixture prepared in a mortar, and consisting of sublimed benzoic acid,

10 grams ; musk, 1 gram ; civet, 1 gram ; gum arabic, 30 grams ; glycerin, 20 grams ; and water, 50 grams. The piece is next folded across the middle, smoothed down with a paper knife, and left to dry under slight pressure. When dry, it will give off an exceedingly fine perfume for a period of years ; it is usually enclosed in a strong silk case. Instead of leather, four to six layers of blotting paper may be treated in the same manner and enclosed in a cover ; and thin layers of cotton wadding, laid between two sheets of paper, may also be scented in the same manner.

To prepare Spanish paste, the following ingredients may be intimately mixed in a mortar, sufficient water being then added, drop by drop, to form a plastic paste :—Ambergris, 20 grams ; benzoin, 40 grams ; musk, 20 grams ; vanilla, 20 grams ; violet root, 20 grams ; cinnamon, 20 grams ; bergamot oil, 40 grams ; rose oil, 30 grams ; gum arabic, 40 grams ; and glycerin, 40 grams. This paste is divided into lumps about the size of hazel nuts, and packed in scent boxes or cassolettes.

For removing the disagreeable stains produced by silver nitrate, the only effective reagent is potassium cyanide. This is poisonous, and therefore must be used with great care ; it is, however, possible to mix this cyanide with soap, the diluted preparation being then known as

PHOTOGRAPHERS' SOAP.

When applied to the stained surface, this preparation will quickly, though not instantaneously, remove the colour. On account of its unstable character, potassium cyanide cannot be added to soap in a hot liquid state, but must be incorporated with the same in the following manner :—Good curd soap is cut into fine shavings, allowed to dry slowly, and then reduced to powder, the latter being afterwards mixed with 10–20 per cent. of potassium cyanide in a finely divided condition. To ensure uniformity of mixing, the mixed powder should be run twice

through the soap mill; which done, the percentage of cyanide in the soap can be guaranteed. The mixing should be effected very quickly, and at the lowest possible temperature, since the cyanide is readily decomposed by warmth and moisture, and loses its efficacy.

For sale, this soap should be packed in tinfoil, either as it is or after conversion into milled soap.

Another way of preparing photographers' soap is by incorporating the above proportions of potassium cyanide with a carefully prepared soft soap, by stirring. This product should be packed in good tins for sale.

SCENT CAKES.

These preparations of scent in the solid form are made of the ingredients given below, mixed with melted paraffin wax at the lowest possible temperature. The melted mass is then cast into circular moulds,—such as the lids of small tin boxes,—from which it is removed when cold and wrapped in tinfoil.

White rose.—Geranium oil, 20; bergamot oil, 20; patchouli oil, 3.

Lavender.—Mitcham lavender oil, 1000; bergamot oil, 500; cinnamon oil, 6; geranium oil, 50; neroli oil, 6.

Ess Bouquet.—Bergamot oil, 600; lavender oil, 200; clove oil, 140; nutmeg oil, 70; coriander oil, 20; santal-wood oil, 70; rose oil, 40; geranium oil, 50; orange oil, 12.

SPECIALTIES FOR THE TEETH.

Odontine.—This preparation may be recommended as a specialty for cleaning the teeth, 10–20 drops being poured on to a tooth brush, previously moistened with water. It is an agreeable and elegant preparation, which is easily introduced, and forms a very saleable article. The method of preparing it is as follows:—

125 grams of Quillaya bark and $3\frac{1}{2}$ grams of persio are digested for several days in a close vessel with a mixture of equal volumes of water and alcohol. To this liquid, measuring about 1 litre, is added a mixture of—Heliotropin, 0.13 gram; peppermint oil, 20 drops; and oil of aniseed, 10 drops, dissolved in 30 c.c. of alcohol. After standing several days in a filled, stoppered bottle, set in a moderately warm place, the solution is filtered if necessary, mixed with 500 grams of glycerin, and filled into bottles holding 2–3 oz.

Another method: 90 grams of powdered myrrh are extracted with a mixture of 720 grams of alcohol and 240 grams of water. At the same time, 45 grams of Spanish soap, cut into shavings, are dissolved in a mixture of 720 grams of alcohol and 240 grams of water. Both are filtered, mixed together, coloured wine-red with alkanna root or alkannin, scented with a mixture of—Lemon oil, 30 drops; peppermint oil, 30; oil of wintergreen, 10; sternanis oil, 30 drops; and, finally, 120 grams of glycerin are added. After standing one or two weeks in a cool place, the liquid is filtered and filled into bottles, after having received an addition of 18 grams of acetic ether.

Still other methods are the following:—

1. Sixteen parts of talc, 8 parts each of medicinal soap and white sugar, 1 part each of gum arabic and Berlin red, 1 part of peppermint oil, 4 parts of glycerin, and 4 parts of distilled water, are mixed to a homogeneous paste.

2. Prepared oyster shells, 96 parts; powdered pumice, 4; violet root, 12; Venetian soap, 20; and Berlin red, 12 parts, are mixed, triturated with 8 parts each of white syrup and glycerin, 1 part of peppermint oil, and $\frac{1}{2}$ part of clove oil; then made up to a uniform paste with 10–12 parts each of alcohol and distilled water.

Owing to the presence of pumice, and consequent attritive action on the teeth, the last recipe cannot be specially recom-

mended. Probably, also, the oyster shells mentioned could be advantageously replaced by precipitated chalk.

Kalodont, a highly popular preparation for cleansing the teeth, is packed in tubes, and can be successfully imitated by means of the following recipe:—Equal parts of soap powder from good neutral soap, finest levigated chalk, and 28° Be. glycerin are carefully mixed, and incorporated with sufficient water to form a thin pulp. This is coloured with soluble soap, red or carmine solution, scented with peppermint oil, warmed and stirred in a porcelain dish over the water bath, re-cooled, and filled into tubes whilst still semi-fluid.

Eau de Botot, which is also popular, is prepared in different ways, the best being the English and French recipes. Ratanhia root, from *Krameria triandra*, a plant indigenous to South America, is frequently used, and furnishes a red-coloured alcoholic extract.

The English recipe runs as follows:—Cedar essence, 4 litres; myrrh essence, 1 litre; ratanhia essence, 1 litre; lavender oil, 20 grams; peppermint oil, 30 grams; rose oil, 10 grams.

French recipe:—Anise, 280 grams; cochineal, 20 grams; mace, 10 grams; clove, 10 grams; cinnamon, 80 grams; alcohol, 3 litres; peppermint oil, 20 grams.

The following recipe, though more complicated, yields a very fine product:—Lemon rind, 100 grams; cloves, 100 grams; horse radish, 100 grams; rose leaves, 100 grams; and soap powder, 100 grams, are finely ground or pounded, suffused with 8 litres of alcohol, and then distilled. In about three weeks the preparation is scented with 25 grams of cassia oil and 160 of peppermint oil, and filled into bottles.

CARBOLIC POWDER AND SALICYLIC POWDER

are in plentiful demand, especially in the summer months, and form very profitable articles.

Carbolic powder is prepared by melting 60 parts of resin

and 70 of stearin by gentle heat, and to the cooled but still fluid mixture are added 25 parts of carbolic acid, the whole being then converted into a uniform powder by careful trituration with 700-800 parts of levigated chalk. The powder is packed in a dredger (which can be kept closed when not in use), and is applied by dusting.

Salicylic powder.—Pure starch powder is added by degrees, with continued stirring, to a 2-3 per cent. alcoholic solution of salicylic acid, the volume of this solution being such that it extends a hand's-breadth above the starch when the latter has settled down. This being accomplished, the alcoholic liquid is poured off, the residue pressed in a close muslin cloth, dried, triturated, and dried again at a temperature of 80° C. The objection that the incorporation of the salicylic acid and starch could be as completely effected by dry mixing, is disproved by examining the mixture under the microscope, which will show that, in the case of a starch containing 2 per cent. of dry salicylic acid, as many as five, six, or even more starch granules can be counted in some parts before one fragment of the acid crystals is seen; whereas, with the method described above, each grain of starch will be found embedded in a rudimentary micro-crystal of the acid.

SALICYLATED TALLOW

is prepared by dissolving 2 parts of salicylic acid in 5 of benzoïn tincture, and adding the solution to a melted digest of 100 parts of mutton tallow with 5 of gum benzoïn, the whole being stirred a few times, and packed into suitable tins.

It is advisable to employ benzoated tallow, since this has a more agreeable odour, and keeps the preparation from becoming rancid, at least for a long time. To save the expense of the tins, the article may be made up in stick form. For this purpose, 100 parts of benzoated tallow are melted with 50 of olive oil and 50 of yellow wax, the melt then receiving an

addition of 4 parts of salicylic acid, previously dissolved in 10 parts of tincture of benzoïn, and the whole finally poured into the same moulds as are usually employed for stick pomades. The sticks are cut into proper lengths, wrapped in tinfoil, and labelled.

STICK POMADES.

Special preparations are frequently employed to keep the hair in place; and a suitable article for this purpose can be obtained from the following recipe:—

Olive oil	90 grams.
Yellow wax	70 „
Spermaceti	10 „
Bergamot oil	2 „
Melissa oil	0·3 „

The odour of this pomade recalls that of honey; but, of course, it can be scented in any desired manner, *e.g.* with geranium oil and a little cumarin.

In preparing stick pomade with vaseline, the basis consists of a mixture of—Ceresin, 100 parts; vaseline, 200 parts—white vaseline for white pomade, and yellow vaseline for yellow or coloured preparations;—tallow, 100 parts; and lard, 50 parts. These materials are melted in the order given, the mass being coloured as required—rose-red with alkannin, brown with umber, black with vine black. When cool enough to be poured into the moulds, it is scented with 6-7 parts of any desired perfume, *e.g.* a mixture of bergamot oil, $2\frac{1}{2}$ parts; petit grain oil, $2\frac{1}{2}$; lavender oil, 1; and clove oil, 1 part.

MENTHOL PREPARATIONS.

The menthol preparations are well adapted for the production of specialties. Menthol is the name applied to the stearoptene isolated from the ethereal oils obtained from various species of mint, of Japanese, Chinese, and North

American origin. This product forms white crystalline masses, with a peculiar smell and taste of peppermint, though that from Chinese oil has a bitter flavour, and smells more like spearmint. It is sparingly soluble in water, but more readily so in alcohol, ether, oil, and chloroform. Pure menthol melts at 43° C. and boils at 212° C.

Since menthol now enjoys extensive use, a few recipes of practical value are subjoined.

MENTHOL MIGRAINE OINTMENT.

Two parts of white vaseline and 85 of lanolin are united in a porcelain basin over the water bath, and to the mixture is slowly added, with stirring, 10 parts of menthol, the liquid being poured off as soon as the mass begins to thicken.

CHLORAL MENTHOL.

This is a mixture of chloral and menthol in equal parts, warmed together on the water bath to a temperature not exceeding 30° C. The resulting thick oily liquid has proved efficacious in facial neuralgia and toothache.

MENTHOLINE, INSTANTANEOUS.

Ten parts of menthol are dissolved in 78 parts of 95 per cent. alcohol, and mixed with 12 parts of strong ammonia. This liniment acts very rapidly in the case of sprains, local smarting, etc.

MENTHOL VINEGAR.

Three parts of menthol and 97 of white vinegar. When largely diluted with rose water, it may be used as a washing fluid.

MENTHOL TOOTHACHE DROPS.

Menthol, 8 parts; chloroform, 8; and 95 per cent. alcohol, 84 parts, mixed together as intimately as possible. In cases of

toothache resulting from caries, a little cotton-wool is steeped in the preparation and inserted in the hollow of the tooth.

MENTHOL ICE.

Ten parts of spermaceti are melted with an equal quantity of white vaseline in a porcelain dish on the water bath, and 10 parts of menthol are added to the mixture.

MENTHOL SNUFF FOR CATARRH.

Menthol, 15 parts; boric acid, 30 parts; and ammonium chloride, 55 parts, are triturated and intimately mixed together in a porcelain mortar.

MENTHOL CREAM.

Menthol cream, frequently used by barbers for cooling the face after shaving, is prepared from—Tragacanth, 1 part; glycerin, 3; alcohol, 4; menthol, 4; water, *quant. suf.* The tragacanth is first suffused with warm water, and left for two to three days with frequent shaking, the glycerin being then added, followed by the menthol dissolved in alcohol.

SOZODONT

is prepared from—White Castile soap, 4 grams; glycerin, 4; water, 1; alcohol, 2 grams; peppermint oil, 5 drops; wintergreen oil, 12 drops. Cochineal is used to impart a red colour.

SAND-ALMOND BRAN

is a very popular toilet preparation, compounded by dissolving 4 parts of borax in 8 of glycerin, the solution being used for impregnating 30 parts of extremely finely powdered sand. The mixture is gradually incorporated with 80 parts of almond bran, and scented with a little oil of bitter almonds.

ALMOND BRAN

is prepared, for cosmetic purposes, in the following manner:—
 Calcined soda, 50 grams ; powdered borax, 100 ; finest violet root
 powder, 250 ; and finest wheat flour or rice meal, 2000, are mixed
 together, 120 grams of fatty oil of almonds being then uniformly
 incorporated therewith. Finally, 100 grams of alcohol and
 about 50 drops of oil of bitter almonds are added, and the
 whole is rubbed through a medium hair sieve.

SCENTED PARAFFIN TABLETS

can also be recommended as cosmetic specialties in the wider
 sense of the term. Paraffin is melted on the water bath and
 mixed with any desired proportion of scent, the whole being
 then cast into moulds, and the tablets employed for rubbing
 the handkerchief or clothing.

Suitable recipes herefor are—To 125 grams of paraffin :
 essence of lavender, 7 grams ; essence of cloves, 3 ; essence of
 geranium, 1 ; essence of bergamot, 7 ; vanillin, $\frac{1}{2}$; glycerin, 3 grams.
 To 250 grams of paraffin : neroli essence, 3 grams ; essence of
 geranium, 3 ; essence of lavender, 3 ; essence of bergamot, 7 ;
 essence of cloves, $\frac{1}{4}$; heliotropin, 1 ; glycerin, 3 grams. To 125
 grams of paraffin : essence of ylang-ylang, 7 ; cumarin, 1 ; essence
 of musk, 1 ; essence of cloves, 3 ; santal-wood essence, 2 ; glycerin,
 3 grams.

MASSAGE COSMETIC.

This is a new specialty, which is largely used in America
 before, during, and after massage. The preparation known as
 "Madame Dornier's Cosmetic," which is claimed to be an
 excellent softener for the hands, is compounded as follows :—

Alum	.	.	.	.	.	.	.	4 parts.
Oak bark.	.	.	.	.	.	.	.	4 "
Aniseed	.	.	.	.	.	.	.	8 "
Thyme	.	.	.	.	.	.	.	8 "

Sage	8 parts.
Rosemary	8 „
Hyssop	8 „
Lavender flowers	8 „
Wormwood	8 „
Camphor	8 „
Peppermint	8 „
45 per cent. alcohol	1000 „

The ingredients are macerated for fifteen days and then filtered. It would be advisable, perhaps, to strain the mixture, then add the alum, and filter after settling.

CHAPTER IX

ANTISEPTIC WASHES AND SOAPS, TOILET AND HYGIENIC SOAPS.

THE cosmetic preparations employed for treating the hair, and which are not infrequently prepared by members of the laity, do not, for the most part, reach that state of perfection that might be expected from the present development of chemistry and physiology; and, in fact, we still find in this branch a number of useless, unsuitable, and even injurious preparations. Nevertheless, large sums are laid out annually on useless preparations claiming to promote the growth of the hair on hairless parts of the cuticle. A preparation of this kind (for forcing the beard) was investigated by Simon, and was found to possess the interesting property of causing intense itching and smarting on the first application, and in one case produced great swelling of the face and frightful smarting, these effects being readily explained by the fact that the preparation proved to be a 5 per cent. solution of cantharides.

Again, washes for the hair and head only too frequently contain ingredients that redden the skin, such as cantharides tincture, mustard tincture, tincture of Spanish pepper or onions, oil of mustard, formic acid, and the like, the use of which cannot be recommended, since none of them is capable of stimulating the growth of the hair, even when constantly applied. In fact, any good results following their use could be equally attained by means of any other hair wash, especially those of an antiseptic character, and are in reality merely the

consequence of the increased attention bestowed on the general care of the hair, more particularly as regards washing the head. On the other hand, the continued use of such concentrated preparations of the ingredients referred to, is by no means free from danger. Simon found, in several instances, in the case of an eau de Quinquine prepared with tincture of cantharides, a short continued use produced itching and smarting of the scalp, and even a rush of blood to the head, with the usual consequences. Hence such materials should only be used under the directions of an experienced medical man, and do not belong to the domain of cosmetics at all.

It is well known that the falling out of the hair is due for the most part to the action of small vegetable parasites, which find their way into and destroy the hair, mostly as the result of continued uncleanness,—especially defective attention,—though in some cases by direct contagion. The hair may also fall out in consequences of general disturbances of nutrition, consumption, loss of power, and illnesses of all kinds, especially nervous complaints and skin diseases; as also by defective ventilation of the scalp, or insufficient nutrition of the same, by constantly wearing head coverings that are either impermeable, too heavy, or too tight, such as waterproof hats and caps, helmets, etc.

Now, whilst in the case of hair falling out as the result of illness, the only way to ensure the success of remedies is to simultaneously improve the general strength of the entire organism, a similar occurrence caused by parasites or defective care of the hair can be arrested by the aid of suitable cosmetic preparations; this also applies to cases resulting from the use of unsuitable head coverings, provided the cause be removed at the same time.

In the first place, it will be necessary to briefly consider the proper care of the hair in health, since on this point there are current a variety of opinions, chiefly based on habit and train-

ing; since, whilst an uneducated individual cares little for cleansing the hair, and looks mainly to pomade, those who are better informed attach the greatest importance to cleanliness, attained by frequent washing. Certainly, the latter is the more correct view, though the moderate use of suitable pomades has its justification. Fat imparts gloss, smoothness, and softness to the hair, and diminishes the evaporation of moisture from the hair and scalp. Hair that is insufficiently greased, suffers considerably from excessive evaporation, which is intensified by perspiration and heat, and thereby loses its gloss, smoothness, and softness, becoming hard and even brittle. If, therefore, the natural fat of the hair is insufficient to keep it glossy and smooth, it must be supplemented by frequent small additions of fat, and this should always be done whenever the hair is washed.

The extremely large surface presented to the atmosphere by the hair renders the latter highly susceptible to the influence of the former. The fat suffers oxidation, and, unless prevented by careful attention, quickly unites with the ever-present dust to form an impermeable incrustation. When insufficient attention is bestowed on the hair, an increased secretion of the natural fat occurs, and this is just as frequent a source of scurf as excessive pomading. On the other hand, if the hair receive proper attention, especially as regards frequent washing, this scurf will not appear; and it is the recognition of this fact that has led to the introduction of the large number of hair washes and the like now in use. As a matter of fact, also, a good hair wash is really useful for the purpose in question.

A good hair wash should free the hair from grease, help to remove dirt and dust, be convenient to use, and exhibit a pleasant odour. It therefore follows that such a hair wash should be slightly alkaline, in order to be able to emulsify the fat. For private use, an aqueous solution may be

employed, but preparations intended for sale should contain an addition of alcohol to ensure rapid drying. An addition of glycerin is also advisable, since this substance makes the hair soft and supple, besides most effectually contributing to the removal of scurf.

Hair washes may be divided into three classes: such as are chiefly mechanical in their action; medicinal washes, for removing scurf, destroying parasites, etc.; and antiseptic washes, possessing the properties of the two other kinds in common.

The following recipes are recommended for ordinary hair washes:—

Rosemary hair wash.—12 grams of potassium carbonate are dissolved in 700 of rose water and shaken up with the following mixture: 90 per cent. alcohol, 300 grams; lavender oil, 1; lemon oil, 1; neroli oil, $\frac{1}{2}$; bergamot oil, 2; rosemary oil, 5 grams. After thorough agitation, the wash is stored for three days in a very cool place, filtered, and mixed with 100 grams of glycerin.

Laurel hair wash.—Ammonium carbonate, 20 grams; borax, 35; rose water, 700; 90 per cent. alcohol, 300; ethereal oil of laurel, $\frac{1}{2}$; pimento oil, $\frac{1}{2}$; bergamot oil, 2 grams. The method of preparation is the same as for the Rosemary wash.

Bay rum water.—Ammonium carbonate, 25 grams; borax, 25; honey, 50; rose water, 700; rum, 300 grams; geranium oil, 10 drops; bergamot oil, 15; rosemary oil, 8; ethereal oil of laurel, 2; pimento oil, 1; oil of bitter oranges, 1 drop. The honey, borax, and ammonium carbonate are dissolved in the rose water, the oils in the rum, and the two solutions are mixed. An addition of glycerin is unsuitable in this case.

Quillaya hair wash.—Borax, 25; 90 per cent. spirit, 450; Quillaya tincture, 1:10 spirit, 100; eau de Cologne or similar scent, 75. The method of preparation is the same as for the Rosemary wash.

Soap hair washes.—(1) Soap spirit, 200; glycerin, 100; scent,

75 ; eau de Cologne, Bouquet d'Amour, eau de Millefleurs, 90 per cent. spirit, 325 ; water, 300 parts.

(2) Medicinal soap or white cocoanut-oil soap, 25 grams ; ammonia, 10 ; rose water, 800 ; glycerin, 100 ; 90 per cent. spirit, 100 ; geranium oil, 2 ; bergamot oil, 1 gram ; oil of bitter almonds, 2 drops.

The best and most agreeable of the above washes are the Quillaya wash and the soap washes. American shampoo fluid, which is prepared with eggs, will only keep a very short time. The last of the above soap washes is exceedingly cheap, highly agreeable, and fully equal to the American preparation.

Athenian hair wash.—Potassium carbonate, 70 grams ; sassafras wood, 250 grams ; rose water, 4 litres ; orange-flower water, 4 litres ; alcohol, 1 litre.

The ingredients are digested together for a month. The potassium carbonate and the alcohol have a cleansing effect on the hair. After using this wash, the hair, when dry, should be treated with a fine pomade or hair oil, in order to restore fat and gloss.

Glycerin and Cantharides wash.—Ammonia, 100 grams ; essence of cantharides, 100 grams ; rosemary water, 8 litres ; glycerin, 300 grams ; rose oil, 20 grams. The cantharides essence is prepared by digesting 50 grams of powdered Spanish fly with 1 litre of strong alcohol. The ammonia acts like potassium carbonate, as a detergent ; the glycerin imparts softness and suppleness to the hair.

Blossom hair wash.—Acacia essence, 200 grams ; jasmine essence, 200 ; orange-flower essence, 290 ; Tonka-bean essence, 100 ; tuberose essence, 200 ; vanilla essence, 100 grams ; rose water, 2 litres ; alcohol, 2 litres.

Soap hair wash.—Rosemary water, 5 litres ; rondeletia perfume, 300 grams ; saffron, 5 ; soap, 30 ; alcohol, 300 grams.

The finely divided soap and the saffron are boiled in a little

distilled water until the first named is completely dissolved, whereupon the other ingredients are added, the whole well mixed, and then set aside for a few days, for the coarser parts of the saffron to settle down. This preparation has a very handsome appearance, and, when filled into cut-glass bottles, has an opalescent sheen resembling mother-of-pearl, though, viewed by transmitted light, it is an almost perfectly transparent saffron-yellow liquid.

Among the medicinal hair waters may be mentioned—

Unna's Spiritus Capillorum Resorcin, 25 grams; castor-oil, 25; 95 per cent. spirit, 750 grams; eau de Cologne, 200 grams. This is a specific against parasitic skin diseases, eczema of the scalp, and scurf. The scalp is first cleansed with a lukewarm soap bath, the hair dried, and the wash applied by rubbing in with a soft cloth, the operation being repeated daily until the desired effect is attained. All the other medicinal hair washes are applied in the same way.

Naphthol hair water.—B-naphthol, 25 grams; Quillaya tincture (1 : 10), 100 grams. This preparation, which is scented with ylang-ylang oil or a little heliotropin and 1–20 of iris oil, is successfully used for combating scurf.

The antiseptic hair waters are also used for removing scurf, curing parasitic diseases of the hair, and for disinfecting the hair and scalp generally. They are applied by pouring a tablespoonful into about $\frac{1}{3}$ th of a litre (120 c.c.) of lukewarm water, and washing the head with a sponge dipped into the liquid. The hair is next dried, combed, and wrapped up with a cloth or hood for three-quarters of an hour, at the end of which time it is again combed, lightly oiled, and arranged in the usual way.

The following recipe is typical of the group :—Antiseptic, 25; glycerin, 100; spirit, 900, with a little scent.

The antiseptic may consist of—Salicylic acid, 25 grams; or

boric acid, 100 grams ; or salol, 20 grams ; or thymol, 30 grams ; or eucalyptol, 40 grams ; or 30 grams of benzoic acid from toluol, and 15 grams of thymol.

The perfume may be either— $\frac{1}{2}$ gram wintergreen oil, 0.1 gram geranium oil, and 0.1 gram orange-flower oil ; or 1 gram cumarin sugar (1:1000), 4 drops rose oil, 4 drops bergamot oil, 2 drops French geranium oil, 2 drops ethereal oil of bitter almonds ; or 1.0 gram cumarin sugar, 2.5 grams peppermint oil, 1.09 gram clove oil, 0.5 gram wintergreen oil ; or, especially for thymol washes, 8 grams peppermint oil, $2\frac{1}{2}$ grams clove oil, 1 gram sassafras oil, and 10 grams cumarin sugar.

The antiseptic hair washes combine the properties of the previous hair waters with those of the true hair washes. They are applied in an undiluted condition, and, in addition to cleansing, disinfect the hair and scalp. They may be recommended not merely as washes, but also as medicaments against scurf, parasitical diseases of the hair, etc., and form the most suitable preparations for use by the laity.

A trustworthy recipe for a wash of this kind is the following:—

Borax, 25 grams ; pure potash soap from olein or olive oil, 40 grams ; glycerin, 100 grams ; rose water, 650 grams ; spirit, 250 grams ; crystallised phenol, creolin, solveol, or similar cresols, 15 grams. The soap is dissolved in water, the glycerin and borax being then added, followed by the antiseptic and scent dissolved in the spirit, the whole being then set in the cool for a day, and finally decanted.

Mention might be made of a number of hair waters, for which stimulative action on the growth of the hair is claimed on account of the presence of quinine or similar alkaloids, Peruvian balsam, or tannin. Peruvian balsam is the most promising, but exerts its favourable effects only when added to pomades, and cannot be recommended for hair waters.

Tannin substances in general retard the growth of the hair, and their sole recommendation is that they counteract excessive perspiration of the head. Quinine and chinchona bark have no stimulative action on the growth of the hair, and merely serve as antiseptics, for which reason quinine should be entirely omitted, its place being taken by salicylic acid. The tonic properties of quinine are only manifested when applied internally; the sole value of this alkaloid as a constituent of hair washes being its antiparasitic, *i.e.* antiseptic, properties.

Nevertheless, since, in despite of all teaching, such preparations are *continually asked for*, the already mentioned excellent preparation of Dieterich for preventing the hair from falling out will now be supplemented by a few recipes.

HAIR RESTORER.

Cantharides tincture	50 grams.
Extract of gall nuts	50 "
Essence of musk	10 "
Carminc	5 "
Alcohol	100 "
Rose water	1 litre.

The gall-nut extract is prepared by digesting 100 grams of powdered gall nuts with 1 litre of alcohol. The extract of chinchona bark, mentioned below, is prepared in a similar way.

TANNO-QUININE HAIR RESTORER.

Extract of chinchona bark	50 grams.
Extract of gall nuts	50 "
Carminc	10 "
Neroli oil	5 "
Nutmeg oil	5 "
Alcohol	100 "
Rose water	1 litre.
Orange-flower water	1 "

BAUME DE MILAN POUR LES CHEVEUX.

Lard	500	grams.
Almond oil	500	„
Spermaceti	50	„
Carmine	10	„
Styrax essence	30	„
Cantharides tincture	20	„
Tolu essence	30	„

BEARD FORCING LOTION.

Lard	500	grams.
Almond oil	500	„
Spermaceti	20	„
Cantharides	20	„
Carmine	10	„
Bergamot oil	5	„
Lavender oil	5	„
Santal oil	5	„

The cantharides are first triturated to a fine powder along with the carmine, the product being incorporated with the ethereal oils, and finally with the other ingredients.

Antiseptic scurf water.—Five parts of tannic acid are stirred up with 20 parts of formaldehyde, and treated with 5 parts of sulphuric acid. The reddish mass is placed in water, filtered, and washed until the washings cease to show any trace of acid, whereupon the residue is dried by gentle heat. Five parts of this product are dissolved in 100 parts of spirit, and furnish a liquid equal to eau de Quinine de Pinaud, and only requiring to be scented. The red colour is fast to light, but does not dye the scalp, nor does the colour come off.

ANTISEPTIC SOAPS: THEIR PREPARATION AND APPLICATION
IN COSMETICS.

Generally speaking, the action of soaps is of a cleansing character; but by suitable adjuncts they may acquire new and valuable disinfectant properties.

In order to ascertain on what the cleansing power of soap is based, Rotondi precipitated a solution of Marseilles (olive oil) soap with salt, dissolved the precipitate in 95 per cent. alcohol, filtered the solution, evaporated the solvent from the filtrate, and repeated the operation, thus producing a pure soap which was completely soluble in boiling water, and had an alkaline reaction towards litmus, without, however, containing any alkali hydroxide or carbonate. This soap gave, on analysis, the following composition:—

	Air dry.	Dried over Sulphuric Acid.
Water	29.41	
Fatty acids (m.p. $41^{\circ}5$ C.) .	64.90	91.43
Sodium oxide	7.82	11.32
Inorganic salts	0.16	0.31

From this soap Rotondi prepared solutions of different strength, and subjected the same to dialysis in the following manner:—(1) 100 grams of the soap were reduced to the smallest possible fragments, and then dialysed for thirty days at 15° C. (2) 100 grams of soap were dissolved in 2 litres of water, and treated as before. (3) 100 grams of the soap were dissolved in 2 litres of boiling water and dialysed for thirty days at 15° C. (4) 100 grams of the soap were dissolved in $\frac{1}{2}$ litre of distilled water and dialysed at a temperature of 80° C. At the end of the experiment the various dialysed and undialysed liquids were evaporated on the water bath, the residues, after drying over sulphuric acid, giving the following results:—

	Percentage of Sodium Oxide in the Residue from the	
	Dialysed Liquids.	Undialysed.
Experiment 1, ...	15.06	10.81
„ 2, ...	15.34	10.74
„ 3, ...	16.12	9.56
„ 4, ...	15.58	9.33

In the first and second experiments particularly, the propor-

tion of dialysed to undialysed soap was very small. The evaporation residues from the dialysed liquids are soluble in water, and the solutions do not become turbid unless exposed to the air for several days. On the other hand, those from the undialysed liquids are soluble in hot water alone, and immediately become turbid on cooling. If the dialysis be sufficiently prolonged, the residue finally obtained is undialysable, and is also insoluble in hot water. Since the dialysed liquids contain no free alkali, Rotondi draws the following conclusions from his experiments:—1. The neutral soaps on a basis of alkali are decomposed by water into basic soaps, soluble in hot and cold water, and into acid soaps insoluble in cold water and only sparingly soluble in hot water. 2. The decomposition of the neutral soap is effected more easily in hot water than in cold, and the rapidity varies according to the concentration and temperature of the solution. 3. The basic soaps dialyse easily; not so the acid soaps. 4. The solution of basic soap obtained by dialysis may still contain neutral soaps, which can again be decomposed into acid and basic soaps, until, finally, the solution contains basic soap only. 5. Neither alkali hydrate nor alkali carbonate is liberated during the decomposition of the neutral soaps by water, as may be easily ascertained by precipitating the basic soap by common salt and dialysing the liquid after filtration.

The action of water on soap can also be demonstrated without dialysis. With this object Rotondi prepared a solution of 100 grams of soap in 1 litre of water by warming, and left the solution to stand twenty-four hours in an open vessel.

He then succeeded, by filtration, in separating an insoluble soap A, and a liquid soap B. The soap A, dried over sulphuric acid, gave a residue containing 10·17 per cent. of sodium oxide. This residue was completely soluble in boiling water; but, on cooling and filtering the solution, an insoluble soap, containing 6·11 per cent. of sodium oxide, was obtained. On evaporation,

the solution B left a residue containing 14.12 per cent. of sodium oxide. On being left to stand in an open vessel for fourteen days, it deposited a fibrous substance, which, when separated by filtration and dried over sulphuric acid, contained 7.84 per cent. of sodium oxide. The filtrate gave, on evaporation, a residue containing 15.97 per cent. On being left to stand fourteen days longer in an open vessel, it gave a further insoluble deposit; but nothing further could be brought to deposit from the filtered residual liquid. The residue on evaporation, however, contained 16.21 per cent. of sodium oxide. This residue proved completely insoluble in cold water, gave a strongly alkaline reaction, was completely precipitable by common salt, and the liquid filtered from the precipitate contained no alkali hydrate or alkali carbonate.

It follows from these experiments that the neutral soaps are decomposed by water into acid and basic soaps; that the basic soaps are completely soluble in cold water; whereas the acid soaps are only very sparingly soluble, even in hot water; and that the basic soaps are completely precipitated by common salt, without any liberation of free alkali. The aqueous solution of the basic soaps gives—especially when hot—a clear liquid with oleic acid and other fatty acids, this liquid also remaining clear on cooling, provided a sufficient quantity of basic soap be present. If, however, the solution be exposed to the air for several days, it becomes turbid, owing to the separation of the mixture of basic and neutral soaps into acid and basic soaps. The basic soaps dissolve the acid soaps in the warm; but the solution becomes turbid on cooling, after a longer or shorter time, according to the proportions of basic and acid soap present. This explains why ordinary soaps are soluble in hot water, although the decomposition is more complete therein than in cold water. The cold aqueous solutions of the basic soaps do not combine with fats, but merely form emulsions. In this manner one can obtain with triolein a mass resembling clotted

cream, and keeping for several months without any separation of oil globules. The mass is also unaltered by warming; but when treated with an addition of 80 per cent. alcohol, the fat is separated and floats above a clear liquid containing no glycerin, a proof that the fat has not been saponified, but only emulsified.

This property is also possessed, though to a smaller extent, by the ordinary neutral soaps, especially in freshly prepared solutions, possibly because they are then more easily decomposed into basic and acid soaps than when older. That no saponification occurs, is evidenced by the fact that the fat can readily be separated by alcohol, and that no glycerin is to be found in the alcoholic solution. The acid soaps do not possess the property of emulsification, even in the warm, or if so, only to a very limited extent. If carbon dioxide be passed through a cold aqueous solution of basic soap, a milky liquid results, which, however, can be reclarified by exposure to the air, by warming, or by the addition of alcohol, though it is difficult to obtain the solution clear by filtering. Common salt will precipitate all the soap dissolved or suspended therein. The residual liquid is free from alkali hydrate or alkali carbonate. The insolubility of basic soaps in water containing carbon dioxide explains the low cleansing power of basic and neutral soaps when used in highly carbonated waters—quite independent of the presence of inorganic salts in the waters.

Unna prepares his medicinal soaps in accordance with the following principles:—He employs only one fat, namely, best beef tallow, and rejects cocoanut oil entirely, on account of its roughening effect on the skin when continuously used. The alkali employed consists of a mixture of 2—in the middle of summer 3—parts of freshly prepared soda lye and 1 part of potash lye, the total quantity being just sufficient to leave the saponified mass perfectly neutral in reaction. These soda-potash soaps are solid, more efficacious than soaps from soda alone, and do not peel like the latter sometimes do. Since,

when permanently incorporated with the skin, neutral soaps abstract fat therefrom, and produce a disagreeable brittleness, attended by congestion and roughness. Unna superfats his soaps, by taking, beyond the amount of fat necessary to complete saponification, an excess—averaging 3–4 per cent. of fat, consisting of 8 parts tallow and 1 part olive oil. These superfatted soaps have the additional advantage of better preserving sundry medicaments, such as acids—salicyelic acid—and unstable salts—sublimate—better than ordinary neutral soaps do. Of course, all the other conditions applicable to the best toilet soaps must also be fulfilled: in particular, the under-lye precipitated by a minimum quantity of salt must be completely removed, and the soap must be dried as perfectly as possible. Unna rejects as useless any secondary addition of glycerin, vasoline, or scent. The superfatted soda-potash soap he terms superfatted stock soap.

From this stock soap the medicinal soaps are prepared by grinding with weighed quantities of the medicaments employed, the mixture being first made with a small quantity of the stock soap, and the bulk of the latter is incorporated afterwards.

The application of these soaps may be of a threefold character. The least efficacious method is by washing in the ordinary manner, the soap being brought to a lather with warm water, rubbed into the skin, and then rinsed off again. The effect is far superior when the affected portions of the skin are lathered, and the soap rubbed off, at once or after a few minutes, with a dry cloth: but the best effect of all is produced by laying the soap lather on thick and allowing it to dry on. Hot or warm water is preferable to cold in using these soaps.

In order to prepare good lathering soaps by the warm process, Schneider recommends saponifying with a quantity of lye not quite sufficient to ensure complete saponification, the deficit

being made good by an equivalent amount of soda solution afterwards.

In such cases where a decomposition of the soap is feared in this wet process, as may happen with salicylic acid,—which may give rise to the formation of an alkali silicate, fatty acid being then liberated,—Schneider recommends the following imitation of the milled-soap process:—

The hardened outer portions of Venetian soap are pared off and the softer interior is rasped. A similar quantity of powdered medicinal soap—the exact amount depending on the percentage of moisture in the Venetian soap—is taken, and the medicaments to be used are mixed with this soap powder if they are in the solid form, but with the rasped Venetian soap when they are partly or entirely liquid. The two soaps are next mixed in a porcelain mortar, and put into tin, wooden, or cardboard moulds lined with tinfoil, the soap being pressed down firmly with the pestle. The moulded blocks are then wrapped in thin parchment paper, paraffin paper, or tinfoil.

A little stearite powder is generally added to medicinal soap by the makers, and this leaves the skin agreeably smooth after use.

Schneider's opinion, that the lathering qualities of boiled soap are improved by using an insufficient quantity of lye at the outset, is controverted by the *Seifensieder Zeitung*, which states that an easier and better way, having the advantage of furnishing a properly saponified soap, is to employ an addition of cocoanut oil.

According to Boiry, cocoanut oil is peculiarly adapted for the preparation of medicinal soaps, and he recommends the following formula:—900 grams of cocoanut oil are boiled in a porcelain pan along with 600 grams of 10° Be. soda lye until the whole has assumed a creamy appearance, whereupon 375 grams of 20° Be. soda lye are added, and boiling is discontinued when a sample of the mass solidifies

on being dropped on a cold plate. He then adds about 500 c.c. of distilled water, heats to boiling, and adds 375 grams of common salt; the soap rises to the surface of the liquid. After cooling, the soap is separated, by decantation, from the layer of glycerin and brine, and is purified by a couple of washings with 20 per cent. brine, followed by a quick washing with cold distilled water, after which it is drained, and pressed to remove the excess of water. This process furnishes a pasty soap, which Boiry employs in the preparation of the following medicinal soaps:—

Borax soap.—900 grams of stock soap and 900 grams of sodium borate are triturated to a uniform mass, which is divided into portions of 100 grams each (or other suitable weight), and pressed into moulds.

Coal-tar soap.—100 grams of tar are gradually triturated with 900 grams of stock soap in a mortar. The same procedure is also recommended for naphthol soap.

Carbolic soap.—50 grams of white carbolic acid are dissolved in about 25 grams of 90 per cent. alcohol, and the solution gradually incorporated with 950 grams of stock soap by trituration.

Sublimate soap.—Five grams of corrosive sublimate are dissolved in 80 grams of alcohol, and the filtered solution incorporated with the soap as above.

Sulphur soap.—This is prepared from 100 grams of washed flowers of sulphur and 900 grams of soap, in the same manner as borax soap.

Antiseptic soaps frequently contain free alkali, which, in many cases, produces troublesome irritation of the skin. Attempts have therefore been made to obviate this defect by the addition of boric acid, basing the practice on the idea that an excess of boric acid will increase the antiseptic properties of the soap. A soap of this kind, containing 5 per cent. of boric acid, is now largely sold. It may, however,

easily be shown that this soap by no means fulfils its object, the boric acid being combined with soda and not in the free state, the result being that a corresponding amount of fatty acid has been set at liberty. Such superfatted soaps form inferior antiseptics; and far greater efficiency is secured by using a soap containing only a very small amount (about 1 per cent.) of boric acid in addition to the antiseptic ingredients. Such a soap is found, by bacteriological examination, to be highly efficacious.

Powdered medicinal soaps have been recommended by Eichhoff as useful substitutes for the fatty medicinal soaps, the method of application being either by making them up into a paste with water, and using like ordinary soaps, or else to rub them into the skin with a woollen cloth. They are then either allowed to dry on, or are covered over with some waterproof material. The stock soap consists of a neutral soap, dried and ground to powder. This is converted into a superfatted soap powder by an addition of 2 per cent. of oleic acid and 3 per cent. of lanolin; or into an alkaline powder by adding $2\frac{1}{2}$ per cent. each of potassium and sodium carbonate. Eichhoff recommends the following preparations:—

Salicylic acid soap powder, with 5 per cent. of salicylic acid; salicylic-resorcin soap powder, with 5 per cent. each of salicylic acid and resorcin; salicylic-sulphur soap powder, with 5 per cent. each of salicylic acid and sulphur; salicylic-resorcin-sulphur soap powder, with 5 per cent. each of all three adjuncts; sulphur soap powder, with 10 per cent. of sulphur; camphor-sulphur soap powder, with 2 per cent. of camphor and 5 per cent. of sulphur; Peruvian balsam soap powder, with 10 per cent. of Peruvian balsam; camphor-sulphur-Peruvian-balsam soap powder, with 2 per cent. of camphor, 5 per cent. of sulphur, and 10 per cent. of Peruvian balsam; naphthol-sulphur soap powder, with 5 per cent. each of B-naphthol and sulphur; naphthol soap powder, with 5 per cent. of B-naphthol; camphor soap powder, with 5 per cent. of camphor; borax soap

powder, with 5 per cent. of borax ; thymol soap powder, with 2 per cent. of thymol ; benzoin soap powder, with 3 per cent. of benzoin ; pumice soap powder, with 20 per cent. of pumice ; chloride of lime soap powder, with 10 per cent. of bleaching powder ; quinine soap powder, with 2 per cent. of sulphate of quinine ; menthol soap powder, with 5 per cent. of menthol ; salol soap powder, with 5 per cent. of salol ; naphthalene soap powder, with 5 per cent. of naphthalene.

Liquid soaps, or soap essences, are coloured or uncoloured scented solutions of soap in alcohol.

The best stock for liquid soaps is Marseilles olive oil soap, which is completely soluble in hot 80 per cent. alcohol, though less perfectly so in cold alcohol. Soaps from tallow and other solid fats, though also completely soluble in hot alcohol, are unsuitable for the preparation of liquid soaps, inasmuch as they become merely translucent on cooling. On the other hand, soaps from vegetable oils are very soluble in alcohol, and offer the advantage of furnishing solutions which remain liquid and clear for a very long time. The presence of potash plays a great part in the preparation of liquid soaps ; nevertheless, an excess must be avoided, and 10–12 parts of pure potassium carbonate are sufficient to impart a slight alkaline reaction to 1000 parts of soap essence and increase its emulsive capacity.

The proportions of ingredients are as follows :—

White olive oil soap, 20 parts ; 80 per cent. alcohol, 100 parts ; pure potassium carbonate, 1–2 parts.

The soap is cut into small strips and placed in a roomy water bath, a closed vessel being necessary to prevent loss of alcohol. The potash and alcohol are then added, and the water bath is slowly heated, the soap being stirred to facilitate solution. When this is accomplished—forty to sixty minutes being required—the vessel is removed from the water bath, the scent and colouring matter—saffron or rosaniline—are added, and the soap is left for some hours to clarify, after which it is poured

off, or, better, filtered. In consequence of its large alcohol content, the product remains liquid at the ordinary temperature, and suffers little alteration even by cold. The scent may be of different kinds: orange oil, lemon oil, or compound scents, like—oil of bitter almonds, 8 parts, and bergamot oil, 2 parts; or oil of bitter almonds, 5 parts; oil of mirbane, 2 parts; and oil of cinnamon, 1 part.

A very fine perfume is furnished by a mixture of—Vanilla tincture, 10 parts; violet root extract, 20 parts; rose extract, 20 parts; orange-flower extract, 50 parts; white potash soap, 10 parts; potash, 1 part.

Robiquet recommends the following proportions in the preparation of liquid soap:—White soap, 1 part; 85 per cent. alcohol, 3 parts; distilled water, 1 part.

A composition of white soap, 2 parts; 80 per cent. alcohol, 4 parts; and soda crystals, 1 part, forms a good detergent; and Regnault gives the following recipe for a medicated soap that will lather freely:—White soap, 20 parts; distilled water, 30; 60 per cent. alcohol, 60; potash, 1 part; and scent, 8–10 per mil.

Saidemann gives the following recipe for his liquid soaps:—

One part of caustic potash is dissolved in an equal quantity of water and well shaken up for ten minutes with 4 parts of oil—olive oil or sesame oil—and $\frac{1}{4}$ part of alcohol. After standing for an hour, with frequent shaking, the mixture is diluted with an equal volume of water, and left for several days before filtering. The soaps have the consistence of thick glycerin, contain an excess of fat, and show a content of up to 70 per cent. of water.

Liquid soaps appear to be specially commendable, since it is only when the soap is liquid that it can be well rubbed into the skin, and removed or left on according to convenience; moreover, because antiseptics are more easily incorporated with liquid soaps than with any others; and, finally, because these soaps must necessarily be prepared from vegetable oils, which are

preferable to animal fats from a sanitary point of view, since these latter may have been derived from animals that have died.

Medicinal sulphur and tar-and-sulphur soaps are prepared by melting a suitable fat and saponifying it in a pan placed over an open fire and fitted with stirring appliances. Meanwhile, 1 per cent. of sulphur is mixed to a soft paste with hot 25° Be. potash lye. The necessary amount of potash lye is run into the fat in a thin stream, followed by the sulphur paste and the scent, the whole being then stirred until the mass has become thick and viscous, whereupon it is transferred to the frames, which are covered up whilst still very warm, and left to stand. On the following day the frames are opened and the soap is cut. The addition of a little glycerin to the soap is advisable, as it increases the delicacy and suppleness of the product.

The same procedure is followed in the preparation of tar-and-sulphur soaps, except that the proportion of sulphur is somewhat diminished and a corresponding percentage of birch-tar oil is added to the soap when the mass commences to thicken, whereas the sulphur and the scent are added to the fat before the lye is run in. If the soap gives off much heat when framed, the frames should be left for about an hour before being covered up.

Medicinal sulphur-sand soap is prepared in the same manner as sulphur soap, except that an addition of 30 per cent. of sand or pumice is stirred into the soap in the pan. Cheap scents alone, *e.g.* oil of lavender, cinnamon, or clove, are used for this class of soap.

Thiosavonal is the name applied to sulphur soaps soluble in water. These are made from sulphonated oils, the following recipe being given by Grube:—

Sulphur soft soap.—The thick sulphonated oil is rendered more fluid by the aid of alcohol, and is gradually mixed, by continued stirring, with an equivalent quantity of potash lye, also diluted with alcohol; the addition of large quantities of

lye at a time will cause a sudden deposition of sulphur, the risk of which, however, diminishes towards the end of saponification. Finally, a slight excess of potash lye is used; the complete saponification of the thio-fatty acids is indicated by the clearing of the liquid, and the fact that a sample will dissolve in water or alcohol to a clear solution. The excess of alkali is neutralised with an ethereal solution of fatty acid. The resulting soap solution is freed from alcohol in a steam bath, and, having been tested for neutrality, is evaporated to the consistence of ointment. 85 parts of this soap are then mixed with 15 of glycerin; the water content of this mixture is about 12 per cent., and the percentage of the potassium compound of the thio-fatty acids is about 5 per cent.

Liquid sulphur soap.—The method of working is the same as above, except that the soap is only evaporated to the consistence of syrup. 88 parts of this liquid soap are mixed with 12 of glycerin, the mixture containing 29.6 per cent. of water and 4 per cent. of the potassium compounds of thio-fatty acids. Both varieties of thiosavonal are readily miscible with large amounts of tar, the more solid one requiring to be warmed first.

Schleich's marble soap is particularly suitable for disinfecting the hands. It is prepared according to the following recipes:—

Clear, fresh, unfilled amber-yellow resin soap, cut	
into fine shavings	750 grams.
Warm water	1500 „

When this mixture has been completely dissolved by the aid of fire heat, there is added—

Wax paste	150 grams.
Stearin paste	150 „

and when these are dissolved, 7000 grams of coarse purified marble dust (No. 4 sieve) are stirred in by degrees.

The product must be sterilised and concentrated to the

consistence of syrup by heating for about an hour and a half, replacing the water lost by evaporation,—about 200 grams,—and stirring.

The wax paste is prepared by melting 100 grams of pure beeswax on the water bath, and adding about 10 grams of ammonia, with stirring. About 150 grams of sterilised distilled water are then stirred in, until a homogeneous emulsion is produced.

The stearin paste is made by melting 100 grams of pure stearin in a basin, stirring in 8–10 grams of ammonia, diluting to a pulp with water, and a few more drops of water and ammonia are added at a time, until a snow-white emulsion, which must be alkaline, is formed.

Marble soap is a light yellow mass, with an agreeable odour recalling beeswax. The consistence is that of paste, and, when rubbed with water, it lathers freely and falls apart without assuming the sticky, stringy appearance of the potash (soft) soaps. Its use makes the hands very soft and delicate, and leaves an exceedingly agreeable feeling on the skin for some time.

COSMETIC TOILET SOAPS AND HYGIENIC SOAPS.

According to G. Hurst, the free alkali in soaps can be determined by dissolving 10 grams of soap in boiling water, and titrating the solution with normal sulphuric acid, in presence of phenolphthalein as indicator: a good hard soap should not consume more than 0.75–1.0 c.c. of the acid. The author, however, demonstrated the unsuitability of this method by showing that, even when 100 c.c. of each of the three solutions—1–10—of superfatted cocoanut-oil soap is treated with 2 c.c. of a 1 per cent. solution of phenolphthalein, a faint rose-red coloration is given; whilst in the case of neutral cocoanut soap and stock soap a darker red is produced,—still more intense with alkaline soap,—a really decided decoloration only occurring when the greater portion of the soda present has been neutralised. The author has also shown that concentrated

solutions of soap are incapable of dissolving any large quantity of unsaponified fat, but merely emulsify same; and he therefore assumes that the fat in superfatted soaps is not present in the form of a triglyceride. As a matter of fact, he has succeeded, by shaking up solutions of superfatted soaps with ether, in effecting the separation of mono- and di-glycerides. Also in cases—such, for instance, as in that of dirt retained by fat—where the solvent capacity of the soap solution seems insufficient, the capacity for forming emulsions may perhaps be assumed the cause of the action of soap.

Wool fat has recently been employed in the preparation of toilet soaps.

In the case of milled soaps it is advisable to stir the wool fat up beforehand with its own weight of water, rose water, or orange water until a uniform pale yellow pulp is produced. On adding this to the well-dried soap shavings, it supplies the mass and frees the finished soap from the presence of beads or streaks of fat, which are frequently observed when wool fat is used without any such preparation. The natural colour of the soap is a creamy tint; and this should not be more than assisted by a weak solution of pale yellow colouring matter, all high colours being avoided.

The proportions of the ingredients are as follows:—

Milled wool fat soap, with 10 per cent. of wool fat.—Well dried soap, 10 kilos; wool fat, 1200 grams—no soap should contain more than this proportion;—rose water, 600 grams; orange water, 600 grams, emulsified together, then mixed with the soap and allowed to draw overnight. The scent is compounded of geranium oil, 25 grams; lemon-rind oil, 10 grams; limetta oil, 10 grams; vanillin, 3 grams—as a 10 per cent. solution; oil of bitter almonds, 1.0 gram; violet root, 15 grams.

In the preparation of cocoanut-oil soap by the cold process, the great point is to get the right proportion of lye, 4–6 parts being taken to 10 of cocoanut oil. Wool fat, being un-

saponifiable, requires no lye ; immaterial whether it be added to the oil or employed in the new practical form of lana-cocoanut oil—100 parts of Cochin oil and 10 of wool fat.

Another recipe runs. The following ingredients—all well clarified—are carefully weighed out :—Cochin cocoanut oil, 5 kilos ; very pale palm oil,—air-bleached at a low temperature, —1 kilo ; wool fat for soap, 1 kilo. A highly caustic soda lye of sp. gr. 1·408, or 42° Be., is used,—2800 grams,—and is followed by 100 grams of water mixed with 100 grams of 28° Be. glycerin to improve the soap, this mixture being well stirred into the warm oil and lye. Finally, the scent—70–100 grams—and colouring, a pale cream yellow, are added. The mass soon thickens, and is then framed in order to secure uniform distribution and allow the air bubbles to rise. The frame is covered up and set in a warm place to induce a strong rise in temperature.

Bearing in mind the fact that almond oil is cheaper than olive oil, Schimmel & Co. have recommended the following method of saponifying the first-named oil by the cold process : After the sweet oil of almonds has been carefully saponified with weak lye, and salted out with a 20° Be. brine, 100 parts of oil are mixed with 34 of soda lye and 17 of potash lye (38° Be.) at 25° C., the mixture being kept stirred as in the case of cocoanut oil soap. The mixture must be stirred twenty-four to thirty-six hours, with intervals of rest, the stirring being performed for an hour after setting the ingredients in the pan (in the morning) ; thereafter the stirring is repeated for a few minutes at the end of every hour. The soap will not be ready for pouring by the evening, so that the pan must be left in a warm room all night, and then well stirred up the following morning, whereupon the soap will speedily thicken, and can be framed. Saponification sets in after another forty-eight hours, provided the mass is warm ; and at the end of thirty-six to forty-eight hours longer the soap will be hard.

This almond oil soap completely replaces olive oil soap for cosmetic preparations, such as Hungarian beard pomade. It dissolves in its own volume of distilled water to a fine clear paste, and is very agreeable to the skin.

Furthermore, 3 parts of cocoanut oil and 1 of sweet oil of almonds, saponified with a 38° Be. soda lye, furnish a very good lathering cocoanut-almond-oil soap, which seems to be very mild on the skin. This soap behaves in use like a good fat soap.

The following data are typical for the production of shaving soap from almond oil :—

Tallow, 85 parts, and sweet almond oil, 15 parts, are saponified with 25 parts of soda lye and an equal quantity of potash lye at 45–46° C. The saponification proceeds smoothly, and the resulting soap holds a brilliant fatty lather and is quite white. The bars do not undergo distortion, but remain rectangular. In view of its properties and the simple method of making, the operation is preferable to the plan of boiling, especially since the product fulfils all the requirements of a good shaving soap.

Shaving soaps by the cold process. The following recipes are given :—

1. Tallow, 29 kilos, and Cochin cocoanut oil, 28 kilos, are melted together, mixed with 15 kilos of sesame oil and 3 kilos of Venice turpentine, and saponified at 37°·5 C. (30° R.) by 37½ kilos of 38° Be. soda—from 120–125 per cent. caustic soda—by stirring in the usual manner; the soap is collected on the surface, framed, and well covered.

2. Tallow, 36; Cochin cocoanut oil, 16; soda lye (37° Be.), 22; potash lye (30° Be.), 6; Venice turpentine, 2.

3. Tallow, 40; cocoanut oil, 8; potash lye (35° Be.), 20.

4. Tallow, 34; lard, 12; cocoanut oil, 6; olive oil, 4; castor-oil, 4; glycerin, 4; soda lye (36° Be.), 28; potash lye (38° Be.), 1.

Scent.—Three parts each of aniseed oil and lemon oil, and 2 parts of lavender oil, employed in the proportion of 4–6 parts per 1000 of soap.

According to the Swedish patent (7398 of 1895) taken out by M. Zadig of Malmö, a hygienic soap can be prepared by adding 3–10 per cent. of cream to a neutral soap. The cream is previously emulsified with 3–10 per cent. of pure lanolin, and is run into the soap as the latter is passing through the mill. With regard to this recipe, it has been rightly observed that the hygienic action of cream cannot be very important. Is it not probable that in summer time the butter fat, which apparently is not saponified, but merely mixed with the soap, would very soon become rancid and spoil the soap?

The makers of Pfund's Milk Soap published three analyses, with reports by Bischoff, Haggenberg, and Fresenius, with regard to which Hefelmann made the following observations:—

Milk soap is neither entirely nor chiefly prepared from pure cow's milk, but is merely a mixture of 80 per cent. of soap and 20 per cent. of dry milk. The product lacks the good qualities of the best toilet soaps prepared according to definite technical principles. Such soaps should be perfectly neutral, pure curd soaps, of constant composition and containing at least 75–80 per cent. of combined fatty acids. Milk soap is not boiled to curd, it has a strongly acid reaction, is very irregular in composition, and, according to reports from various sides, contains only 52–60 per cent. of combined fatty acids instead of the 64 per cent. mentioned in the guarantee analysis. The deficit of 20 per cent. of combined fatty acids, or 22 per cent. of dry soap, is replaced by 20 per cent. of milk constituents; of these, however, the milk fat alone has any cleansing action, all the others—about 15 per cent.—being merely so much worthless ballast. The pure milk fat—5 per cent.—which is claimed to be so valuable, was found by Hefelmann and Haggenberg to be mixed with unsaponified soap fat, completely rancid, and consisting, to the extent of $\frac{4}{5}$ and $\frac{3}{5}$ respectively, of free fatty acids. The milk sugar had also undergone considerable decomposition, the natural ratio of milk

sugar to casein in milk being reversed in the milk soap. Hefelmann concludes that every good pure neutral toilet soap with 75-80 per cent. of combined fatty acids, is far superior to such a milk soap.

Soap spirit is one of those articles which are prepared with more or less difficulty. It is a prevalent opinion that the addition of spirit facilitates the saponification of oil. This, however, is not altogether correct; in contrast to caustic soda, the saponification of oil with a caustic potash of the prescribed purity is an impossible matter, the presence of at least a small quantity of certain salts—such as potassium carbonate or chloride—or alcohol being absolutely essential. The addition of alcohol must be confined within certain limits; for, if these be exceeded, the saponification is considerably retarded, in addition to which there is a considerable waste of the spirit.

A standard soap spirit may be easily prepared in the following manner, in a very short time and with very little waste of spirit:—

60 parts of oil are heated with 70 parts of caustic potash lye for ten to twelve minutes in a long-necked flask on the steam bath. 25 parts of spirit are then added in 2-3 portions, at intervals of a minute, and the flask is repeatedly tilted. After about five minutes the saponification will be complete, and the remainder of the spirit and water may then be added, care being taken to keep the flask in continuous motion after the addition of the alcohol, and to leave it only a few seconds at a time on the steam bath. The loss of spirit will then not exceed 1 per cent.

Solid soap spirit has been recommended by Vollbrecht for disinfecting the skin and hands; and though it would appear to be chiefly intended for medical use, it may also be employed in the ordinary care of the skin, especially in cases where efficient disinfection is in question.

60 grams of fine almond soap are dissolved in a sufficient quantity of 97 per cent. spirit with the assistance of the water

bath, and the solution is then made up to 1 litre with spirit. The mass solidifies on cooling, and can be cut into sticks or filled into tubes.

For purposes of disinfection, the hand or skin is rubbed over with this soap by a brush, but without water. The alcohol evaporates and leaves adhering to the skin a thin layer of soap, which, however, does not make the hand too slippery.

Bacteriological experiments show that this soap will give results comparable with those furnished by the officinal soap spirit. This soap has the advantage of high stability and cheapness.

Concerning the employment of violet root powder in the preparation of toilet soaps, it is advisable to purchase the violet root or powder, the latter for choice, from a firm of known reputation.

Good violet powder should readily form into pills or globules when rubbed between the thumb and finger. Any one desiring to reduce the root to powder himself must employ a specially constructed mill or a mortar; if the latter be used, it should be covered over with a piece of leather to prevent the dust flying about the room. In using violet root powder in the preparation of toilet soaps by the cold process, it should be stirred into the mill shortly before saponification, the lye not being added until after the powder and oil are uniformly mixed. For milled soaps, the powder is intimately mixed with the soap cuttings, and the mixture run two to three times through the mill.

To prepare violet root tincture from the roots, 1 kilo of the powdered root is shaken up with 2 kilos of 95-96 per cent. spirit, the flask being agitated several times daily, and the operation repeated for ten to twenty days. The tincture is then filtered off, and the powder treated over again in the same manner with another 2 kilos of similar spirit. This second extract has a finer and more agreeable odour, and is employed for the preparation of "extracts."

A new and special use for vaseline has been introduced by

the celebrated Viennese surgeon, Professor Gersuny, namely, as a subcutaneous injection for restoring plumpness to sunken portions of the flesh. Thus injected, vaseline retains its position, and does not affect the tissues. Gersuny has applied this treatment with success in several cases, and has reported on the same in the *Zeitschrift für Heilkunde*, 1900. Since that time the author has practised the method, and found it act satisfactorily in every case. The vaseline used is not the viscous kind, but a mixture of paraffin-vaseline oil, known as "Unguentum Paraffini"; and it has been demonstrated that paraffin, melting at 40° C., when injected into the corporeal tissues remains *in situ*, and does not produce any local irritation.

Gersuny applied this discovery to the production of a small swelling in such parts of the body where it seemed necessary to do so. Thus, for instance, in the case of a male subject whose upper jaw had to be removed, the consequent falling in of the cheeks was counteracted by this injection. Truly remarkable results have been attained by the new method as a substitute for sphincter muscles, a kind of flap valve being formed by the injected vaseline mixture. Paraffin injections have also been successfully practised for improving the articulation in cases of so-called wolf's throat; and the new method is also in use for suppling stiffened joints, the trials made in this direction having been crowned with success. Very good results have also been obtained with paraffin injections in rectifying deformities, replacing lost bone, filling out the depressions left by contracted scars, etc.

Whether this method can also be extended to the domain of cosmetics must be left undecided for the present, since, through the hardening of the paraffin, the injected portions do not exhibit the same softness and roundness as the natural flesh. Nevertheless, the method is one of high importance for surgico-cosmetic purposes as for surgery in general, and has certainly a promising future before it.

CHAPTER X.

COSMETIC SECRET PREPARATIONS FOR THE SKIN, COMPLEXION, TEETH, MOUTH, ETC.

IN many cases, cosmetic secret preparations are the cause why cosmetics do not enjoy the reputation that is really their due. Some of these preparations have been compounded in ignorance of the properties and action of the ingredients used; not infrequently these ingredients are directly contradictory among themselves; and oftentimes—which is the most to be deplored—they are directly injurious.

This is the domain of the empiric and ignorant manufacturer, since, on the one hand, hardly any of his customers is in a position to test the efficacy and harmlessness of the article; and, on the other, human vanity is exceedingly great with regard to the concealment of bodily defects. And the more that certain defects are declared incurable by serious science, the more the experienced cosmetic practitioner emphasises that the production of hair is an impossibility, that wrinkles are inevitable attendants on age, and that cuticular impurities are removed with greater certainty and less danger by a suitable mode of life; so much the more does ignorance turn towards quackery, and not infrequently such “patients” are put through treatments which any medical man would fear to apply, and which the patients themselves would not stand from any regular scientific practitioner.

An acquaintance with these secret preparations is necessary to the cosmetic manufacturer; in the first place, because he must

be informed on every point connected with his sphere of labour ; and, secondly, because he is in a position to eliminate the bad, replace it by good, and thus convert many worthless preparations into articles of utility.

SECRET PREPARATIONS FOR THE HAIR.

Fixoline.—A balsamic hair restorer, compounded of wax, lard, Peruvian balsam, alcohol, and inert aromatic substances. Is without effect.

Seebald's Hair Tincture consists of—Resorcin, 4·0 ; burnt sugar, 0·6 ; lemon oil, 2·5 ; Peruvian balsam, 1·5 ; 80 per cent. spirit, 92·0 parts. Is ineffectual.

Javol.—This is a wash for the head and hair, compounded by W. Anhalt of Colberg, and consisting of—Beef tallow, 1 gram ; lemon oil, 5 grams ; tincture of quinine, 15–20 grams ; potassium carbonate, 0·2 gram, and sufficient water to make up 100 grams. According to this formula, the preparation contains about 1·2 per cent. of a soap prepared with potash instead of soda.

Eau Sublime de Feuilles.—A hair dye made by Chrynet of Paris, is guaranteed innocuous by the maker ; but, being a 1½ per cent. aqueous solution of lead acetate with glycerin and sulphur, cannot be so considered. It is sold at the rate of 5 francs (4s.) per bottle (200 grams).

Seegar's Blonde Hair Dye consists, according to Hofmaas, of a solution of 1 per cent. pyrogallol and 1 per cent. cupric chloride. The brown and black dyes also contain the same ingredients in a more concentrated form. The preparation may be injurious to health.

Jonke's Colour Restorative for the Hair consists of an ammoniacal solution of silver nitrate, with an addition of alcohol and glycerin. It is neither better nor worse than many others of the same kind.

Esillag's Beard-forcing Pomade is an ordinary fat pomade,

with traces of bergamot oil, Peruvian balsam, and similar adjuncts.

Esillag's Tea Hair Wash consists of camomile flowers, and is of no importance.

Jonke's "Non plus ultra" Hair Dye is on the one hand a scented solution of pyrogallie acid, and on the other an ammoniacal solution of silver nitrate. No better or worse than numerous others.

Eau de Quinine Pinaud contains, according to Tscheppe, no trace of quinine or any other chinchona base or characteristic constituent of this bark. Metallic salts, tannin, salicylic acid, and cantharides are also entirely lacking.

Eau de Zenoble, a hair dye, chiefly consists of a solution of sodium hyposulphite, sulphate, and acetate, with free acetic acid, and a sedimental deposit of lead sulphide. It is probably prepared by precipitating a solution of lead acetate with dilute sulphuric acid, and dissolving the precipitate in sodium hyposulphite. The lead sulphide found results from spontaneous decomposition.

Krell's Beard Tincture (Fr. Netter) consists of linseed oil, castor-oil, charcoal, saltpetre, a little sulphur, and pounded bread crust, coloured either light or dark.

Vienna Cosmetic, Seegar's Hair Water, contains large proportions of iron, copper, pyrogallie acid, and free hydrochloric acid. May be dangerous to health.

Heiderich's Hair Fluid contains large quantities of lead acetate, and may be very dangerous to health.

Plumbiferous Hair Waters. The following plumbiferous preparations are sold in the Netherlands:—Allen's Hair Restorer, Ayro's Hair Invigorator, Eau de Castile, Eau d'Iuns, Eau de Florida, Eau Magique, Eau Hamilton, Eau d'Amerique, Melrose's New Hair Wash, New London Hair Restorer, Rosetter's Hair Restorer, Royal Windsor Hair Restorer. All of them are open to objections based on their constitution.

Hair Regenerator, a preparation sold in Austria, contains metallic ingredients.

Schneeweiss's Rapid Hair Dye contains copper, and may therefore be injurious.

Trixogene Hair Restorer is an alkaline liquid, containing small quantities of alcohol, ammonia, boric acid, salicylic acid, and glycerin, the great bulk, however, being merely water.

Aphrodite Hair Dye is an acid alcoholic liquid containing free hydrochloric acid, cupric chloride, ferric chloride, and pyrogallic acid. May be injurious to health.

Phoenix Hair Dye.—A case of poisoning following the use of this dye for a short time has been confirmed by Pollak; the symptoms, eczema and œdema, disappeared as soon as the preparation was discontinued. The toxic properties of the dye are attributable to paraphenylenediamine, which is a powerful blood poison, converting hæmoglobin into acid hæmatin.

Eckert's Hair Wash consists of water, alcohol, glycerin, lead acetate, fruity ether, and a sediment of undissolved lead acetate and precipitated sulphur. May be directly injurious to health.

Moustache Balsam, for promoting the growth of the beard, consists of fat, wax, and scent.

Scurf Essence (Lautenschlaeger, of Bischofsheim).—A scented ammonia soap, with alcohol and glycerin.

Oriental Depilatory consists of a mixture of liver of sulphur, calcium sulphide, calcium carbonate, and charcoal. There are many other better, cheaper, and more reliable means for attaining the same end.

Capillarine Hair Restorer (A. Altenkirck, Berlin) consists of absolute alcohol, onion juice, French brandy, Peruvian balsam, burdock-root oil, and juniper-berry oil, with an addition of fat or tallow.

Kneifel's Hair-forcing Tincture consists of ethereal oils, with extractive substances from onions and chinchona bark.

Ristoratere dei Capelli (Rizzi Frati, Florence) contains lead, and may therefore be injurious.

Braun and Jacoby's Eau Capillaire, advertised as an antiseptic cosmetic for promoting the growth of the hair, consists of—Quinine, 0·14 per cent. ; glycerin, 5 per cent. ; alcohol, 68·9 per cent. ; water, 25 per cent. ; Peruvian balsam, 2 per cent. ; and inert scented substances, with traces of lead. It does not fulfil its purpose in any respect, and, indeed, may be injurious.

New Balsamic Hair Dye.—This balsam contains 3·652 per cent. of lead acetate in addition to precipitated sulphur, scent, and small quantities of glycerin. May be injurious.

Erasmus Wilson's Hair Promoter consists of 300 parts of almond oil, 300 of ammonia, 500 of rosemary spirit, 60 of cantharides tincture, and 35 of lemon oil.

Io, a hair dye sold in Vienna, consists of paraphenylenediamine, and, like the similarly composed preparation "Phoenix," may be dangerous to health.

Örigo, a hair dye prepared by W. Jorgensen of Copenhagen, is free from lead, but consists of an ammoniacal solution of bismuth, containing a large amount of sulphur in suspension, a little glycerin, and strongly scented with rose oil. It is not uninjurious.

Kohol, advertised as a hair dye, contains Indian ink suspended in a solution of gum-arabic in thin quince mucilage. It is entirely harmless, but not permanent.

Aureol consists of 1 per cent. of methol, 0·3 per cent. of amidophenol hydrochlorate, and 0·6 per cent. of monoamido-phenylamine, the whole dissolved in 50 per cent. spirit containing 0·5 per cent. of sodium hyposulphite. The hair must, of course, be first cleansed of fat by soap or soda. The aureol is then mixed with an equal amount of hydrogen peroxide and applied to the hair with a fine comb. After two to three hours a dark brown colour is developed, which can be deepened by repeating the operation.

Glycoblazol (A. Sarg's Sohn & Co., Liesing, near Vienna) is a scented liquid, containing 35.22 per cent. by weight of alcohol, 61.64 per cent. of glycerin, and 0.19 per cent. of a substance—probably capsaicin—allied to cardol.

Seebald's Tincture for preventing Scurf and the Falling out of the Hair, consists of an extract of fresh orange peel in dilute alcohol, with 5 per cent. of resorcin and 3 per cent. of Peruvian balsam. This preparation has been found, in many instances, to produce reddening of the skin and pustules, and may therefore be injurious.

Pinaud's Eau de Quinine has the composition—Crameria tincture, 2; cantharides tincture, 1; alcohol, 50; lavender spirit, 5; glycerin, $7\frac{1}{2}$; and quinine sulphate, 1.

Sterling's Ambrosia is a North American hair dye, containing about 1 per cent. of lead acetate. May be injurious.

Hegewald's Antiphilathron, for preventing the falling out of the hair, is a filtered extract of 2.5 gall nuts with a mixture of 50 of strong spirit and 30 of water, scented with several ethereal oils.

Bittner's Aricine Pomade, for maintaining and invigorating the growth of the hair, is an ordinary pomade, and does not contain any aricine at all. It is sold at a price equal to 1s. per oz.

Fragelli's Beard Producer consists of a spirituous extract of lavender, benzoin, cinnamon, and santal-wood oil.

Royer's Beard-producing Pomade is an ointment compounded of 1 part of red chinchona powder and $1\frac{1}{2}$ parts of wax hair pomade. It is sold at a price equal to 1s. 6d. per oz.

Bergmann's Beard-producing Tincture is an alcoholic extract of tree bark, scented with a little oil of rosemary and thyme.

Royer's Beard-producing Tincture consists of 10 parts of common salt, 130 parts of imitation French brandy (containing fusel oil), and 2 of mace tincture. It is innocuous.

Elsner's Beard Dye consists of two liquids; the one contains

pyrogallic acid, and the other is an ammoniacal solution of silver nitrate.

Netter's Beard Tincture consists of linseed oil, castor-oil, charcoal, saltpetre, with a little sulphur and pounded bread crust. Over 4s. per bottle is the price charged for this "eminently rational" preparation.

Onion Solution for the Beard, which is claimed to be prepared from the juice of the onion, and to raise a strong, full beard on the chins of youths in their fifteenth year, consists of dilute scented spirit coloured with gentian tincture. The price is equal to 4s. 3d. per bottle.

Baume Circassienne, a Viennese hair dye, is a solution of lead acetate in aromatic water, with a small addition of sulphur. May be injurious.

Wortley's Berenizon, a stimulant for the hair, consists of Peruvian balsam, 3 parts; castor-oil, 3; cantharides, 4; alcohol, 85; and rose water, 40 parts.

Spring of Life, a Hamburg hair dye, contains lead acetate, and may be injurious.

Brilliantine, for beautifying the beard, is composed of castor-oil, 7; almond oil, 50; glycerin, 22; Jockey Club extract, $2\frac{1}{2}$; and spirit, 125 parts. Any fat would do just as well.

Pinaud's Brilliantine is a scented liquid which separates into two layers, and consists of about $\frac{3}{4}$ of olive oil and $\frac{1}{4}$ alcohol.

Wallritz's Cactus Pomade, for promoting the growth of the hair, is prepared by crushing 125 parts of cactus (spines and all) in a mortar, and boiling the same in a vessel, which must not be of copper, an addition of turmeric and indigo being made until a green colour is produced. The strained liquid is stirred until cold, with 750 parts of water, 60 of glycerin, 15 of tannin, $7\frac{1}{2}$ of rosemary oil, and 4 parts of fennel oil.

Lohse's Chromacome Hair Dye consists of two liquids, the one being a tincture of gall nuts, and the other a solution of iron acetate with a little silver nitrate.

Comachrome, for dyeing the hair black, is a solution of silver nitrate with pyrogallie acid.

Buehligen's Hair Preservative is a mixture of arnica tincture, 10 parts : glycerin, 5 : water, 60 : and alcohol, 10 parts.

Henry's Cosmetic for preventing the hair falling out or turning grey, consists of 180 parts of spirit, 3 of lemon oil, 1 part each of bergamot oil, rosemary oil, and lavender oil. The price charged for this "wonderful" but altogether inefficacious preparation is at the rate of 3s. for 4 oz., an extra charge of 9½d. being levied for the "rational" directions for use.

Funke's Crinine is an ammoniacal solution of silver nitrate.

Eau d'Afrique, for dyeing the hair black, consists of three liquids to be applied in succession: (1) A solution of 3 parts of silver nitrate in 100 of water; (2) an 8 per cent. aqueous solution of sodium sulphide; (3) a solution of silver nitrate, like the No. 1, but scented.

Eau d'Apollon is a scented aqueous solution of lead acetate, which is harmless.

Eau de Bahama, for dyeing the hair black, is a solution of lead acetate, containing sulphur in suspension, and scented with aniseed oil. This preparation may be injurious.

Kamprath & Schwartz's Eau de Capille hair dye is a mixture of 16 parts glycerin, 8 of sodium hyposulphite, 1 of lead acetate, 2 of precipitated sulphur, and 130 of water, scented with a small quantity of eau de Cologne. Containing lead, it may be injurious.

Bergmann's Ice Pomade, for curling the hair, is ordinary pomade.

Auricomus Golden Hair Wash is an aqueous solution of hydrogen peroxide.

Kreller's Milanese Hair Balsam consists of—Beef marrow, 40

parts; chinchona extract, 5 parts; 1 part each of Peruvian balsam, styrax, and bergamot oil, and $\frac{1}{2}$ part of lemon oil.

Migaree's Moustache Balsam is composed of fat and resin.

Pinkas' Invigorating Hair Oil is a solution of Peruvian balsam and extract of walnut shells, with a little cinnamon tincture, in strong spirit.

Karig's Crinochrome consists of a solution of 10 parts pyrogallie acid in a mixture of rectified wood vinegar and spirit of wine, 500 parts of each, and a solution of silver nitrate, 30 parts, in 900 parts of distilled water, with sufficient ammonia to redissolve the precipitate first formed.

Nieske's Patent Birch oil Balsam, advertised as an innocuous preparation, and as a solution of the pith of South American birch, really consists of an aqueous solution of lead acetate and a 10 per cent. precipitate of a mixture of lead carbonate and sulphur. It is therefore far from being innocuous.

Franz's Puritas Rejuvenator for Grey Hair consists of—Glycerin, 40 parts; water, 106; soda crystals, 3; cadmium sulphide, 15; and zinc sulphide, 1–3 parts.

Rostel's Recreation, for stimulating the growth of the hair, is a solution (10 per cent.) of glycerin and potash in water.

Polt's Reseda Curling Pomade is a simple pomade of yellow wax, cocoanut oil, olive oil, with an agreeable scent of mignonette and orange flowers.

Grolich's Roborantium, to remedy baldness, is dilute eau de Cologne with a little glycerin. The preparation has the highly valuable property of being absolutely innocuous. Pinkas' preparation, sold under the same name, for promoting the growth of the hair and beard, is a weak alcohol, with additions of sulphuric ether, acetic ether, liquidambar, rose water, and traces of clove oil and glycerin.

Milk of Roses Extract, a Viennese preparation for dyeing the hair, contains 10 parts of lead acetate, 15 of glycerin, 5 of sulphur, and 70 parts of an aqueous liquid scented with rose oil.

Lautenschlager's Scented Flavour is a scented rose soap, dissolved in alcohol and glycerin.

Liquid Shampoo contains 2 parts each of Quillaya bark extract and castile Soap, 1 of glycerin, 4 of alcohol, and 7 of rose water.

American Shampoo Liquid contains—Rosa, 1000 parts; spirit of wine, 120; ammoniacal tincture, 3; ammoniac carbonate, 5; and potash, 10 parts.

Shampoo Powder is a scented mixture of 22.5 Soda, 50 calcined soda, and 15 Quillaya.

Tincture de Lévrière, a so-called vegetable hair dye, consists of two liquors, one of which (a) is a solution of silver nitrate coloured with aniline blue, the other (b) a dilute solution of calcium pentasulphide.

Brogue's Tallow ("The Nearest and Best Hair Preservative") is a rose-scented liquid, consisting of 1 part of pure glycerin, 10 parts of water, and a little sulphur in suspension.

Allen's Wood's Hair Restorer for Renewing, Strengthening, Beautifying, and Curbing the Hair consists of—Sulphur, 56; lead acetate, 80; glycerin, 100; and aromatic water, 200 parts. Another preparation consists of—Prepared sulphur, 17 parts; cinnamon, 2; glycerin, 200; lead acetate, 26.5; and water, 630 parts.

Agnes Tamm's Depilatory for ladies chiefly consists of melted pine resin.

Mulder's Hair Ointment is an aqueous solution of rose leaves with about 5 per cent. of carbolic acid.

Kogel's Hair-growing Powder consists of rosewood fat, containing, in addition to scent, a little of a sulphur oil, presumably oil of mustard.

Seebald's Hair Tincture contains in 100 parts, 40 of resorcin, 9.6 of camphor, 2.5 of cedar oil, 1.5 of Peruvian balsam, and 92 of spirit.

SECRET PREPARATIONS FOR THE SKIN AND COMPLEXION.

Poudre Antiasthmatique Clery consists of a mixture of powdered sage leaves, belladonna leaf powder, and saltpetre. The belladonna is said to be efficacious, but the powder cannot fulfil its purpose.

Lewinsohn's Freckle Cream consists of 10·8 per cent. of "white precipitate," basic bismuth nitrate, lanolin, and ethereal oils. It cannot do what is claimed for it, and may be dangerous.

Lana Cream is a preparation bearing the trademark of the Norddeutsche Wollkaemmerei of Bremen, and, according to Lohmann, is a yellowish white ointment prepared by the aid of wool fat. The water incorporated with the wool fat contains free hydrochloric, sulphuric, and nitric acids, the latter especially in large proportion, and must therefore produce an injurious effect on the skin.

Grolich's Face Ointment (C. J. Dahms) is a white precipitate ointment mixed with bismuth white and scented with rose oil. May be injurious to health.

Legran's Freckle Ointment consists of a fat scented with rose water, and containing bismuth nitrate.

Spitzer's Face Pomade is a mixture of fats scented with rose water, and containing white precipitate. Probably injurious to health.

Milk of Lilies, a wash for the skin, contains rose water, glycerin, and magnesium silicate.

Poudre de Riz de Java (Bourgeois, Paris) contains 25·5 per cent. of zinc oxide and 74·5 per cent. of talc, but no trace of rice. The price charged is equivalent to 3s. 3d. per 3 oz. tin, whereas the real value is not more than sixpence. Moreover, the mixture may be injurious.

Crème Iris consists of 0·5 per cent. borax, 2 per cent. powdered talc, 10 per cent. zinc oxide, 87·5 per cent. glycerin

ointment, scented with tuberose extract. Is objectionable on account of the zinc oxide present.

Eau de Lys de Lohse has the following composition:— Powdered talc, 4 per cent. ; zinc oxide, 8 per cent. ; glycerin, 6 per cent. ; rose water, 82 per cent. Is objectionable by reason of the zinc oxide.

Gross & Reichardt's Betular is a scented liquid consisting of 11 parts of a solution of menthol-betulin, 19 parts boroglycerin, 20 parts birch tincture, and 50 parts aromatic birch water.

Crème Lefebvre, a remedy for freckles, is a yellow ointment composed of fat and wax, with a little sublimate. May be poisonous.

Eau de Circassiennes du Thompson, a remedy for freckles, consists of 30 parts zinc oxide triturated with 200 parts of scented water.

Vienna Face Water is composed of about 12-13 parts of commercial (plumbiferous) zinc oxide, $1\frac{1}{2}$ –2 parts of bismuth oxychloride, and 86–87 parts water, with a few drops of glycerin, rose water, and carbolic acid. It may be injurious.

Ayer's Recamier Cream is a mixture of zinc oxide with glycerin, scented with rose oil spirit, and contains a little mercuric chloride.

Rejuvenating Pills, said to be of Parisian origin, contain strychnine, and have been denounced by M. and A. Jolles of Vienna.

Spitzer's Freckle Ointment contains mercuric and mercurous salts.

Albadermine, a remedy for freckles, consists of 2 solutions : (a) contains 7.5 potassium iodide and 0.35 iodine dissolved in 12.5 glycerin and 120 of extract of rose leaves ; (b) is a solution of 75 sodium hyposulphate in 500 rose water.

Albion, a Parisian specific for wrinkles, and to whiten the

skin, is lead chloride and mercuric chloride in an aromatic water.

Fagner's Amandine is a cosmetic soap crême. 60 parts of powdered gum and 90 of powdered Venetian soap are mixed with 180 of honey and 1000 parts of almond oil, 5 egg yolks and 125 parts of milk of almonds are added by degrees. The whole is coloured with carmine and scented with oil of bitter almonds.

Praeger's Axeromaticum, a remedy for sweaty hands, is ordinary water coloured rose-red with carmine and scented with a little lemon oil. This "precious" remedy is sold at the rate of 1s. 6d. per 2 oz.

Baby Powder consists of 25 parts boric acid, 500 of wheaten flour, 440 of Venetian talc, and 10 of geranium oil.

Balsamisk Amikos, for the toilet, is a scented solution of borax.

Lengiel's Birch Balsam, a beautifier and specific against wrinkles, pockmarks, freckles, etc., consists of 5 parts sodium silicate (waterglass), 2 of potash, 1 of soap, 5 of gum arabic, and 10 of glycerin, in 400 parts of water; but no trace of birch sap. Any soap will work the same miracle without costing 4s. per bottle, like this preparation.

Bismuth Powder, for beautifying the skin, contains no bismuth, but consists of chalk and clay.

Blanc Camélia, a white cosmetic for the face, consists of 5 parts magnesium carbonate, 10 parts each of bismuth subchloride and precipitated chalk, and 75 parts of the whitest asbestos, with 0.5 per cent. of ultramarine.

Blanc de Perles, a beautifying wash, consists of scented water containing a little alcohol, and showing a dense white sediment of calomel and lead carbonate.

Pollan's Cosmetic Caustic, for warts, birthmarks, etc., is a mixture of equal parts of fused caustic potash and medicinal soap, with 8 parts of slaked lime.

Molfenter's Chemical Preparation for warts, is fuming hydrochloric acid.

Parisian Complexion Balsam, for rough and cracked skin, consists of equal parts of glycerin and egg albumin faintly perfumed. Of course, glycerin would be just as efficacious by itself.

Irvine's Compound Chinese Tablet of Alabaster, a skin beautifier, consists simply and solely of calcium carbonate.

Simmerling's Cosmeticum, for skin diseases, freckles, etc. Sweet almonds, 30 parts, and bitter almonds, 15, are pounded to an emulsion with 330 of water, then strained and mixed with 25 parts of benzoin tincture and 15 of lemon juice.

Cosmetic Vinegar is a mixture of 60 parts benzoin tincture, 10 parts Peruvian balsam, 150 each of eau de Cologne and Hofmann's balsam, and 300 parts of acetic acid, clarified by settling and filtration.

Rowland's Kalydor Cosmetic Wash, an American beautifier, is an emulsion of 30 parts of blanched bitter almonds and 500 parts of rose water, containing in solution 0.5 of mercuric chloride.

Rottmanner's Cosmetic Paste consists of a white wax ointment with a little sulphur, bismuth oxide, and zinc oxide.

Crème de Beauté, a beautifier, is an emulsion of sweet and bitter almonds; of course, without any action at all.

Crème du Liban Cosmetic is a mixture of olive oil, 35 parts, 5 parts each of wax and spermaceti, 5 of blanched sweet almonds, 20 of bismuth subnitrate, 30 of talc, 2 of benzoin tincture, 10 drops of Peruvian balsam, 5 drops each of rose oil and orange-flower oil, with sufficient olive oil to form a tough pulpy mass. Useless as a cosmetic.

Crème Simon à la Glycérine, a remedy for redness of the skin, blackheads, etc., consists of a mixture of zinc white, glycerin, and talc, scented with a little patchouli oil.

Pohlmann's Face Powder.—White lead, 14; talc, 7; mag-

nesia, 1; coloured with a little carmine, and scented with volatile oils.

Querloin's Eau Cosmetique is a mixture of bitter almond water, 30; rose water, 600; lead acetate, 1; spirit, 50; and benzoin tincture, 1.

Kreller's Eau d'Attirona, a beautifying soap, is an alcoholic extract of cloves and cinnamon, with 4 parts of soda soap and a few drops of peppermint oil.

Stenard's Eau de Beauté consists of 122 parts orange-flower water, 0.45 mercuric chloride, and 2.5 calomel.

Eau de Blanc de Perles contains 15 per cent. of white lead.

Eau de Castille, for freckles, contains sodium hyposulphite and lead acetate.

Eau de Naples, a beautifier, consists of—Borax, 12 parts; distilled water, 100; rose water, 50; camphor, 1; and benzoin tincture, 4 parts. Harmless.

Renard's Eau des Princesses consists of—Calomel, 2.5; corrosive sublimate, 0.45; orange-flower water, 122 parts.

Viola's Ephelidène, a beautifying wash and wrinkle smoother, is a slightly alkaline, weak alcoholic extract of styrax. Harmless.

Esprit d'Amaranth, for freckles, contains in bottle No. 3, which must be applied first—One part of corrosive sublimate and 30 of spirit (containing fusel oil); in bottle No. 2, over 2 parts of sublimate; and 3 parts of the same chemical in bottle No. 1.

Extrait de Camélias, a cosmetic largely sold in Paris, consists of white precipitate powder finely distributed in water.

Honora, an American beautifier, is a solution of 4.307 grams of potash in 50 c.c. of an extract of soap and violet root, with a little musk tincture.

Kaloderma, for rendering the skin soft and delicate, is composed of potash soap, glycerin, camphor, and comminuted

almonds. The same effect could be more cheaply produced with glycerin by itself.

Rowland's Kalydor, a toilet preparation, is compounded as follows:—100 parts of bitter almonds are made into an emulsion with 500 parts of rose water, the product being strained and mixed with 7·5 of sal-ammoniac and 15 of cherry-laurel water, followed by a solution of 0·10 sublimate in 15 of alcohol.

Candes & Co.'s Lait Antéphélique, a wash for freckles and other defects of the skin, contains 10 parts of corrosive sublimate, 1 of sal-ammoniac, 140 of egg albumin, 7 of lead sulphate, 2 of camphor, 840 of water.

Lait de Hébé, for freckles, is prepared in the following manner:—250 parts of Marseilles soap are dissolved in 500 of water containing a little potash, and the solution is treated with a solution of magnesium sulphate until the precipitation of soap ceases. The precipitate is collected, drained, and stirred up with 900 parts of 33 per cent. alcohol, 15 parts each of benzoic acid, sternalis oil, and lemon oil, 5 parts each of caraway oil, verbena oil, and oil of thyme, and 10 parts of Tolu balsam tincture being afterwards added. The preparation cannot produce the effect claimed.

Lilionese, a beautifier, consists of a saturated solution of potash, scented with a few drops of cinnamon oil and rose oil.

Gschleiden's Milk of Lilies is a mixture of rose water, glycerin, and magnesium silicate.

Schubert's Milk of Lilies is a solution of 3 per cent. of borax, 2 per cent. of benzoin tincture, and 95 per cent. of water.

Nieske's Menyl, a specific for redness of the nose, consists of—No. 1, a spirituous, slightly perfumed solution of benzoic acid, salicylic acid, and thymol; No. 2, of zinc white, talc powder, and a trace of phenol. Another preparation of the same name consists of an aqueous extract of violet root, with

a not inappreciable deposit of calomel. The first of these articles is entirely inefficacious, the other may be injurious.

Vogel's Odalin, a beautifying wash for freckles, moles, brittle skin, etc., is a solution of borax in glycerin water scented with rose oil, and coloured red with fuchsine.

Oleagine, a skin purifier, is an ordinary soap made of lard and soda, with starch and scented oils. Harmless.

Rother's Pasta Cosmetica, for pimples on the face, is an ointment of sulphur, styrax, and fat.

Pasta Mack, a toilet preparation, consists of 27 per cent. of rice meal and 73 per cent. of effervescing powder.

Rix's Pasta Pompadour, for freckles and moles, is a mixture of finely ground blanched almonds (bitter) and cold cream or wax pomade.

Lehmann's Pomade is a cosmetic compounded of almond oil, 20; white wax, 10; spermaceti, 5; bismuth sub-nitrate, 1; white precipitate, 1; glycerin, 3 parts, and a little scent.

Mueller's Poudre d'Adonis is claimed to impart to the face the delicacy of that of a girl of sixteen. It consists of 60 parts of prepared talc powder, coloured with a little madder lake, and slightly scented.

Bichler's Pilachrine, a specific for red noses, consists of water with about 10 per cent. of alcohol, and containing coarse sand and sulphur in suspension.

Cobbe's Rosalind, a New York cosmetic for tinting the face, fingers, and lips, consists of 10 parts of eosine, 30 each of white wax and spermaceti, and 410 of vaseline.

Rosy Cheeks without Rouge consists of powdered siliceous sponge. Under the microscope the preparation exhibits numerous siliceous needles, which penetrate the skin, producing mechanical irritation, and thus reddening the face.

Rouge de Théâtre, an American preparation for beautifying the skin, consists of talc, chalk, starch, and a little carmine to produce a red colour.

Sarah Bernhardt Powder La Diaphane consists of 50 grams each of powdered steatite and rice meal, and 28 of zinc white. The scent for the white powder is—Bergamot oil, 3 ; ylang-ylang oil, 2 ; neroli oil, 2 ; eau de Cologne, 20 grams : for red—Bergamot oil, 3 ; rose oil, 2 ; white rose extract, 2 grams ; 10 drops each of Ceylon cinnamon oil and musk tincture, coloured with carmine : for yellow—Bergamot oil, 3 ; clove oil, 1 ; cedarwood oil, 1 : and patchouli oil, 1 ; extract of new mown hay, 20 grams, the colouring matter being equal parts of light and dark cadmium yellow : for black—Portugal oil, 4 ; lavender oil, 1 ; petit grain oil, 1 : Ess Bouquet extract, 15 grams ; musk tincture, 20 drops : the colouring matter consisting of moist prepared vine black.

Viennese Face Wash consists of 12–13 parts of commercial zinc oxide, $1\frac{1}{2}$ –2 of bismuth oxychloride, and 86–87 of water, with a few drops of glycerin, rose oil, and a trace of carbolic acid.

Seiffert's Schneewittchen, a beautifier, is an aqueous solution of borax and glycerin, scented with rosewood or similar rose-perfumed oils.

Mizerski's May Dew Beautifier is prepared in the following manner:—100 parts of alum are dissolved in water and precipitated by a solution of 100 parts of soda crystals, the precipitate being washed with water and dissolved in 50 parts of concentrated vinegar, then mixed with 60–80 parts of damp, freshly levigated clay, followed by 30 of glycerin and sufficient water to make the whole up to 400 parts. A little eau de Cologne is added for its perfume.

Mueller's Oriental Milk of Beauty, a "world-famed" beautifier, for defects of the skin, consists of—Prepared talc, 12·5 ; glycerin, 10 ; borax, 0·5 ; eau de Cologne, 12 parts, with a little musk and 109 parts of water.

Kohlmann's Milk of Beauty.—Sweet almond oil, glycerin, and gum arabic, 7·6 of each, are emulsified with enough straw-

berry water to make 400 parts, 4 parts of Calidor essence and 15 of benzoin tincture being then added. The Calidor essence is composed of—Mace oil, 2; patchouli extract and jasmine extract, 12 each; Peruvian balsam and benzoin, 2 each; and Tolu balsam, 1 part.

Hudson's Beauty of Venus Paste is a very soft white ointment, consisting of 18 per cent. white wax, 4 per cent. stearin, $50\frac{2}{3}$ per cent. castor-oil, 18 per cent. glycerin, $1\frac{1}{2}$ per cent. precipitated sulphur, 5 per cent. moisture, and 3 per cent. perfumed oils, chiefly lemon oil and bergamot oil.

Lewicki's "Dew of Beauty" Water (maker's recipe).—Dew is collected on the grass in summer, distilled, and treated to an addition of 15 grams of pimpernels and 30 of powdered alum per $\frac{1}{4}$ litre of distillate, the whole being then boiled gently for an hour and strained through a close cloth. Four grams of calcined magnesia are partially dissolved in 50 of rectified spirit and 100 of the strongest wine vinegar, in such a manner that the undissolved portion of the magnesia forms with the vinegar a white mixture, which may be scented. This is mixed with the first liquid.

Rottmanner's Beauty Water, an "only reliable remedy for facial eruptions," consists of 1 part camphor and 5 of milk of sulphur in 50 parts of rose water.

Hoefeld's Freckle Specific is—(a) 15 grams of white precipitate ointment mixed with a little wax; (b) a wash containing 0.24 gram of camphor, 8.0 of benzoin tincture, 15.0 of soap spirit, and 125.0 of rose water.

Kimball's Freckle Specific is a solution of 1 part of potassium acetate in 50 parts of water, mixed with a little rose oil and alcohol.

Riedl's Freckle Ointment consists of—Paraffin, 18 parts; almond oil, 6; glycerin, 4; milk of sulphur, 2; colocynth tincture, 2; tannin, 1; with a few drops of thyme oil and rosemary oil.

Russ's Freckle Wash contains—Benzoin tincture, 1; Hofmann's balsam, 2; and rose water, 100 parts.

Sesser's Medicinal Toilet Mystery, for imparting a juvenile freshness and beauty to the complexion—(a) 60 grams of toilet wash cream, a mixture of fat, soap powder, and rose water; (b) toilet ice, a soft camphor-like ointment of fat, a little wax and spermaceti, and cocoanut oil; (c) 100 grams of herb milk, consisting of rose water, a good deal of prepared talc, and a little glycerin and zinc oxide; (d) 60 grams of Teint Poudre, consisting of a mixture of talc with a small quantity of white magnesia, and tinted with cochineal.

Tecklenburg's Milk of Venus consists of 5 parts of benzoin tincture and 200 of rose water.

Polt & Gruber's Venus Paste consists of 50 per cent. of a mixture of lard and cocoa butter, 6 per cent. of soap powder, 12 per cent. of glycerin, 3 per cent. of white levigated bole, 5 per cent. of gum arabic, $1\frac{1}{2}$ per cent. of styrax or benzoin, $2\frac{1}{2}$ per cent. of water; the whole scented with various ethereal oils.

Grecian Water, a cosmetic, consists of 189 parts of scented water with about $7\frac{1}{2}$ parts of white precipitate, and may be poisonous.

Manicere is a paste of zinc oxide and carmine, scented with bergamot oil.

Vincent's Poudre Refraichissante consists of tartaric acid and sodium bicarbonate.

SECRET PREPARATIONS FOR THE TEETH, MOUTH, ETC.

Odol.—According to the investigations of the Breslau Testing Station, this preparation contains—80 per cent. alcohol, 97 parts; salol, $2\frac{1}{2}$; saccharin, 0.004; peppermint oil, 0.5, with a trace of clove oil and caraway oil. The active ingredient is therefore salol, which has an antiseptic action.

Ulrich's Mouth Wash is an aqueous solution of sal-ammoniac

and potassium chloride, treated with a little spirits of camphor and alcohol until clear. It is of no real importance.

Rothe's Tooth Wash contains peppermint oil, carbolic acid, and strong alcohol. Is useless for the preservation of the teeth.

Harting's Mouth and Tooth Wash consists of eau de Cologne, alcohol, and carbolic acid. Is highly unsuitable for the purposes in view.

Detroit's Tooth Essence and Tooth Balsam contains peppermint oil and carbolic acid in addition to alcohol. Is highly unsuitable for the purposes specified.

Eau dentifrice aromatique pour entretenir la beauté des dents is a mixture of ratanhia extract, cinnamon, aniseed, and peppermint oil.

Sach's Mouth Wash consists of ratanhia extract, peppermint oil, myrrh, and alcohol.

Ebermann's Mouth Wash is a mixture of ratanhia extract, alcohol, clove oil, and peppermint oil.

Bergmann's Tooth Wash is a reddish liquid containing alcohol, peppermint oil, and aniseed oil.

Van Buskirk's Tragacanth-Sozodont consists of—Alcohol, 30 grams; water, 45; soap, $7\frac{1}{2}$ grams; and 2 drops of wintergreen oil. The soap is dissolved in the alcoholic mixture, then coloured with an extract of red santal wood, the wintergreen oil is added, and the liquid made up to 400 grams with water. The tooth powder used with this preparation consists of—Kieselguhr, 40 parts; powdered violet root, 125; and precipitated chalk, 250 parts, slightly scented with clove oil. Is, on the whole, an ordinary tooth soap; but the addition of kieselguhr is objectionable, owing to its attritive action on the teeth.

Cosmine.—This mouth wash has the following composition:—Formaldehyde, 0·327 per cent.; alcohol, 58·04 per cent. by weight; water, 41·0 per cent.; extract of myrrh and ratanhia,

0·32 per cent. : saccharin, 0·027 per cent. ; with a little ethereal oil. The liquid is of a red colour, and smells of peppermint and geranium oil. There are, however, much better and perfectly harmless mouth washes.

Abraham's Formagene, for filling decayed teeth, consists of two small bottles containing $2\frac{1}{2}$ grams, the one being filled with a mixture of creosote and clove oil, with a little phenol and alcoholic solution of formaldehyde. The other contains 2 grams of a yellowish white powder, free from organic matter, and setting like cement when mixed with water. It has the composition—Silica, 8·44 per cent. ; carbon dioxide, 19·84 per cent. ; alumina, 7·06 per cent. ; lime, 52·46 per cent. ; magnesia, 2·58 per cent. ; and zinc oxide, 7·52 per cent. The alumina is certainly present in the form of silicate, the magnesia and part of the lime as carbonates, and the rest of the lime—23·9 per cent.—as oxide. Traces of iron and potassium compounds are also detectable.

Agathodont Mouth Wash consists of an extract of 10·0 goosefoot, 5·0 Para cress, 3·0 pyrethrum in 100·0 spirit, with 1 per cent. of salicylic acid. The latter alone is of any efficacy, the other ingredients being worthless ; salicylic acid, however, is not to be recommended as a mouth wash.

Algotine, a mouth and tooth wash, is an aqueous solution of saltpetre, rendered aromatic by peppermint oil, myrrh tincture, and cinnamon tincture.

Popp's Anatherine Mouth Water.—Myrrh, 1 part ; guaiacum resin, 4 ; saltpetre, 1,—macerated in 120 parts of corn brandy and 180 of scurvy grass spirit for one night, then distilled to 240 parts of distillate ; to which are added 1 part each of rue, scurvy grass, rose leaves, black mustard, horse radish, pyrethrum, chinchona bark, lycopodium, sage, betiver root, and alkanna root. The whole is digested for fourteen days, then strained, filtered, and 1 part of spirits of nitric ether is added to each 120 parts of the filtrate. Notwithstanding the

trouble involved in its production, this preparation is of no importance.

Antiscorbutic Water, a mouth wash, is a mixture of 5 drops each of orange peel oil, mace oil, curly-mint oil, and sage oil, with 2 grams of mustard oil and 100 grams each of scurvy grass spirit and ordinary dilute alcohol.

Jackson's Balsam Water, for the mouth, is a distillate from benzoin, guaiacum resin, myrrh, tolu balsam, orange peel, angelica root, cinnamon, vanilla, and peppermint; coloured with alkanna red, and qualified with scurvy grass spirit. Despite—or rather, *because of*—the number of ingredients, the preparation is worthless.

Eau de Batot, a mouth wash, consists of 500 grams of cedar-wood tincture, 125 each of myrrh and ratanhia tincture, and 5 drops of peppermint oil.

Krohn's Fluid Ozone, a mouth wash, is an aqueous solution of sodium permanganate, contaminated with a little common salt and Glauber salt. There are far better mouth washes.

Pohlmann's Cosmetic Mouth Wash is prepared by digesting 8 parts each of aniseed and cinnamon, 4 parts each of guaiacum and benzoin, and $\frac{2}{3}$ part of pyrethrum with 240 parts of highly rectified alcohol for several days, then filtering and adding $\frac{1}{5}$ part of peppermint oil and 30 parts of scurvy grass water.

Nail Powder, for polishing the finger nails, is a scented mixture of 200 parts zinc oxide and 2 of carmine.

Kothe's Tooth Beautifier is a mixture of 1 part alum with 3 of calcium carbonate, scented with peppermint oil. When old, the mixture also contains gypsum. Is certainly not to be recommended for the teeth.

Müller's Antiseptic Tooth Water consists of $\frac{1}{4}$ part thymol, 3 parts benzoic acid, 2 parts eucalyptus tincture, and 750 parts water.

Kothe's Tooth Wash is a solution of 0.3 parts of salicylic

acid (containing carbolic acid) in 100 of 65 per cent. alcohol, a few drops of peppermint oil being added.

Pauli's Azymol is an antiseptic for the mouth and skin. According to the tests made at the Aufrecht Chemical and Bacteriological Institute, Berlin, it chiefly consists of a spirituous solution of salicylic acid, saccharin, vanillin, and menthol, presumably containing ratanhia tincture and peppermint oil, and coloured with fuchsin. A similar product would be furnished by the following recipe:—

Menthol	1·0 gram.
Peppermint oil	2·0 grams.
Saccharin	1·0 gram.
Vanillin	0·5 „
Fuchsin	trace.
Ratanhia tincture	4·0–5·0 grams.
96 per cent. spirit	92·0 „

CHAPTER XI.

TESTING AND EXAMINING THE MATERIALS EMPLOYED IN THE MANUFACTURE OF COSMETICS.

THE present section will deal with the testing of such materials as have not already been sufficiently treated of in this connection earlier in the book. A little repetition is inevitable.

Value is chiefly laid on the methods affording a ready means of detection and recognition.

Adulteration of olive oil with oils from the seeds of cruciferous plants can be detected by the sulphur present in the latter. With this object, 1-2 grams of caustic soda are dissolved in 20-30 grams distilled water, and heated to gentle boiling with 30 grams of the suspected oil, the mixture being stirred with a piece of bright silver plate. If cruciferous oils be present, the silver will become tarnished with black specks of silver sulphide.

As, however, it is often the custom to clean out olive oil casks with lyes containing sulphur, it is advisable, before applying the above test, to wash the oil with hot water, and examine the washings for sulphur. If the latter be present, it may be eliminated from the oil by repeated washings with hot water.

Commercial glycerin is somewhat impure, yellow, malodorous; and, since pure glycerin is preferable for most purposes, it may be purified in the following manner:—

The glycerin is first diluted with—preferably distilled—

water, then shaken up for some time with coarsely granulated animal charcoal, and filtered through loose asbestos, the filtrate being then concentrated to syrupy consistence on the water bath.

Pure glycerin should be inodorous, syrupy, and not darker than a dull amber yellow; nor should it redden blue litmus paper. Glycerin with an acid reaction is unsuitable for use. When heated on a clean platinum plate, it should volatilise completely, without leaving any residue. Glycerin contaminated with grape sugar or saccharose leaves a black carbonaceous residue, frequently of considerable amount. When glycerin is rubbed between the fingers it should not give off any fatty odour, nor should it turn brown or give a brown precipitate when treated with its own volume of concentrated sulphuric acid.

Glycerin adulterated with syrup is insoluble in two vols. of a mixture (1-2) of ether and alcohol, whereas pure glycerin is completely soluble therein.

Commercial glycerin frequently contains lime, which spoils it for use for many purposes, internal application in particular. This lime is due to the fact that in the saponification process the tallow is treated with a little lime and water under very high steam pressure in closed vessels, the resulting water separating from the lime soap being freed from lime by oxalic acid. The presence of lime in glycerin is detected by the following test: -The glycerin is treated with an equal volume of alcohol containing $\frac{1}{100}$ of sulphuric acid, and left for several hours; a precipitate indicates lime. Another test is by treating the glycerin with ammonium oxalate, which renders it turbid when lime is present.

In some cases commercial glycerin is purified by dilution with its own volume of water, and boiling for forty-eight hours with finely powdered litharge; then filtered, and the dissolved lead thrown down from the filtrate by sulphuretted hydrogen

gas, the treatment being completed by filtering through coarse bone black, and concentrating to syrup on the water bath. When the lead has not been completely removed, the glycerin will give a black or brownish turbidity with yellow ammonium sulphide, whereas pure glycerin remains clear.

The percentage of water in glycerin is determined by heating a given weight at 110° C. for an hour and then weighing.

Organic impurities can be detected by mixing the glycerin with 2 vols. of water and adding lead acetate. The amount of the precipitate will be in accordance with the quantity of impurities present.

As already stated, brown and yellow glycerin is adulterated with molasses, white glycerin with glucose. This can be detected by agitation with chloroform, the sugar settling down to the bottom, leaving the insoluble glycerin floating above.

Sodium bichromate gives no coloration when warmed with pure glycerin.

When shaken up with an equal volume of water, glycerin froths the more abundantly in proportion to the amount of impurities present.

For testing lanolin, Astolfi recommends a method based on comparative colorimetric tests, and capable of being extended to a quantitative process. He treats 0.1 gram of lanolin with 10 c.c. of concentrated sulphuric acid, and thus obtains an intensely red liquid with greenish fluorescence. When the lanolin is adulterated, the intensity of this coloration diminishes in proportion as the non-cholesterin adjuncts increase. On shaking the liquid up with 10 c.c. of chloroform, the latter assumes a bright red colour when the sample is pure, but is less red when water, extraneous fats, etc., are present. Also a black ring is observable at the plane of contact between the two liquids when the lanolin is pure, but is less decided in adulterated samples. On comparing these results with those furnished by pure wool fat under the same treatment, an

approximate idea of the extent of adulteration in any sample may be gained.

Testing the purity of Soap Powder.—A reagent is prepared by mixing 50 c.c. of 85 per cent. alcohol and 50 c.c. of concentrated acetic acid, 15 c.c. of which are then added to 1 gram of the soap powder, the whole being shaken up and heated to boiling.

Pure soap powder gives an almost clear solution, any impurities settling down to the bottom. The clear liquid is poured off and diluted with 2 vols. of water, whereupon the fatty acids collect as an oily layer on the surface. Soda, chalk, and magnesia, the usual adulterants, effervesce when dissolved in the acid reagent.

A simple method of testing the purity of benzine consists in dipping a cotton rag in the sample of benzine, and then allowing the latter to evaporate without moving the rag. If the benzine be pure, scarcely any smell will remain; but the longer and more strongly the cotton rag smells after the benzine has evaporated, the greater the impurity of the latter, and the less is it suitable for cosmetic purposes. Cotton will retain the smell of benzine far longer than wool.

Testing Bitter Almond Oil.—In addition to the particulars given in the section on materials, the following simple test may be mentioned:—A drop of the oil is applied to filter paper, which is then ignited, and the vapours caught in an inverted moist beaker. On rinsing out with water, filtering, and adding silver nitrate, a white precipitate will indicate adulteration with artificial benzaldehyde, which invariably contains chlorine.

A good test for artificial water of bitter almonds, based on the presence of chlorine products in commercial benzaldehyde, is afforded by a solution of hydrogen peroxide containing pure sodium peroxide, but free from hydrochloric acid. This reagent oxidises benzaldehyde to benzoic acid even at a moderate temperature, the latter then combining with the sodium, so that

even an aldehyde chlorinated in the benzol nucleus is retained. For this test 10 c.c. of the suspected water is mixed with 20 c.c. of hydrogen peroxide solution (free from HCl), and treated with 2-3 grams of pure sodium hydroxide free from chlorine, the whole being warmed to 40-50° C. for fifteen to thirty minutes, until the odour of bitter almonds has entirely disappeared. The liquid is then evaporated to dryness, gently incinerated, the residue taken up with water supersaturated with nitric acid, filtered, and the filtrate treated with silver nitrate solution. Water distilled from almonds will give not more than an insignificant opalescence under this test, even after long standing; but a decided chlorine reaction will be obtained if commercial benzaldehyde has been used. The same test may also be employed for testing bitter almond oil, 2-3 drops of the oil being taken.

The best water of bitter almonds is that prepared by long distillation, and containing 0.6-0.7 per cent. of benzaldehyde; since such a water will keep the longest, and will allow the action of the benzaldehyde-cyanhydrin to be exerted most completely.

Testing ethereal oil of Cherry Laurel and Bitter Almonds.—Hager recommends the following method for detecting adulteration with other ethereal oils, and distinguishing artificial water of bitter almonds:—

Five drops of the oil are dissolved in 2 c.c. of alcohol and treated with 2-3 drops of a 10 per cent. solution of mercurous nitrate. No reduction occurs in the case of natural or artificial cherry laurel oil or ethereal oil of bitter almonds; but when other oils are present, a grey precipitate is at once formed. This, however, does not apply to Ol. Cassian cin., Succini, Petran, Lithantracis, Vincæ, Vitis vinifer, and wintergreen oils,—these, therefore, and phenol being undetected by the above test. In such cases the almond oil under examination is dissolved in 6 vols. of dilute alcohol (sp. gr. 0.895), a turbidity then indicating adulteration with one of the oils mentioned.

Cherry laurel oil gives a clear solution with 3 vols. of the alcoholic solvent. Artificial water of bitter almonds—2–3 c.c.—will generally give with 3 drops of mercurous nitrate solution a faint white or grey turbidity, the distilled water giving an immediate dark grey coloration or turbidity with this reagent.

The purity of vaseline may be tested, according to Michle, by dissolving 2 grams in 5 grams of chloroform, shaking up with 10 c.c. of water, and adding 1 drop of phenolphthalein solution. Pure vaseline, free from acid and alkali, will not give any colour change, but a strong red coloration ensues on adding 1 drop of $\frac{n}{10}$ potash.

On stirring 10 grams of vaseline with 10 of water for a quarter of an hour on the water bath, leaving to cool, and pouring off the water, the latter should be perfectly neutral, give no sulphuric acid reaction with barium chloride, and leave no ponderable residue when evaporated on a watch-glass. Vaseline is browned by 98 per cent. sulphuric acid on the water bath, but weaker—73 per cent.—acid has no action except when a badly purified—technical—vaseline is present.

The best way is to melt 10 grams of the vaseline in the water bath, and add 50 drops of 73 per cent. sulphuric acid, continuing to heat and stir the mixture for a quarter of an hour. Pure vaseline will not alter the acid, neither colouring it brown nor forming a dark ring at the plane of contact on settling. If 5 grams of vaseline are heated half an hour, with stirring, on the water bath, along with 5 grams of sodium carbonate and 25 grams of water, the solution being poured off when cold and supersaturated with dilute hydrochloric acid, it will remain clear, provided neither resins nor fatty acids be present.

For the rapid determination of menthol in peppermint oil, Kleber recommends—

About 5 grams of the oil, accurately weighed, are treated with a known quantity—about 5 c.c.—of acetic anhydride, and boiled for half an hour under a reflux condenser. In the mean-

time a similar quantity of the acetic anhydride is titrated with normal caustic soda and phenolphthalein. After the boiled liquid has cooled down, the condenser is rinsed with water, which is added to the acetylated mixture, the latter being then titrated with normal caustic potash. The difference in the number of c.c. consumed in the two titrations when multiplied by 0.156 gives the amount of menthol in the oil. The entire operation does not take more than fifty minutes.

The following observations on the effect of various medicinal substances on the teeth have been published by Maurel :—

Copper sulphate turns the sound enamel of the teeth a muddy yellow colour ; it seems to attack the cement, and stains both this and the dentine a permanent green, which shows through the enamel.

Potassium chlorate has no action on the teeth.

Silver nitrate does not corrode the teeth, but imparts a black metallic sheen thereto.

Alum is very destructive to the enamel. Its action on the dentine and cement varies, and is but weak ; the cement is less affected than the dentine.

Alcohol has no action on dental enamel.

Benzoin tincture does not affect the structure, but, when constantly used, colours the dentine and cement brown.

Peppermint tincture does not affect the hard tissue, but stains the roots of the teeth brown.

Eau de Cologne has no action.

Tobacco juice does not act on the structure of the teeth, but stains them more or less brown. The enamel is only slightly affected.

Ordinary white wax paper for wrapping cosmetic preparations can be prepared in different ways, the simplest being to melt white wax at a temperature of over 100° C. on a brine bath, and when very fluid a fine soft paper is drawn quickly through the wax, or the latter is applied by the brush. This

gives the best, but most brittle, kind of wax paper. The brittleness may be diminished by mixing the wax with other suitable ingredients; such as 1 part of liquefied white Venice turpentine and $\frac{1}{2}$ part of bleached caoutchouc to 10 parts of white wax, the whole being melted on the water bath. This mixture, applied with a wide brush whilst hot, gives a paper that will not crack even when folded, and will stand moisture and water very well. When cold, the covering of wax can be beautifully polished by rubbing with a soft cloth.

Since it is impossible to print on waxed paper, the paper must be printed before waxing, water inks being generally used, since these are less affected by the hot wax than oil inks.

The following official (Excise) test for alcoholic perfumes and cosmetic washes, the alcohol content of which cannot be ascertained immediately, is prescribed in Germany:—

50 grams of the perfume, or the like, are well shaken up in a separating funnel with 50 grams of water and 50 of petroleum benzine (0.69–0.71 sp. gr.). After a rest of at least twelve hours, the gravity of the under stratum is determined with a Westphal balance or pycnometer at 15° C., and the amount of alcohol is calculated from the result, the percentage being obtained by multiplying by 2.

Should the perfume, etc., contain resin or other extractives, 50 grams are mixed with 50 grams of water, and from the mixture a quantity of not less than 90 grams is distilled, the distillate being made up to 100 grams with water, and treated as above. Acids, when present, are neutralised with caustic soda before the distillation. Preparations rich in glycerin—*e.g.* brilliantine—are diluted with 2 vols. of water, 150 grams of the diluted liquid being distilled to furnish about 100 grams, and made up to that quantity for treatment as in the other cases.

An empirical method of testing cosmetic liquids is by dilution with water to detect the oils by means of their divergent volatility. When a scented spirit or mouth wash is poured

into a wine-glass of water, the oils are immediately separated, and spread over the surface of the water. On leaving the liquid uncovered, the oils are volatilised in succession, the lightest first and the heaviest last. This operation may take some weeks, and, in order to be able to check the various stages, it is necessary to have several glasses, which must be marked with the dates in order to prevent confusion, the mixing being performed at definite intervals.

Take, for example, a mouth wash. To examine this, a fresh mixture is prepared, containing 10 grams of water and 10 drops of the liquid, every day for a week. At the end of that time there will then be a collection of seven mixtures, each of which will have a different scent, according to the volatility of the ethereal oils present. The qualitative composition of the mixed oils can be fairly well ascertained by this test.

In the foregoing case the predominance of peppermint oil soon disappears and other oils make their appearance, vanishing in their turn, so that the seven glasses form a whole gamut of characteristic odours, until, finally, the most stable of all are alone apparent. In this way, by a little experience, a bouquet can be easily analysed into its component scents.

Of course, the difficulty is greater when oils of equal volatility are present; nevertheless, even in this event a careful examination will enable a fairly reliable estimate to be formed.

The same method is also to some extent quantitative, so far as the comparison of scented liquids is in question.

According to the quantity of oils present, the greater or narrower will be the turbid zone upon the water; and although this may be disturbed by the presence of other substances, it nevertheless gives some idea of the quantity of the oils when the odour is also taken into consideration. Of course, the same proportions must be maintained in the test mixtures, and the number of drops of the essences carefully counted with a dropping-glass.

When it is believed that the different oils have been identified, a mixture is made in accordance with the recipe thus obtained : but alcoholic dilutions of the various oils are used to prevent the results being spoiled by an irremovable excess of any constituent.

The method, however, entails patience, time, and the possession of an experienced and critical faculty of smell ; and the odour of the mixture must be tested at frequent intervals, success being often attained suddenly.

THE END

INDEX.

- ABRAHAM'S Formagene, 229.
Absolute alcohol, 20, 21.
Absorption or enfleurage for obtaining perfumes, 16.
Acetic acid, detection of sulphuric or hydrochloric acid in, 30.
Acetic acid, glacial, 29.
Acetic acid, preparation of, 29.
Acetic ether, 32.
Acetic ether, decomposition of, 33.
Acetic ether, preparation of, 32.
Acetoxylol, 50.
Acetylene liberated from alcohol and calcium carbide, 20.
Acid, benzoic, 24.
Acid, pyrogallie, 37.
Adjuncts, cosmetic, 142, etc.
Adulterants for juniper oil, 65.
Adulterants of oil of cloves, 62.
Adulterated cochineal, 51.
Adulteration of cinnamon oil, 66.
Adulteration of lard, 73, 78.
Adulteration of lemon oil, 58.
Adulteration of oil of bitter almonds, 27.
Adulteration of oil of lavender, 60.
Adulteration of olive oil, 69.
Adulteration of patchouli oil, 65.
Adulteration of peppermint oil, 61.
Adulteration of Peruvian balsam, 79.
Adulteration of spermaceti, 72.
Adulteration of vaseline, 82.
Adulteration of wax, 71.
Æsthetic character of cosmetic art, 2.
Afrique, Eau d', 118.
Agathodont mouth wash, 229.
Agnes Teumel's depilatory, 217.
Ajowan oil, 54.
Ajowan oil, composition of, 54.
Albadermine, 220.
Albion, 220.
Alcohol, 20.
Alcohol, absolute, 20, 21.
Alcohol as an adulterant of oil of bitter almonds, 27.
Alcohol as a test for benzol and benzine, 24, 25.
Alcohol, deodorisation of, 21.
Alcohol, effect on teeth, 238.
Alcohol mouth wash, 98.
Alcohol, reaction with calcium carbide, 20.
Alcoholic perfumes, official test for, 239.
Alcohol rectified by potassium permanganate, 22.
Alcohol rectified by sodium peroxide, 22.
Alcoholic solution of thymol as mouth wash, 96.
Alcohol, solvent power of, 20.
Alcohol, table of specific gravity of, 22.
Algotine, 229.
Alkaline tests for musk, 48.
Alkanna, 147.

- Albanna red, 51.
 Allen's World's Hair Restorer, 217.
 Almond bran, 177.
 Almond oil, properties of, 67.
 Almond oil shaving soap, 203.
 Almond oil soap, toilet, 202.
 Almond oil, tests for purity, 67.
 Almond paste, 141.
 Almond soap *crème*, 141.
 Almonds, oil of, 66.
 Almonds (oil), oil of, 26, 51.
 Almonds (crushed), preparation of, 136.
 Alopecia areata, 93.
 Alum, effect on teeth, 238.
 Amandine, 141.
 Ambergis, 46.
 Ambergis, properties of, 47.
 American castoreum, 47.
 American face powder, 169.
 American hair washes, 154.
 American peppermint oil, 60, 64.
 American shampoo liquid, 217.
 Ammonia test for almond oil, 67.
 Ammonium carbonate, 26.
 Ammonium carbonate composition of, 26.
 Amyl valerianate, preparation, etc., of, 33.
 Anacardic acid, 116.
 Anathérine, Eau, 163.
 Aniline dyes for colouring, 146.
 Animal products used for cosmetics, 46, etc.
 Aniseed oil, 63.
 Antiscorbutic water, 230.
 Antiseptic hair washes, recipes for, 184, 187.
 Antiseptic mouth wash (Huchard's), 101.
 Antiseptic scurf water, 187.
 Antiseptic soaps, 179, etc.
 Antiseptic soaps, preparation of, 187, etc.
 Antiseptic sticking plaster, 127, 138.
 Antiseptic washes, 179, etc.
 Aphrodite hair dye, 211.
 Apparatus for absorption or effluage, 16, 17.
 Apparatus for distillation, 13, 14.
 Apparatus for extraction, 17.
 Apparatus for maceration, 15.
 Apple, essence, 116.
 Arsenicals and arsenic, volatility of, 4.
 Artificial bay rum, 140.
 Artificial essence of violets, 143.
 Artificial musk, 48, 49, 50.
 Artificial oil of violets, 43.
 Artificial oil of wintergreen, 41.
 Artificial vanilla essence, 143.
 Artificial vaselines, 82.
 Asiatic styrax, 79.
 Athenian hair wash, 183.
 Aureol, 152, 212.
 Ayer's Balm, 216.
 Ayer's Becaumier Cream, 219.
 Baby powder, 220.
 Baldness, 92.
 Baldness, Finsen light treatment for, 93.
 Baldness, Jackson's treatment of, 92.
 Baldness, Lassar's treatment of, 93.
 Baldness, neurotic, 93.
 Baldness, parasitic, 93.
 Balsam, Peruvian, 79, 185.
 Balsam soap, scent for, 125.
 Balsamisk, Amikos, 220.
 Bandoline, 132.
 Bandoline, rose, 140.
 Bandolines, 190.
 Barium sulphide paste, 121.
 Barium sulphide, preparation of, 121.
 Baume, 187.
 Baume, Circassienne, 214.
 Baume de Milan pour les cheveux, 187.
 Bay oil, 54.
 Bay oil, composition of, 54.
 Bay rum, 122.
 Bay rum, artificial, 140.
 Bay rum, preparation of, 123.

- Bay rum water, 182.
 Beard and moustache, wax for, 132.
 Beard forcing lotion, 187.
 Beard, specialties for, 157.
 Becchi and Welmann's tests for vegetable oils in lard, 75.
 Benzaldehyde or benzol hydride, 26.
 Benzine or benzol, 24.
 Benzine, rectification of, 25.
 Benzine, testing purity of, 235.
 Benzoic acid, 24.
 Benzoic acid, preparation of, 24.
 Benzoin, 80.
 Benzoin as a retarder of rancidity, 95.
 Benzoin, Calcutta, 8.
 Benzoin, Penang, 81.
 Benzoin, preservative qualities of, 24.
 Benzoin, properties of, 81.
 Benzoin, Siam, 80.
 Benzoin soap powder, 196.
 Benzoin, Sumatra almond, 80.
 Benzoin tincture, effect on teeth, 238.
 Benzol and benzine tested by iodine and alcohol, 24, 25.
 Benzol hydride or benzaldehyde, 26.
 Bergamot oil, 54, 63.
 Bergamot oil, composition of, 54.
 Bergamot oil, properties of, 54.
 Bergamot oil, tests for purity of, 55.
 Bergmann's beard-producing tincture, 214.
 Bergmann's ice pomade, 216.
 Bergmann's tooth wash, 228.
 Bernegan's toilet cream prepared with egg yolk, 127, 128.
 Bersch's recipe for pure fats for hair pomade, 94.
 Betulin, 145.
 Bichler's pilachrine, 224.
 Birch leaf and St. John's wort ointment, 114.
 Birch-tar oil, 55.
 Birch-tar oil, composition of, 55.
 Bismuth hair dye, 119.
 Bismuth powder, 220.
 Bismuth white, 45.
 Bismuth white, preparation of, 45.
 Bismuth white, solubility of, 45.
 Bitter almond, oil of, 26, 55.
 Bitter almond oil, testing purity of, 235.
 Bitter almond water, testing of, 236.
 Bittner's aricine pomade, 213.
 Black dyes for hair, 116-119.
 Blacks, 148.
 Blanc camélia, 220.
 Blanc de bismuth, 45.
 Blanc de perles, 45, 221.
 Blanc végétal, 107.
 Bleu végétal pour les veines, 113.
 Blonde hair dye, 119, 120.
 Blossom hair wash, 183.
 Blue (vegetable), for veins, 113.
 Blue (vein), for theatrical make-up, 114.
 Blues, 147.
 Boiry's cocoanut oil soaps, 193.
 Borate de Potasse, Eau de, 104.
 Borax, 29.
 Borax, colour of, 29.
 Borax, preparation of, 29.
 Borax soap, 194.
 Borax soap powder, 195.
 Borax, solubility of, 29.
 Borax tooth tincture, 104.
 Botot, Eau de, English recipe, 104.
 Botot, Eau de, French recipe, 103.
 Bottle, Florentine, 14.
 Bran, almond, 177.
 Bran, sand-almond, 176.
 Brilliant transparent pomade, 128.
 Brilliantines, 100, 132, 214.
 Brilliantine, recipes for, 158.
 Braun and Jacoby's eau capillaire, 212.
 Broers' harmless hair dyes, 150.
 Brown, 145, 147, 148.
 Brugier's Tolma, 217.
 Buccal cavity, care of, 95.
 Buchligen's hair preservative, 215.
 Burns and roughenings of skin, ointment for, 114.
 Butylxytol, 49.

Butyric ether, 33.
 Butyric ether, preparation of, 33.

CAKES, scent, 170.
 Cajeput oil, 63.
 Calcium carbide treated with alcohol, 20.
 Calcium carbonate, 46.
 Calcium carbonate, decomposition of, 46.
 Calcium carbonate, preparation of, 46.
 Calcium carbonate, solubility of, 46.
 Calcium pentasulphide, 38.
 Calcium sulphide, 37.
 Calcium sulphide, preparation of, 37, 120.
 Calcutta benzoin, 81.
 Calmus oil, 63.
 Caloderm, 111.
 Campeachy wood extract for colouring, 53.
 Camphor, 56.
 Camphor, crude, 56.
 Camphor, laurel, 56.
 Camphor powder, grated, 57.
 Camphor powder, preparation of, 57.
 Camphor (pure), properties of, 56.
 Camphor soap powder, 195.
 Camphorated vaseline ice, 128.
 Camphor-sulphur-Peruvian-balsam soap powder, 195.
 Camphor-sulphur soap powder, 195.
 Canadian castoreum, 47.
 Candes & Co.'s Lait Antéphélique, 223.
 Cantharides and glycerine hair wash, 183.
 Cantharides, injurious effect of, 179, 180.
 Cantharides pomade, 130.
 Capillarine hair restorer, 211.
 Caraway oil, 64.
 Carbolic powder, 172.
 Carbolic soap, 194.
 Carbon dioxide for enfleurage, 17.
 Carbon disulphide, 17.

Carbon disulphide, extraction by, 7.
 Cardol, 116.
 Care of buccal cavity, 95.
 Care of the hair, 92, 180.
 Care of the skin, 86, 87.
 Care of teeth, 95.
 Carmine, 146.
 Carmine or cochineal, 51, 53.
 Carmine, test for, 52.
 Carthamine red, 146.
 Cashew apple, 116.
 Cassia oil, 64.
 Castoreum, 47.
 Castoreum, American, 47.
 Castoreum, Canadian, 47.
 Castoreum, properties of, 47.
 Castoreum, Siberian, 47.
 Castoreum scent, initiation of, 8.
 Catarrh, menthol snuff for, 176.
 Chalk, levigated, 46.
 Chapped and cracked hands, 136.
 Chemical products used in manufacture of cosmetics, 19, etc.
 Cherry laurel and bitter almonds, testing ethereal oil of, 236.
 China pomade, 130.
 Chinchona bark, 186.
 Chinese gelatine, 91, 94.
 Chloral menthol, 175.
 Chloride of lime soap powder, 196.
 Chloroform, 23.
 Chloroform, decomposition of, 23.
 Chloroform, extraction by, 7.
 Chloroform, preparation of, 23.
 Chloroform salicylide, 23.
 Chromachrome, 215.
 Cinnamon cassia oil, 66.
 Cinnamon oil, adulteration of, 66.
 Cinnamon oil, properties of, 66.
 Cinnamon soap, scent for, 125.
 Citronella oil, 58, 64.
 Civet, 50.
 Civet, properties of, 51.
 Clove soap, scent for, 125.
 Cloves, oil of, 62.
 Cloves, oil of, 64.
 Coal-tar soap, 194.
 Cobbe's Rosalind, 225.
 Cochineal, adulterated, 51.
 Cochineal, preparation of, 51.

- Cochineal, properties of, 51.
 Cochineal (pure), tests for, 52.
 Cochineal, spent, 51.
 Cochineal or carmine, 51, 53.
 Cocoonut-almond oil soap, 203.
 Cocoonut oil for medicinal soaps, 193.
 Cocoonut-oil soap, 201, 202.
 Cold cream, Idelson's, 129.
 Cold cream, preparation of, 91.
 Cold process shaving soaps, 203.
 Colour of borax, 29.
 Coloured starch, 108.
 Colouring by aniline dyes, 146.
 Colouring by betulin, 146.
 Colouring cosmetic preparations, 145, etc.
 Colouring matter, violet, 145, 148.
 Colours, green, 147.
 Complexion and skin, secret preparations for, 218-228.
 Composition of ajowan oil, 54.
 Composition of bay oil, 54.
 Composition of bergamot oil, 54.
 Composition of birch-tar oil, 55.
 Composition of ionone, 42.
 Composition of menthol, 39.
 Composition of oil of bitter almonds, 55.
 Compound quinine and glycerine hair tonic, 154.
 Confectionery containing nitrobenzol, 32.
 Copper sulphate, effect on teeth, 238.
 Cosmetic adjuncts, 142, etc.
 Cosmetic art, aesthetic character of, 2.
 Cosmetic art, hygienic importance of, 2.
 Cosmetic pomades, 126-134.
 Cosmetic secret preparations, 208, etc.
 Cosmetic specialties, 142, 148, etc.
 Cosmetic toilet soaps, 200, etc.
 Cosmetic vinegar, 221.
 Cosmetic washes, official test for, 239.
 Cosmetics and medicine, 3.
 Cosmetics, animal products used in preparation of, 46, etc.
 Cosmetics, chemical products used in manufacture of, 19, etc.
 Cosmetics distinguished from perfumery, 1.
 Cosmetics, oils and fats used in preparation of, 54, etc.
 Cosmetics, testing materials employed in manufacture of, 232, etc.
 Cosmire, 229.
 Cracked and chapped hands, 136.
 Cream, cold, preparation of, 91.
 Cream, glycerin, 158.
 Creams for shaving, 164.
 Creams for washing, 166.
 Crème de Beauté, 221.
 Crème de Mauve, 140.
 Crème du Liban cosmetic, 222.
 Crème iris, 219.
 Crème Lefebvre, 219.
 Crème of almond soap, 141.
 Crème Simon à la Glycérine, 222.
 Crèmes, 91.
 Crude camphor, 56.
 Crushed almonds, preparation of, 136.
 Cumarin, 9, 29.
 Cumarin, preparation of, 29.
 Curling essence for hair, 155.
 Curling washes for hair, recipes for, 155.
 Curly mint oil, 64.
 DARKENING hair, preparations for, 116, etc.
 Darkening red hair, preparations for, 120.
 Decomposition of acetic ether, 33.
 Decomposition of ammonium carbonate, 26.
 Decomposition of calcium carbonate, 46.
 Decomposition of chloroform, 23.
 Decomposition of nitrobenzol, 31.
 Deodorisation of alcohol for cosmetic purposes, 21.
 Deodorisation of benzine, 25.
 Depilatories, 99, 117, 120-122.

- Depilatories, effect of, 100.
 Depilatories, recipes for, 161, 162.
 Depilatories, their advantages and disadvantages, 161.
 Depilatory (patent) of Huettemann and Zrzawy, 122.
 Detection of mineral acids in acetic acid, 30.
 Detection of nitrobenzol in confectionery, 32.
 Detection of nitrobenzol in liquids, 31.
 Detection of nitrobenzol in soaps, etc, 31.
 Detection of sulphuric or hydrochloric acid in acetic acid, 30.
 Dettroit's tooth essence and tooth balsam, 228.
 Diachylon plaister, substitute for, 137.
 Dieterich's preparation for promoting growth of hair, 156.
 Dieterich's recipe for Quillaya bark tincture, 126.
 Disagreeable smells, removal of, 135.
 Distillation for obtaining perfumes, 5, 12-15.
 Distillation of nitrobenzol, 31.
 Distillation, various appliances for, 13, 14.
 Distilled glycerin, 35.
 Dornier's cosmetic, 177.
 Dry rouge paste, 107.
 Dye for hair, blond, 119, 120.
 Dye for hair, harmless, 153, 156.
 Dye for hair, henna, 150.
 Dye for hair, walnut shell, 151.
 Dye, pomade, for beard and whiskers, 157.
 Dyeing the hair, 92.
 Dyes for hair, 99, 116, 150.
 Dyes for hair, black, 116-119.
 Dyes (tannin) for hair, 117.
 EAU Anathérine, 103.
 Eau d'Afrique, 118, 215.
 Eau d'Apollon, 215.
 Eau de Bahama, 215.
 Eau de Batot, 230.
 Eau de Blanc de Perles, 222.
 Eau de Borate de Potasse, 104.
 Eau de Botot, 98.
 Eau de Botot, English recipe, 104, 172.
 Eau de Botot, French recipe, 103, 172.
 Eau de Capille (Kamprath & Schwartz's), 215.
 Eau de Castille, 222.
 Eau de Circassiennes du Thompson, 219.
 Eau de Cologne, effect on teeth, 238.
 Eau de Cologne, Fehr's recipe for, 144.
 Eau de Cologne, imitation of, 9, 124.
 Eau de Cologne soap, scent for, 125.
 Eau de Hebe, 132.
 Eau de Lys de Lohee, 219.
 Eau de Naples, 222.
 Eau de Permanganate de Potasse, 104.
 Eau de Quinine Pinaud, 210.
 Eau de Salvia, 105.
 Eau de Violettes, 105.
 Eau de Zenoble, 210.
 Eau dentifricée aromatique pour entretenir la beauté des dents, 228.
 Eau Lenticuleuse, 133.
 Eau Sublime de Feuilles, 209.
 Ebermann's mouth wash, 228.
 Eckhart's hair wash, 211.
 Eclectic hair tonic, 155.
 Effect of depilatories, 100.
 Effect of medicinal substances on teeth, 238.
 Egg yolk as a cosmetic, 127, 128.
 Elaidin test for almond oil, 67.
 Elderberry colouring matter, 53.
 Elderflower scent, imitation of, 9.
 Elsner's beard dye, 214.
 Emulsions, 87, 88.
 Emulsions containing glycerin, 88.
 Emulsions, fats used in preparation of, 89.

- Emulsions, two groups of, 88.
 "Enfleurage" by carbon dioxide, 7, 17.
 Enfleurage or absorption, apparatus for, 16, 17.
 Enfleurage or absorption for obtaining perfumes, 16.
 "Enfleurage," perfumes obtained by, 7.
 English peppermint oil, 60, 64.
 English smelling salts, 168.
 Erasmus Wilson's hair promoter, 212.
 Errors in choice of tooth powders, 96.
 Esillag's beard - forcing pomade, 209.
 Esillag's tea hair wash, 210.
 Esprit d'Amaranth, 222.
 Ess Bouquet, imitation of, 9.
 "Essence de Lavande Mont Blanc," 59.
 Essence of jargonelle pear, 33.
 Essence of violets, artificial, 143.
 Essences, scent, 124.
 Essences, soap, 124, 167.
 Ether, acetic, 32.
 Ether and sulphuric acid, mixing of, 19.
 Ether, butyric, 33.
 Ether, nitrous, 34.
 Ether, sulphuric, 19.
 Ether, valerianic, 33.
 Ethereal oil of cherry laurel and bitter almonds, testing of, 236.
 Ethereal oils (light), adulterating oil of bitter almonds, 27.
 Ethereal oils, tests for, 63.
 Eucalyptus oil, 58, 64.
 Eucalyptus tooth paste, 103.
 Experiments by Dr. Karl Roese on value of mouth washes, 97.
 Experiments on soaps, Rotondi's, 188, etc.
 Expressed liquid, separation of, 12.
 Extract of heliotrope, 9.
 Extract of walnut shells, 151.
 Extraction, apparatus for, 17.
 Extraction by carbon disulphide, 7.
 Extraction by chloroform, 7.
 Extraction by petroleum ether, 7.
 Extraction by sulphuric ether, 7.
 Extraction, methods for obtaining perfumes by, 17.
 Extraction, perfumes obtained by, 5, 7.
 Extrait de Camélias, 223.
 Extrait végétal, 141.
 FACE powder, American recipes for, 109.
 Face powder, rose, 110.
 Face powder, salicylic acid, 99.
 Face powder, white, 110.
 Face powders, 162.
 Face powders, preparation of, 105, etc.
 Fagner's Amandine, 220.
 Falling of hair, causes of, 180.
 Fat in hair, 181.
 Fats for hair pomade, 94.
 Fats used in preparing emulsions, 89.
 Fehr's recipe for Eau de Cologne, 144.
 Fennel oil, 64.
 Finsen light treatment for baldness, 93.
 Fixateur à la rose pomade, 130.
 Fixoline, 209.
 Florentine bottle, 14.
 Formalin mouth wash, 101.
 Formalin tooth paste, 101.
 Fragelli's beard producer, 213.
 Frangipani sachet powder, 141.
 Franz's Puritas rejuvenator for grey hair, 216.
 Freckle washes, 132, 133, 134.
 Freckles and moles, preparations for removal of, 98.
 Freckles, preparations for, 132-134.
 French Eau de Botot, 172.
 French milk added to ethereal oils, 10.
 French smelling salts, 168.
 French white, 42.
 Frohman's tooth soap, 102.
 Frothing mouth wash, 159.

- Fumigating paper, incombustible, 140.
 Fumigating papers, 138, 139.
 Funke's Crinine, 215.

 GAULTHERIA oil, 40.
 Gaultheria oil, preparation of, 40.
 Gaultheriline, 41.
 Gelatine, Chinese, 91, 94.
 Gelées, glycerin, 127.
 General cosmetic preparations, 85, etc.
 Geranium oil, 64.
 Geranium soap, scent for, 125.
 German peppermint oil, 60.
 Glacial acetic acid, 29.
 Gloss, skin, 111.
 Glycerin, 34.
 Glycerin and Cantharides hair wash, 183.
 Glycerin as a preservative, 88.
 Glycerin cream, 158.
 Glycerin, dilution with water, 35.
 Glycerin, distilled, 35.
 Glycerin for maceration, 16.
 Glycerin for skin, 87.
 Glycerin gelatine, 158.
 Glycerin gelées, 127.
 Glycerin hair restorer, 123.
 Glycerin in aqueous solutions of different densities, 35.
 Glycerin in emulsions, 88.
 Glycerin, maceration by, 6.
 Glycerin, preparation of, 34.
 Glycerin, purification of, 232-234.
 Glycerin solution, refining of, 34.
 Glycoblazol, 213.
 Grease powder, manufacture of, 105.
 Grecian water, 227.
 Green colour of olive oil, 70.
 Green colours, 147.
 Grolich's face ointment, 218.
 Grolich's Roborantium, 216.
 Gross & Reichardt's Betulinar, 219.
 Growth of hair, preparations for promoting, 156.
 Gschleiden's milk of lilies, 224.

 HAIR, brilliantine for stiffening, 158.
 Hair, care of, 92, 180.
 Hair, causes of falling out, 180.
 Hair curling essence, 155.
 Hair curling washes, recipes for, 155.
 Hair dye, bismuth, 119.
 Hair dye from walnut shells, 151.
 Hair dye of henna, 151.
 Hair dyes, 99, 117, etc.
 Hair dyes, black, 116-119.
 Hair dyes, (harmless), Broers', 150.
 Hair dyes, tannin, 117.
 Hair, fat in, 181.
 Hair pomade, Bersch's recipe for pure fats for, 94.
 Hair pomades, 94.
 Hair, preparations for darkening, 116, etc.
 Hair, preparations for promoting growth of, 156.
 Hair regenerator, 211.
 Hair restorer, glycerin, 123.
 Hair restorers, 117-122, 123, 186, 187, etc.
 Hair, secret preparations for, 209-218.
 Hair, soap for washing, 94.
 Hair tonic, eclectic, 155.
 Hair tonic, quinine, 155.
 Hair wash (superior), 122.
 Hair washes, American, 154.
 Hair washes (antiseptic), recipes for, 184, 185.
 Hair washes (medicinal), recipes for, 184.
 Hair washes (ordinary), recipes for, 182, etc.
 Hair washes, specialties in, 150.
 Hair, washing of, 92.
 Halogen acid, 50.
 Hands, cosmetic for, 160.
 Hands, pâte au musc for washing, 111.
 Hands, protective wash for, 135.
 Hands (sweaty), remedy for, 135.
 Hard paraffin, 78.
 Harmless hair dye, 153, 156.
 Harmless hair dye, Joseph's, 156.

- Harmless hair dyes, Broers', 150.
 Harting's mouth and tooth wash, 228.
 Hebe, Eau de, 132.
 Hefelmann's observations on milk soap, 204.
 Hegewald's Antiphilathron, 213.
 Heiderich's hair fluid, 210.
 Heliotrope extract, 9.
 Heliotropin, 9, 36.
 Heliotropin, artificial, 36.
 Heliotropin, preparation of, 36.
 Henna as hair dye, 150.
 Henry's cosmetic for hair, 215.
 Herb soap, scent for, 125.
 Hermite's thymol solution for mouth wash, 102.
 Hoefeld's freckle specific, 227.
 Honora, 223.
 Hot milk added to ethereal oils, 10.
 Hübl or iodine, value in adulterated lard, 77.
 Huchard's antiseptic mouth wash, 101.
 Huchard's salol tooth powder, 102.
 Hudson's beauty of Venus paste, 226.
 Huettemann and Zrzawy's patent depilatory, 122.
 "Huiles antiques," or pomades, 15.
 Hungarian pomade, 131.
 Hydraulic press for extracting perfumes, 5, 11.
 Hydrochloric acid in acetic acid, detection of, 30.
 Hydrogen peroxide for delicate skins, 114.
 Hygienic, importance of cosmetic art, 2.
 Hygienic soaps, 166, 200, etc.
 ICE, camphorated vaseline, 128.
 Ice, menthol, 176.
 Idelson's cold cream, 120.
 Imitation of castoreum scent, 8.
 Imitation of Eau de Cologne, 9, 124.
 Imitation of elderflower scent, 9.
 Imitation of Ess Bouquet, 9.
 Imitation of lily of the valley scent, 9.
 Imitation of moss rose perfume, 9.
 Imitation of smell of poplar buds, 8.
 Imitation of strawberry smell, 8.
 Imitation of white rose perfume, 8.
 Incombustible fumigating paper, 140.
 Indigo-carmine, 147.
 Ingredients used in preparation of cosmetics, 4.
 Injected vaseline to restore plumpness, 206, 207.
 Injurious effect of cantharides, 179, 180.
 Instantaneous mentholine, 175.
 Io, 212.
 Iodine as a test for benzol and benzine, 24.
 Iodine or Hübl value in adulterated lard, 77.
 Iodoform, removal of smell of, 135.
 Ionone, 42.
 Ionone, composition of, 42.
 Ionone, preparation of, 42, 142.
 Ionone, properties of, 43.
 Irvine's compound Chinese tablet of alabaster, 221.
 Isobutyl alcohol, 50.
 Isodibutylene acid, 50.
 Isopropyltoluol, 50.
 JACKSON on care of hair, 93.
 Jackson's balsam water, 230.
 Jargonelle pear, essence of, 33.
 Javol, 209.
 Jonke's colour restorative for the hair, 209.
 Jonke's "Non plus ultra" hair dye, 210.
 Joseph's hair dye (harmless), 156.
 Juniper oil, 65.
 Juniper oil, adulteration of, 65.
 Juniper oil, properties of, 65.
 KALODERMA, 223.
 Kalodont, 172.
 Karig's Crinochrome, 216.
 Karl Roese's experiments on value of mouth washes, 97.
 Kimball's freckle specific, 227.

- Kniefel's hair-forcing tincture, 212.
 Kogler's hair-renewing oil, 218.
 Kohol, 212.
 Kosmin, 98.
 Kothe's tooth beautifier, 231.
 Kothe's tooth wash, 231.
 Köttstorfer method of finding saponification value in adulterated lard, 77.
 Krell's beard tincture, 210.
 Kreller's Eau d'Attirona, 222.
 Kreller's Milanese hair balsam, 216.
 Krinochrome, 118.
 Krohn's fluid ozone, 230.
- LANA cream, 218.
 Lait de Hébé, 223.
 Lang's table of glycerin in aqueous solutions of different densities, 35.
 Lanolin, 83.
 Lanolin, manufacture of, 105.
 Lanolin milk, 127.
 Lanolin, properties of, 83.
 Lanolin rose cream, 152.
 Lanolin, testing of, 84, 234.
 Lanolin toilet cream, 126.
 Lard, 72.
 Lard (adulterated), Hübl or iodine value, 77.
 Lard (adulterated), saponification value, 77.
 Lard, adulteration of, 73-78.
 Lard, properties of, 72.
 Lard, test for purity of, 72.
 Lard, tests of Becchi and Welmanns, for vegetable oils, 75.
 Lassar's treatment of baldness, 93.
 Laurel camphor, 56.
 Laurel hair wash, 182.
 Laurel tincture, 140.
 Lautenschloeger's scurf essence, 217.
 Lavender, oil of, 59, 64.
 Leaf rouge, 113.
 Leather, Spanish, 168.
 Legran's freckle ointment, 218.
 Lehmann's pomade, 224.
- Leistikow's pomade for preventing fall of hair, 157.
 Lemon oil, 58, 64.
 Lemon oil, adulteration of, 58.
 Lemon oil, properties of, 58.
 Lemon-grass oil, 58.
 Lengiel's birch balsam, 220.
 Lenticuleuse, Eau, 133.
 Levigated chalk, 46.
 Levigated chalk as tooth powder, 97.
 Lewicki's dew of beauty water, 226.
 Lewinsohn's freckle cream, 218.
 Light treatment (Finsen) for baldness, 93.
 Lilionese, 133, 224.
 Lily of the valley scent, imitation of, 9.
 Lip colouring matter, 112.
 Lip salve (excellent), 129.
 Liquid medicated soap, Regnault's, 197.
 Liquid pearl white, 112.
 Liquid soap, Robiquet's, 197.
 Liquid soap, Saidemann's, 197.
 Liquid shampoo, 217.
 Liquid soaps, 196, etc.
 Liquid sulphur soap, 199.
 Liquid vegetable rouge, 112.
 Liquids containing nitrobenzol, 31.
 Liquidambar, 79.
 Liver of sulphur, preparation, etc., of, 37.
 Lohse's chromacome hair dye, 215.
 Lotion for forcing beard, 187.
- MACE oil, 64.
 Maceration, apparatus for, 15.
 Maceration by glycerin, 6, 16.
 Maceration for obtaining perfumes, 5, 6, 15.
 Madder red, 53.
 Manicere, 228.
 Manufacture of cosmetics, chemical products used in, 19.
 Marble soap, Schleich's, 199.
 Marble-dust soap, 166.
 Marseilles olive-oil soap, 196.

- Marshmallow soap, scent for, 125.
Massage cosmetic, 177.
Materials employed in manufacture of cosmetics, testing, 232, etc.
Mauve, Crème de, 140.
Meals and pastes, 90.
Medicinal hair washes, recipes for, 184.
Medicinal soaps, powdered, 195.
Medicinal soaps, recipes for, 194.
Medicinal soaps, Unna's, 191.
Medicinal sulphur and tar soaps, 198.
Medicine and cosmetics, 3.
Melanogene, 118.
Menthol, 39.
Menthol as a solvent, 39.
Menthol, chloral, 175.
Menthol, composition of, 39.
Menthol ice, 176.
Menthol in oil of peppermint, 40.
Menthol migraine ointment, 175.
Menthol snuff for catarrh, 176.
Menthol soap powder, 196.
Menthol preparations, 174.
Menthol, properties of, 39.
Menthol toothache drops, 175.
Menthol vinegar, 175.
Mentholine, instantaneous, 175.
Messina oil, 54.
Methylsalicylic ether, 41.
Migaree's moustache balsam, 216.
Migraine menthol ointment, 175.
Milan, Baume de pour les Cheveux, 187.
Milk (fresh) added to ethereal oils, 10.
Milk (hot) added to ethereal oils, 10.
Milk of lilies, 219.
Milk of roses, 141.
Milk of roses extract, 217.
Milk soap, Hefelmann's observations on, 204.
Milk, vegetable, 90.
Milled soaps, 201.
Mineral acids in acetic acid, detection of, 30.
Mixing of ether and sulphuric acid, 19.
Mizerski's May dew beautifier, 226.
Moles, preparations for, 132-134.
Moles and freckles, preparations for removal of, 98, 134.
Molfentner's chemical preparation, 221.
Moss rose perfume, imitation of, 9.
Moustache and beard, wax for, 132.
Moustache balsam, 211.
Mouth, secret preparations for, 228, 231.
Mouth wash for smokers, 96.
Mouth wash (frothing), 159.
Mouth wash of alcoholic solution of thymol, 96.
Mouth wash of 40 per cent. alcohol, 98.
Mouth washes, 96, 101, etc.
Mouth washes, Dr. Karl Roese's, experiments on value of, 97.
Mouth wash tablets, 159.
Mueller's Oriental milk of beauty, 226.
Mueller's poudre d'Adonis, 224.
Mulder's hair balsam, 218.
Müller's antiseptic tooth water, 231.
Musk, 47.
Musk, artificial, 48, 49, 50.
Musk paste for washing the hands, 111.
Musk, properties of, 48.
Musk, substitute from turpentine, 49.
Musk, tests with alkalis for, 48.
Musk tincture, 144.
NAIL powder, 231.
Naphthalene soap powder, 196.
Naphthol hair water, 184.
Naphthol soap powder, 195.
Naphthol-sulphur soap powder, 195.
Nerolin, 9.
Nerve pencil, 159.
Nervous headache, Leistikow's pomade for preventing fall of hair through, 157.
Netter's beard tincture, 214.
Neurotic baldness, 93.

- Neutral soaps, 188, etc.
 New balsamic hair dye, 212.
 Nicotine soap, 166.
 Nieske's menyl, 224.
 Nieske's patent birch-oil balsam, 216.
 Nitrobenzol as an adulterant of oil of bitter almonds, 27.
 Nitrobenzol as a perfume for fats and soaps, 31.
 Nitrobenzol, decomposition of, 31.
 Nitrobenzol, distillation of, 31.
 Nitrobenzol, light, 31.
 Nitrobenzol in confectionery, a test for, 32.
 Nitrobenzol in liquids, a test for, 31.
 Nitrobenzol in soaps, etc., 31.
 Nitrobenzol, preparation of, 30.
 Nitrous ether, 34.
 Nitrous ether, preparation of, 34.
 Nitrous ether, properties of, 34.
 Nutmegs, 80.
 Nutmegs, good and bad, 80.

 ODOL, 98, 228.
 Odontine, 170, 171.
 Oil for hair, walnut, 151.
 Oil of almonds, 66.
 Oil of bitter almonds, 26, 55.
 Oil of bitter almonds, adulteration of, 27.
 Oil of cloves, 62.
 Oil of cloves, adulterants of, 62.
 Oil of cloves, preparation of, 62.
 Oil of cloves, properties of, 62.
 Oil of cloves (pure), tests for, 62.
 Oil of lavender, 59.
 Oil of lavender, adulterants for, 60.
 Oil of lavender, properties of, 59.
 Oil of lavender, tests for, 60.
 Oil of peppermint, 60.
 Oil of peppermint, menthol in, 40.
 Oil of violets, artificial, 43.
 Oil of wintergreen, 40.
 Oil of wintergreen, artificial, 41.
 Oil of wintergreen, preparation of, 40.

 Oils and fats used in cosmetics, 54, etc.
 Oils, scented, and pomades, 6.
 Ointment for burns and rough skin, 114.
 Ointment of birch leaf and St. John's wort, 114.
 Oleagine, 224.
 Oleolisse Tonique de Piver, 140.
 Oleum eucalypti, 59.
 Oleum eucalypti Australe, 58.
 Olive oil, 69.
 Olive oil, adulteration of, 69, 70.
 Olive oil, green colour of, 70.
 Olive oil soap, Marseilles, 196.
 Olive oil, testing of, 232.
 Onion solution for the beard, 214.
 Orange-flower oil, 64.
 Ordinary powder, 163.
 Oriental depilatory, 211.
 Origo, 212.

 PALM oil, 146.
 Palmarosa oil, 64.
 Paper [plaisters for cosmetic purposes, 136.
 Paper (white wax), preparation of, 238.
 Papers, fumigating, 138, 139.
 Paraffin, 78.
 Paraffin, hard, 78.
 Paraffin, properties of, 78.
 Paraffin tablets, scented, 177.
 Paraisobutylxylol sulphonic acid, 50.
 Parasitic baldness, 93.
 Parisian complexion balsam, 221.
 Pasta Mack, 224.
 Paste, almond, 141.
 Paste of barium sulphide, 121.
 Paste, Spanish, 168, 169.
 Pastes and meals, 90.
 Patchouli oil, 64.
 Patchouli oil, adulteration of, 65.
 Patchouli oil, properties of, 64.
 Patchouli oil, test for, 65.
 Pâte au musc for washing the hands, 111.
 Patent depilatory of G. Huettemann and J. Zrzawy, 122.

- Pauli's Azymol, 231.
 Pearl white, 45.
 Pearl white (liquid), 112.
 Penang benzoin, 81.
 Pencil, nerve, 159.
 Peppermint oil, 40, 60, 61, 64, 237.
 Peppermint oil, adulteration of, 61.
 Peppermint oil, American, 60, 64.
 Peppermint oil, English, 60, 64.
 Peppermint oil, German, 60.
 Peppermint (oil of), menthol in, 40.
 Peppermint oil, properties of, 60.
 Peppermint oil, testing purity of, 61, 237.
 Peppermint tincture, effect on teeth, 238.
 Perfume of nitrobenzol for fats and soaps, 31.
 Perfume, spray, 168.
 Perfumery distinguished from cosmetics, 1.
 Perfumes, alcoholic, official test for, 239.
 Perfumes for strong scents, 9.
 Perfumes obtained by absorption or enfleurage, 16.
 Perfumes obtained by distillation, 5, 12-15.
 Perfumes obtained by "enfleurage," 7.
 Perfumes obtained by extraction, 5, 7, 17.
 Perfumes obtained by maceration, 5, 6, 15.
 Perfumes obtained by pressure, 5.
 Permanganate de Potasse, Eau de, 104.
 Perspiration, remedy for smell of, 135.
 Peruvian balsam, 79, 185, 195.
 Peruvian balsam, adulteration of, 79.
 Peruvian balsam, properties of, 79.
 Peruvian balsam soap powder, 195.
 Petroleum ether, 17.
 Petroleum ether, extraction by, 7.
 Phoenix's hair dye, 211.
 Photographer's soap, 169.
 Pinaud's brilliantine, 214.
 Pinaud's Eau de Quinine, 213.
 Pinka's Roborantium, 217.
 Piperin, 36.
 Piperonal, 36.
 Pistachio powder, 110.
 Piver, Oleolisse Tonique de, 140.
 Plaister, author's antiseptic, 138.
 Plaister, salicylic acid, 136.
 Plaister, styptic, 136.
 Plaister, sublimate, 136.
 Plaisters, defects in Redoin's instructions for making, 137.
 Plaisters (paper) for cosmetic purposes, 136.
 Plumbiferous hair waters, 210.
 Pohlmann's cosmetic mouth wash, 230.
 Pohlmann's face powder, 222.
 Pollan's cosmetic caustic, 221.
 Polt & Gruber's Venus paste, 227.
 Polt's reseda curling pomade, 216.
 Polymeric propylenes, 50.
 Pomade, brilliant transparent, 128.
 Pomade, cantharides, 130.
 Pomade, China, 130.
 Pomade dye for beard and whiskers, 157.
 Pomade fixateur, à la rose, 130.
 Pomade hongroise, 131.
 Pomade for preventing fall of hair, Lestikow's, 157.
 Pomade, scurf, 129.
 Pomade (tube) for beard and whiskers, 158.
 Pomade, walnut, 151.
 Pomades and scented oils, 6.
 Pomades, cosmetic, 126-134.
 Pomades for hair, 94.
 Pomades or "huiles antiques," 15.
 Pomades, rancid, 95.
 Pomades, stick, 100, 174.
 Pomades, wax, 100.
 Poplar buds' smell, imitation of, 8.
 Popp's antherine mouth water, 230.
 Potassium chlorate, effect on teeth, 238.
 Potassium disulphide, 37.
 Potassium permanganate for rectifying alcohol, 22.

- Potassium permanganate water, 104.
- Potato starch, purified, 38.
- Poudre Antiasthmatique Clery, 218.
- Poudre de pistaches, 110.
- Poudre de riz, 107, 163.
- Poudre de riz à la rose, 108.
- Poudre de riz de Java, 219.
- Poudre de serail, 107.
- Poudres des Dames Rosée, 108.
- Powder, frangipani sachet, 141.
- Powder, nail, 231.
- Powder, rose, 108.
- Powder, shampoo, 155.
- Powders, coloured, 108.
- Powders for teeth, 96.
- Powders, soap, 163.
- Powders, toilet, 99.
- Powdered medicinal soaps, 195.
- Praeger's axeromaticum, 220.
- Preparation for softening water, 164.
- Preparation of acetic acid, 29.
- Preparation of acetic ether, 32.
- Preparation of amyl valerianate, 33.
- Preparation of barium sulphide, 121.
- Preparation of bay rum, 123.
- Preparation of benzoic acid, 24.
- Preparation of bismuth white, 45.
- Preparation of borax, 29.
- Preparation of butyric ether, 33.
- Preparation of calcium carbonate, 46.
- Preparation of calcium sulphide, 37, 120.
- Preparation of chloroform, 23.
- Preparation of cochineal, 51.
- Preparation of cold cream, 91.
- Preparation of cosmetics, ingredients used in, 4.
- Preparation of cumarin, 29.
- Preparation of emulsions, fats used in, 89.
- Preparation of face powders and rouges, 105, etc.
- Preparation of hymol, 44.
- Preparation of ionone, 42, 142.
- Preparation of liver of sulphur, 37.
- Preparation of nitrobenzol, 30.
- Preparation of nitrous ether, 34.
- Preparation of oil of cloves, 62.
- Preparation of pyrogallie acid, 37.
- Preparation of starch, 38.
- Preparation of tin oxide, 45.
- Preparation of valerianic ether, 33.
- Preparation of vanillin, 39.
- Preparations for darkening red hair, 120.
- Preparations for promoting growth of hair, 156.
- Preparations for removal of freckles and moles, 98.
- Preparations, secret cosmetic, 208, etc.
- Preservative, glycerin as a, 88, 89.
- Preservative, salicylic acid as a, 89.
- Presses (hydraulic) for obtaining scents, 11.
- Presses (screw), for obtaining scents, 11.
- Pressure to obtain perfumes, 5.
- Properties of almond oil, 67.
- Properties of ambergris, 47.
- Properties of benzoin, 81.
- Properties of bergamot oil, 54.
- Properties of castoreum, 47.
- Properties of cinnamon oil, 66.
- Properties of civet, 51.
- Properties of cochineal, 51.
- Properties of ionone, 43.
- Properties of juniper oil, 65.
- Properties of lanolin, 83.
- Properties of lard, 72.
- Properties of lemon oil, 58.
- Properties of menthol, 39.
- Properties of musk, 48.
- Properties of nitrous ether, 34.
- Properties of oil of bitter almonds, 55.
- Properties of oil of cloves, 62.
- Properties of oil of lavender, 59.
- Properties of paraffin, 78.
- Properties of patchouli oil, 64.
- Properties of peppermint oil, 60.
- Properties of Peruvian balsam, 79.
- Properties of pure camphor, 56.
- Properties of pyrogallie acid, 37.
- Properties of rosemary oil, 68, 69.

- Properties of safrol, 42.
 Properties of spermaceti, 71, 72.
 Properties of styrax, 79.
 Properties of vanillin, 39.
 Properties of vaseline, 81, 82.
 Properties of ylang-ylang oil, 65.
 Protective wash for hands, 135.
 Provence oil, 69.
 Pumice soap powder, 196.
 Pure camphor, properties of, 56.
 Pure cochineal, tests for, 52, 53.
 Purifying glycerin, 232-234.
 Purity of materials, testing, 232, etc.
 Purity of wax, test for, 71.
 Pyrogalllic acid, 37.
 Pyrogalllic acid, preparation of, 37.
 Pyrogalllic acid, properties of, 37.

 QUERLOIN'S Eau Cosmetique, 222.
 Quillaya bark tincture, Dieterich's recipe, 126.
 Quillaya bark tincture, preparation of, 122.
 Quillaya hair wash, 182.
 Quinine, 186.
 Quinine and glycerine hair tonic, compound, 154.
 Quinine hair tonic, 155.
 Quinine soap powder, 196.

 RANCID pomades, 95.
 Raspberry colouring matter, 53.
 Ratanhia extract, 53.
 Ratanhia root, 103, 147.
 Recipes for medicinal soaps, 194.
 Recovery of scents, 5.
 Rectification of alcohol by potassium permanganate, 22.
 Rectification of alcohol by sodium peroxide, 22.
 Rectification of benzine, 25.
 Red alkanna, 53.
 Red colour, elderberry, 53.
 Red currant berries for colouring matter, 53.
 Red from cochineal, 51, 52, 53.
 Red from red wine, 53.
 Red hair, preparations for darkening, 120.

 Red, madder, 53.
 Red, raspberry, 53.
 Red, rose, 53.
 Red, santal wood, 53.
 Redoin's recipes for plaisters, 137.
 Reds, 145, 146, 147.
 Refining, glycerin solution, 34.
 Regnault's liquid medicated soap, 197.
 Rejuvenating pills, 219.
 Remedies for moles, 134.
 Remedy for smell of perspiration, 135.
 Remedy for sweaty hands, 135.
 Remedies for warts, 134.
 Removal of disagreeable smells, 135.
 Removal of scurf, 92.
 Renard's Eau des Princesses, 222.
 Restorers for hair, 116, etc., 122, 123, 186, 187.
 Riedl's freckle ointment, 227.
 Ristoratore dei Capelli, 212.
 Rix's Pasta Pompadour, 224.
 Riz, poudre de, 107, 163.
 Robiquet's liquid soap, 197.
 Rose bandoline, 140.
 Rose cream, lanolin, 152.
 Rose face powder, 110.
 Rose oil, 64.
 Rose powder, 108, 110, 163.
 Rosemary hair wash, 182.
 Rosemary oil, 64, 68.
 Rosemary oil, properties of, 68, 69.
 Roses, milk of, 141.
 Rostel's Recreation, 216.
 "Rosy cheeks without rouge," 225.
 Rothe's tooth wash, 228.
 Rother's Pasta Cosmetica, 224.
 Rotondi's experiments on soaps, 188, etc.
 Rottmanner's beauty water, 227.
 Rottmanner's cosmetic paste, 221.
 Rouge, 98.
 Rouge de Théâtre, 225.
 Rouge en feuilles, 113, 147.
 Rouge en pâte, 113.
 Rouge en tasses, 147.
 Rouge, liquid vegetable, 112.

- Rouge paste, dry, 107.
 Rouges, preparation of, 105, etc.
 Rough skin and burns, ointment for, 114.
 Rowland's Kalydor, 223.
 Rowland's Kalydor cosmetic wash, 221.
 Royer's beard-producing pomade, 214.
 Russ's freckle wash, 227.

 SACH's mouth wash, 228.
 Sachet powder frangipani, 141.
 Saffron, 146.
 Safrol, 42.
 Safrol, preparation of, 42.
 Safrol, properties of, 42.
 Sage water, 105.
 Saidemann's liquid soap, 197.
 Salicylated tallow, 173.
 Salicylic acid as a preservative, 89.
 Salicylic acid as a retarder of rancidity, 95.
 Salicylic acid face powder, 99.
 Salicylic acid plaister, 136.
 Salicylic acid powder, 111.
 Salicylic acid soap powder, 195.
 Salicylic powder, 172.
 Salicylic - resorcin soap powder, 195.
 Salicylic - resorcin - sulphur soap powder, 172.
 Salicylic-sulphur soap powder, 195.
 Salol soap powder, 196.
 Salol tooth powder, (Huchard's), 102.
 Salve for lips (excellent), 129.
 Salvia, Eau de, 105.
 Sand-almond bran, 176.
 Santal wood colouring matter, 53.
 Santal wood oil, 64.
 Sapo viridis, alcoholic solution of, 94.
 Saponification value in adulterated lard, 77.
 Sarah Bernhardt Powder la Diaphane, 225.
 Scent cakes, 170.
 Scent essences, 124.
 Scented oils and pomades, 6.

 Scented paraffin tablets, 177.
 Scented starch, 107.
 Scents and aromatics, volatility of, 4.
 Scents for soaps, 125.
 Scents obtained by pressure, 5, 11, 12.
 Scents, recovery of, 5.
 Schimmel & Co.'s almond oil soap, 202.
 Schleich's marble soap, 199.
 Schneeweiss's rapid hair dye, 211.
 Schneider's soaps, 192.
 Schubert's milk of lilies, 224.
 Screw presses for obtaining scents, 11.
 Scurf essence (Lautenschlaeger's), 211.
 Scurf pomade, 129.
 Scurf, removal of, 92.
 Scurf soap, 157.
 Scurf water, antiseptic, 187.
 Secret cosmetic preparations, 208, etc.
 Secret preparations for the hair, 209.
 Secret preparations for skin and complexion, 218-228.
 Secret preparations for teeth, mouth, etc., 228-231.
 Seebald's hair tincture, 209, 218.
 Seebald's tincture for preventing scurf and falling out of hair, 213.
 Seegar's blonde hair dye, 209.
 Seegar's hair water, 210.
 Seiffert's Schneewittchen, 225.
 Separation of expressed liquid, 12.
 Sérail, poudre de, 107.
 Sesser's medicinal toilet mystery, 227.
 Shampoo powder, 155, 217.
 Shampoos, 154.
 Shaving creams, 164.
 Shaving soaps, 203.
 Shaving soaps by cold process, 203.
 Siam benzoin, 80.
 Siberian castoreum, 47.
 Silver nitrate, effect on teeth, 238.
 Silver nitrate stains, removal of, 169.

- Simmerling's Cosmétique, 221.
 Singeing the hair, 92.
 Skin and complexion, secret preparations for, 218-228.
 Skin beautifiers, 98.
 Skin, care of, 86, 87.
 Skin gloss, 111.
 Skin, hydrogen peroxide wash for, 114.
 Smalt, 147.
 Smelling salts, English, 168.
 Smelling salts, French, 168.
 Smokers, mouth wash for, 96.
 Snuff (menthol) for catarrh, 176.
 Soap essences, 124, 167, 196, etc.
 Soap for washing hair, 94.
 Soap hair washes, 182, 183.
 Soap, hygienic, 166.
 Soap, marble-dust, 166.
 Soaps and washes, antiseptic, 179, etc.
 Soap, medicinal sulphur-sand, 198.
 Soap, nicotine, 166.
 Soap, photographers', 169.
 Soap powder, testing purity of, 235.
 Soap powders, 163.
 Soap powders, recipes for, 195.
 Soap scents, 125.
 Soap, scurf, 157.
 Soap spirit, 205.
 Soaps, antiseptic, preparation of, 187, etc.
 Soaps, cosmetic toilet, 200, etc.
 Soaps, etc., containing nitrobenzol, 31.
 Soaps, hygienic, 200, etc.
 Soaps, liquid, 196, etc.
 Soaps (medicinal) cocoanut oil for, 193.
 Soaps, medicinal sulphur and tar, 198.
 Soaps, milled, 201.
 Soaps, perfumed with safrol, 42.
 Soaps, recipes for medicinal, 194.
 Soaps, Rotondi's experiments on, 188, etc.
 Soaps, Schneider's, 192.
 Soaps, shaving, 203.
 Soaps, Unna's medicinal, 191.
 Sodium peroxide for rectifying alcohol, 22.
 Soft soap, sulphur, 198.
 Softening water, preparation for, 164.
 Solid rouge, 113.
 Solid soap spirit, 205.
 Solubility of acetic ether, 33.
 Solubility of ammonium carbonate, 26.
 Solubility of bismuth white, 45.
 Solubility of borax, 29.
 Solubility of calcium carbonate, 46.
 Solubility of carmine, 52.
 Solubility of cumarin, 29.
 Solubility of heliotropin, 36.
 Solubility of menthol, 39.
 Solubility of nitrobenzol, 31.
 Solubility of nitrous ether, 34.
 Solubility of paraffin, 78.
 Solubility of thymol, 44.
 Soluble starch as prepared by Wroblewski, 38.
 Solvent power of alcohol, 20.
 Sozodont, 176.
 Spanish leather, 168.
 Spanish paste, 168, 169.
 Specialties for the teeth, 170.
 Specialties in hair washes, 150.
 Spent cochineal, 51.
 Spermaceti, 71.
 Spermaceti, adulteration of, 72.
 Spermaceti, properties of, 71, 72.
 Spermaceti, test for purity of, 72.
 Spirit, soap, 205.
 Spitzer's face pomade, 218.
 Spitzer's freckle ointment, 220.
 Spray perfume, 168.
 Spring of Life, 214.
 Stains caused by silver nitrate, removal of, 169.
 Starch, 38.
 Starch, coloured, 108.
 Starch for toilet purposes, 107.
 Starch, preparation of, 38.
 Starch, purified potato, 38.
 Starch, scented, 107.
 Starch, soluble, 38.
 Stenard's Eau de Beauté, 222.
 Sterling's ambrosia, 213.

- Stick pomades, 100, 131, 174.
 Sticking plaister, antiseptic, 137.
 Sticking plaister (antiseptic), author's recipe for, 138.
 Stiffening hair, brilliantine for, 158.
 St. John's wort and birch leaf ointment, 114.
 Still for distillation, 13, 14.
 Strawberry scent, imitation of, 8.
 Strong scents, perfumes for, 9.
 Styptic plaister, 136.
 Styrax, 79.
 Styrax, properties of, 79.
 Sublimate plaister, 136.
 Sublimate soap, 194.
 Substitute for diachylon plaister, 137.
 Sulphur and tar medicinal soaps, 198.
 Sulphur in ether, test for, 20.
 Sulphur, liver of, 37.
 Sulphur sand medicinal soap, 198.
 Sulphur soap, 194.
 Sulphur soap, liquid, 199.
 Sulphur soap powder, 195.
 Sulphur soft soap, 198.
 Sulphuric acid and ether, mixing of, 19.
 Sulphuric acid in acetic acid, detection of, 30.
 Sulphuric ether, 19.
 Sulphuric ether, extraction by, 7.
 Sulphuric ether test for almond oil, 68.
 Sumatra almond benzoin, 80.
 Sweaty hands, preparations for, 132-134.
 Sweaty hands, remedy for, 135.
 TABLE of specific gravity of alcohol, 22.
 Tablets, mouth wash, 159.
 Tallow, salicylated, 173.
 Tannin, 186.
 Tannin hair dyes, 117.
 Tanno-quinine hair restorer, 186.
 Tecklenburg's milk of Venus, 227.
 Teeth, care of, 95.
 Teeth, effect of various substances on, 238.
 Teeth, secret preparations for, 228-231.
 Teeth, specialties for, 170.
 Terpinol, 9.
 Tertiary butyltolylmethane, 50.
 Test for carmine, 52.
 Test for patchouli oil, 65.
 Test for pure lard, 72.
 Test for purity of bergamot oil, 55.
 Test for purity of lanolin, 84.
 Test for purity of spermaceti, 72.
 Test for purity of vaseline, 82.
 Test for purity of wax, 71.
 Test for thymol, 44, 45.
 Testing materials in cosmetics, 232, etc.
 Tests for adulterants in oil of cloves, 62.
 Tests for benzine, 24, 25.
 Tests for ethereal oils, 63.
 Tests for oil of lavender, 60.
 Tests for pure cochineal, 52, 53.
 Tests for pure peppermint oil, 61.
 Tests for purity of almond oil, 67.
 Theatrical make-up, vein blue for, 114.
 Thiosavonal, 198, etc.
 Thyme oil, 64.
 Thymol, 44.
 Thymol, alcoholic solution of, 96.
 Thymol, preparation of, 44.
 Thymol soap powder, 196.
 Thymol, solubility of, 44.
 Thymol solution for mouth wash (Hermites), 102.
 Thymol, test for, 44, 45.
 Tincture de Leytens, 217.
 Tincture, laurel, 140.
 Tincture, musk, 144.
 Tincture of Quillaya bark, preparation of, 122.
 Tincture, violet root, 206.
 Tin oxide, 45.
 Tin oxide, preparation of, 45.
 Tobacco juice, effect on teeth, 238.
 Toellner's glycerin hair restorer, 123.
 Toilet (cosmetic) soaps, 200, etc.
 Toilet cream, lanolin, 126.
 Toilet powders, 99.

- Toluol, 50.
 Tonique de Piver, Oleolisse, 140.
 Tooth pastes, 101, etc.
 Tooth powder of levigated chalk, 97.
 Tooth powders, 96.
 Tooth powders, errors in choice of, 96.
 Toothache drops, menthol, 175.
 Transparent brilliant pomade, 128.
 Trixogene hair restorer, 211.
 Tube pomade for beard and whiskers, 158.
 Turkish water, 153.
 Turmeric, 146.
 Turpentine as an adulterant for lemon oil, 58.
 Turpentine product as substitute for musk, 49.
 Two groups of emulsions, 88.

 ULRICH's mouth wash, 228.
 Ultramarine, 147.
 Unguentum paraffini, 207.
 Unna's medicinal soaps, 191.
 Use of cosmetics, 1.

 VALERIANIC ether, 33.
 Valerianic ether, preparation of, 33.
 Van Buskirk's Tragacanth-Sozodont, 228.
 Vanilla essence, artificial, 143.
 Vanilla soap, scent for, 125.
 Vanillin, 9, 39.
 Vanillin, preparation of, 39.
 Vanillin, properties of, 39.
 Vaseline, 81.
 Vaseline, adulteration of, 82.
 Vaseline, consistence of, 81.
 Vaseline ice, camphorated, 128.
 Vaseline injected for restoring plumpness, 206, 207.
 Vaseline, properties of, 81, 82.
 Vaseline, test for purity of, 82.
 Vaseline, testing purity of, 237.
 Vaseline, 83.
 Vegetable blue for veins, 113.
 Vegetable milk, 90.
 Vegetable oils in lard, Becchi and Welmann's tests for, 75.
 Vegetable rouge (liquid), 112.
 Vein blue for theatrical make-up, 114.
 Veins, cosmetic for, 160.
 Veins, vegetable blue for, 113.
 Vienna cosmetic, 210.
 Vienna face water, 219.
 Viennese face wash, 225.
 Vincent's poudre rafraichissante, 228.
 Vinegar, menthol, 175.
 Viola's ephelidène, 222.
 Violet colouring matter, 145, 148.
 Violet powder, 111, 162.
 Violet root, essence of, 142.
 Violet root powder in toilet soaps, 206.
 Violet root tincture, 206.
 Violet water, 105.
 Violets, artificial essence of, 143.
 Violets, artificial oil of, 43.
 Vogel's odalin, 224.
 Volatility of scents and aromatics, 4.

 WALLRITZ's cactus pomade, 214.
 Walnut hair oil, 151.
 Walnut pomade, 151.
 Walnut shells as hair dye, 151.
 Warts, 221.
 Warts, preparations for, 132-134.
 Warts, remedies for, 134.
 Wash for removal of disagreeable smells, 135.
 Washes and soaps, antiseptic, 179, etc.
 Washes for curling hair, recipes for, 155.
 Washes for freckles, 132, 133, 134.
 Washes for hair, American, 154.
 Washes for hair (antiseptic), recipes for, 184, 185.
 Washes for hair (medicinal), 184.
 Washes for hair, recipes for, 182, etc.
 Washing the hair, 92.
 Washing creams, 166.
 Water softening preparation, 164.
 Wax, 71.
 Wax, adulteration of, 71.
 Wax for moustache and beard, 132.

- Wax pomades, 100.
Whiskers, specialties for, 157.
White, bismuth, 45.
White face powder, 110.
White, French, 112.
White, pearl, 45.
White, liquid pearl, 112.
White Peruvian balsam, 79.
White powder, 163.
"White rose" perfume, imitation of, 8.
White wax paper for wrapping, preparation of, 238.
Windsor soap, scent for, 125.
Wintergreen, oil of, 40, 64.
Wortley's berenizon, 214.
YELLOW hair dye, 119, 120.
Yellows, 146.
Ylang-ylang oil, 64, 65.
Ylang-ylang oil, properties of, 65.
Yolk of egg as a cosmetic, 127, 128.
ZADIG's hygienic soap, 204.

Catalogue

OF

SCOTT, GREENWOOD & CO.'S

Special Technical Works

FOR

MANUFACTURERS, PROFESSIONAL MEN, STUDENTS,
COLLEGES AND TECHNICAL SCHOOLS
BY EXPERT WRITERS

INDEX TO SUBJECTS.

PAGE	PAGE	PAGE
Agricultural Chemistry ... 4	Dyeing Woollen Fabrics ... 28	Paper-pulp Dyeing... ... 21
Air, Industrial Use of ... 10	Dyers' Materials ... 25	Petroleum 8
Alum ... 4	Enamelling Metal ... 21, 22, 32	Pigments, Chemistry of ... 12
Alumina Sulphates ... 4	Enamels ... 21	Plumbers' Work ... 30, 31
Ammonia ... 5	Essential Oils ... 7	Porcelain Painting... ... 19
Aniline Colours ... 6	Evaporating Apparatus ... 14	Pottery Clays ... 19
Animal Fats ... 2	External Plumbing... ... 30	Pottery Manufacture ... 16, 18
Anti-corrosive Paints ... 3	Fats ... 2, 3	Pottery Recipes ... 16
Architectural Pottery ... 17	Gas Firing ... 30	Power-loom Weaving ... 23
Artificial Perfumes... ... 7	Glass-making Recipes ... 16	Preserved Foods ... 34
Balsams ... 11	Glass Painting ... 19	Recipes for Oilmen, etc. ... 10
Bleaching ... 26	Glue Making... ... 11	Resins... ... 11
Bone Products ... 11	Glue Testing... ... 8	Risks of Occupations ... 13
Brick-making ... 31	Greases ... 2, 4	Rivetting China, etc. ... 19
Burnishing Brass ... 31	History of Staffs Potteries ... 20	Röntgen Ray Work ... 13
Carpet Yarn Printing ... 17	Hops ... 33	Scheele's Essays ... 8
Ceramic Books ... 16-20	Hot-water Supply ... 32	Silk Dyeing ... 27
Charcoal ... 11	India-Rubber ... 13	Smoke Prevention ... 30
Chemical Essays ... 8	Inks ... 7	Soaps ... 2
Chemistry of Pottery ... 20	Iron-corrosion ... 3	Spinning ... 28
Clay Analysis ... 17	Iron, Science of ... 36	Staining Marble, Bone, &c. ... 35
Coal-dust Firing ... 30	Japanning ... 32	Steam Drying ... 14
Colour Matching ... 27	Lacquering ... 31	Sweetmeats ... 34
Colliery Recovery Work ... 29	Lake Pigments ... 6, 10	Terra-cotta ... 17
Colour-mixing for Dyers ... 27	Lead and its Compounds... 12	Testing Paint Materials ... 6
Colouring Pottery ... 17	Lead Ores ... 13	Testing Yarns ... 22
Colour Theory ... 24	Leather Industry ... 15	Textile Fabrics ... 22-29
Combing Machines... ... 29	Leather-working Materials ... 14	Textile Materials ... 24
Compounding Oils ... 2	Lubricants ... 2, 4	Timber ... 34
Condensing Apparatus ... 14	Lubricating Oils ... 4	Varnishes ... 5
Cosmetics ... 7	Manures ... 4, 11	Vegetable Fats ... 3
Cotton Dyeing ... 28	Mineral Pigments ... 10	Waste Utilisation ... 5
Cotton Spinning ... 28, 29	Oil and Colour Recipes ... 10	Water, Industrial Use of ... 10
Dampness in Buildings ... 35	Oil Boiling ... 5	Wood ... 33, 34
Decorating Pottery ... 17	Oil Refining ... 5	Wood Waste Utilisation ... 33
Decorators' Books ... 32	Oils ... 2, 4, 7	Wood Dyeing ... 35
Decorative Textiles ... 22	Ozone, Industrial Use of ... 10	Wool Dyeing ... 27
Dictionary of Paint Materials ... 9	Painters' Books ... 32	X-Ray Work ... 13
Drying Oils ... 7	Painting Glass ... 19	Yarn Numbering ... 23
Drying with Air ... 14	Paint Manufacture... ... 12	Yarn Testing ... 22
Dyeing Marble ... 35	Paint Materials ... 6, 9	
	Paint-material Testing ... 6	

SCOTT, GREENWOOD & CO.,

19 LUDGATE HILL, LONDON, E.C.

Tel. Address: "PRINTERIES, LONDON".

Tel. No. 5403, Bank.

Books on Oils, Soaps, Colours, Chemicals, Glue, Varnishes, etc.

THE PRACTICAL COMPOUNDING OF OILS, TALLOW AND GREASE FOR LUBRICATION, ETC.

By AN EXPERT OIL REFINER. 100 pp. 1898. Demy 8vo. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., **Introductory Remarks** on the General Nomenclature of Oils, Tallow and Greases suitable for Lubrication.—II., **Hyrocarbon Oils**.—III., **Animal and Fish Oils**.—IV., **Compound Oils**.—V., **Vegetable Oils**.—VI., **Lamp Oils**.—VII., **Engine Tallow, Solidified Oils and Petroleum Jelly**.—VIII., **Machinery Greases: Loco and Anti-friction**.—IX., **Clarifying and Utilisation of Waste Fats, Oils, Tank Bottoms, Drainings of Barrels and Drums, Pickings Up, Dregs, etc.**—X., **The Fixing and Cleaning of Oil Tanks, etc.**—Appendix and General Information.

Press Opinions.

"This work is written from the standpoint of the oil trade, but its perusal will be found very useful by users of machinery and all who have to do with lubricants in any way."—*Colliery Guardian*.

"The properties of the different grades of mineral oil and of the animal and vegetable non-drying oils are carefully described, and the author justly insists that the peculiarities of the machinery on which the lubricants are to be employed must be considered almost before everything else. . . . The chapters on grease and solidified oils, etc., are excellent."—*The Ironmonger*.

SOAPS. A Practical Manual of the Manufacture of Domestic, Toilet and other Soaps. By GEORGE H. HURST, F.C.S. Illustrated with Sixty-six Engravings. 390 pp. 1898. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

Chapters I., **Introductory**.—II., **Soap-maker's Alkalies**.—III., **Soap Fats and Oils**.—IV., **Perfumes**.—V., **Water as a Soap Material**.—VI., **Soap Machinery**.—VII., **Technology of Soap-making**.—VIII., **Glycerine in Soap Lyes**.—IX., **Laying out a Soap Factory**.—X., **Soap Analysis**.—Appendices.

Press Opinions.

"We think it is the most practical book on these subjects that has come to us from England so far."—*American Soap Journal*.

"Much useful information is conveyed in a convenient and trustworthy manner which will appeal to practical soap-makers."—*Chemical Trade Journal*.

"Works that deal with manufacturing processes, and applied chemistry in particular, are always welcome. Especially is this the case when the material presented is so up-to-date as we find it here."—*Bradford Observer*.

ANIMAL FATS AND OILS: Their Practical Production, Purification and Uses for a great Variety of Purposes. Their Properties, Falsification and Examination. A Handbook for Manufacturers of Oil and Fat Products, Soap and Candle Makers, Agriculturists, Tanners, Margarine Manufacturers, etc., etc. By LOUIS EDGAR ANDÉS. Sixty-two Illustrations. 240 pp. 1898. Demy 8vo. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Introduction. Occurrence, Origin, Properties and Chemical Constitution of Animal Fats. Preparation of Animal Fats and Oils. Machinery. Tallow-melting Plant. Extraction Plant. Presses. Filtering Apparatus. Butter: Raw Material and Preparation, Properties, Adulterations, Beef Lard or Remelted Butter, Testing. Candle-fish Oil. Mutton-Tallow. Hare Fat. Goose Fat. Neatsfoot Oil. Bone Fat: Bone Boiling, Steaming Bones, Extraction, Refining. Bone Oil. Artificial Butter: Oleomargarine, Margarine Manufacture in France, Grasso's Process, "Kaiser's Butter," Jahr & Munzberg's Method, Filbert's Process, Winter's Method. Human Fat. Horse Fat. Beef Marrow. Turtle Oil. Hog's Lard: Raw Material Preparation, Properties, Adulterations, Examination. Lard Oil. Fish Oils. Liver Oils. Artificial Train Oil. Wool Fat: Properties, Purified Wool Fat. Spermaceti: Examination of Fats and Oils in General.

Press Opinions.

"The descriptions of technical processes are clear, and the book is well illustrated and should prove useful."—*Manchester Guardian*.

"It is a valuable work, not only for the student, but also for the practical manufacturer of oil and fat products."—*Transactions of the American Chemical Society*.

"The work is very fully illustrated, and the style throughout is in strong contrast to that employed in many such treatises, being simple and clear."—*Shropshire Leader*.

"An important handbook for the 'fat and oil' industry, now a large one."—*Newcastle Chronicle*.

"The latest and most improved forms of machinery are in all cases indicated, and the many advances which have been made during the past years in the methods of producing the more common animal fats—lard, tallow and butter—receive due attention."—*Glasgow Herald*.

VEGETABLE FATS AND OILS: Their Practical Preparation, Purification and Employment for Various Purposes, their Properties, Adulteration and Examination. A Handbook for Oil Manufacturers and Refiners, Candle, Soap and Lubricating Oil Makers, and the Oil and Fat Industry in General. Translated from the German of LOUIS EDGAR ANDÉS. Ninety-four Illustrations. 340 pp. Second Edition. 1902. Demy 8vo. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Statistical Data. General Properties of the Vegetable Fats and Oils. Estimation of the Amount of Oil in Seeds. Taste of Vegetable Fats and Oils, with French and German Nomenclature, Source and Origin and Percentage of Fat in the Plants from which they are Derived. The Preparation of Vegetable Fats and Oils: Storing Oil Seeds: Cleaning the Seed. Apparatus for Grinding Oil Seeds and Fruits. Installation of Oil and Fat Works. Extraction Method of Obtaining Oils and Fats. Oil Extraction Installations. Press Moulds. Non-drying Vegetable Oils. Vegetable-lyring Oils. Solid Vegetable Fats. Fruits Yielding Oils and Fats. Woolsortening Oils. Soluble Oils. Treatment of the Oil after Leaving the Press. Improved Methods of Refining with Sulphuric Acid and Zinc Oxide or Lead Oxide. Refining with Caustic Alkalies, Ammonia, Carbonates of the Alkalies, Lime. Bleaching Fats and Oils. Practical Experiments on the Treatment of Oils with regard to Refining and Bleaching. Testing Oils and Fats.

Press Opinions.

"Concerning that and all else within the wide and comprehensive connection involved, this book must be available to every one directly or indirectly interested in the matters it treats of."—*Commerce*.

"A valuable and highly interesting book of reference."—*Manufacturing Chemist*.

IRON - CORROSION. ANTI-FOULING AND ANTI-CORROSIVE PAINTS. By LOUIS EDGAR ANDÉS. Sixty-two Illustrations. 275 pp. Translated from the German. Demy 8vo. 1900. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Iron-rust and its Formation—Protection from Rusting by Paint—Grounding the Iron with Linseed Oil, etc.—Testing Paints—Use of Tar for Punting on Iron—Anti-corrosive Paints—Linseed Varnish—Chinese Wood Oil—Lead Pigments—Iron Pigments—Artificial Iron Oxides—Carbon—Preparation of Anti-corrosive Paints—Results of Examination of Several Anti-corrosive Paints—Paints for Ship's Bottoms—Anti-fouling Compositions—Various Anti-corrosive and Ship's Paints—Official Standard Specifications for Ironwork Paints—Index.

Press Opinions.

"This is a very valuable book."—*Bristol Mercury*.

"Will be of great service to paint manufacturers, engineering contractors, ironfounders, shipbuilders and others."—*Engineer and Iron Trades Advertiser*.

"Deals with the subject in a manner at once practical and scientific, and is well worthy of the attention of all builders, architects and engineers."—*The Builder*.

"The book is very readable and full of valuable information."—*Railway Engineer*.

"The author goes fully into his subject."—*Journal of Gas Lighting*.

"The book is an exceedingly useful record of what has been done in connection with iron preservation, and will undoubtedly prove to be of much value to railway engineers, shipowners, etc."—*Fairplay*.

"Will be particularly useful to iron manufacturers, shipbuilders and shipowners. . . ."—*Ironmonger*.

"... The book is a welcome contribution to technological literature."—*Western Daily Mercury*.

"The author has evidently thoroughly investigated and mastered the subject of iron-corrosion, its cause and its prevention."—*Iron and Steel Trades Journal*.

THE MANUFACTURE OF ALUM AND THE SULPHATES AND OTHER SALTS OF ALUMINA AND IRON. Their Uses and Applications as Mordants in Dyeing and Calico Printing, and their other Applications in the Arts, Manufactures, Sanitary Engineering, Agriculture and Horticulture. Translated from the French of LUCIEN GESCHWIND. 195 Illustrations. Nearly 400 pp. Royal 8vo. 1901. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

Part I., **Theoretical Study of Aluminium, Iron, and Compounds of these Metals.**—Chapters I., Aluminium and its Compounds.—II., Iron and Iron Compounds.
 Part II., **Manufacture of Aluminium Sulphates and Sulphates of Iron.**—Chapters III., Manufacture of Aluminium Sulphate and the Alums.—IV., Manufacture of Sulphates of Iron.
 Part III., **Uses of the Sulphates of Aluminium and Iron.**—Chapters V., Uses of Aluminium Sulphate and Alums—Application to Wool and Silks—Preparing and using Aluminium Acetates—Employment of Aluminium Sulphate in Carbonising Wool—The Manufacture of Lake Pigments—Manufacture of Prussian Blue—Hide and Leather Industry—Paper Making—Hardening Plaster—Lime Washes—Preparation of Non-inflammable Wood, etc.—Purification of Waste Waters.—VI., **Uses and Applications of Ferrous Sulphate and Ferric Sulphates.**—Dyeing—Manufacture of Pigments—Writing Inks—Purification of Lighting Gas—Agriculture—Cotton Dyeing—Disinfectant—Purifying Waste Liquors—Manufacture of Nordhausen Sulphuric Acid—Fertilising.
 Part IV., **Chemical Characteristics of Iron and Aluminium.**—Analysis of Various Aluminous or Ferruginous Products.—Chapter VII., Aluminium.—Analysing Aluminium Products.—Alumina—Alumina—Sodium Aluminate—Aluminium Sulphate. Chapter VIII., **Iron.**—Analytical Characteristics of Iron Salts—Analysis of Pyritic Lignite—Ferrous and Ferric Sulphates—Ror²¹ Mordant—Index.

MANUAL OF AGRICULTURAL CHEMISTRY. By HERBERT INGLE, F.I.C., Lecturer on Agricultural Chemistry, the Yorkshire College; Lecturer in the Victoria University. 388 pp. 11 Illustrations. 1902. Demy 8vo. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d. net.

Contents.

Chapters I., Introduction.—II., The Atmosphere.—III., The Soil.—IV., The Reactions occurring in Soils.—V., The Analysis of Soils.—VI., Manures, Natural.—VII., Manures (continued).—VIII., The Analysis of Manures.—IX., The Constituents of Plants.—X., The Plant.—XI., Crops.—XII., The Animal.—XIII., Foods and Feeding.—XIV., Milk and Milk Products.—XV., The Analysis of Milk and Milk Products.—XVI., Miscellaneous Products used in Agriculture.—Appendix.—Index.

LUBRICATING OILS, FATS AND GREASES: Their Origin, Preparation, Properties, Uses and Analyses. A Handbook for Oil Manufacturers, Refiners and Merchants, and the Oil and Fat Industry in General. By GEORGE H. HURST, F.C.S. Second Edition. Sixty-five Illustrations. 313 pp. Demy 8vo. 1902. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Chapters I., **Introductory.** Oils and Fats, Fatty Oils and Fats, Hydrocarbon Oils, Uses of Oils.—II., **Hydrocarbon Oils.** Distillation, Simple Distillation, Destructive Distillation, Products of Distillation, Hydrocarbons, Paraffins, Olefins, Naphthenes.—III., **Scotch Shale Oils.** Scotch Shales, Distillation of Scotch Oils, Shale Retorts, Products of Distilling Shales, Separating Products, Treating Crude Shale Oil, Refining Shale Oil, Shale Oil Stills, Shale Naphtha Burning Oils, Lubricating Oils, Wax.—IV., **Petroleum.** Occurrence, Geology, Origin, Composition, Extraction, Refining, Petroleum Stills, Petroleum Products, Cylinder Oils, Russian Petroleum, Deblowing Mineral Oils.—V., **Vegetable and Animal Oils.** Introduction, Chemical Composition of Oils and Fats, Fatty Acids, Glycerine, Extraction of Animal and Vegetable Fats and Oils, Animal Oils, Vegetable Oils, Rendering, Pressing, Refining, Bleaching, Tallow, Tallow Oil, Lard Oil, Neatsfoot Oil, Palm Oil, Palm Nut Oil, Coconut Oil, Castor Oil, Olive Oil, Rape and Colza Oils, Arachis Oil, Niger Seed Oil, Spermin Oils, Whale Oil, Seal Oil, Brown Oils, Lardine, Thickened Rape Oil.—VI., **Testing and Adulteration of Oils.** Specific Gravity, Alkali Tests, Sulphuric Acid Tests, Free Acids in Oils, Viscosity Tests, Flash and Fire Tests, Evaporation Tests, Iodine and Bromide Tests, Elaidin Test, Melting Point of Fat, Testing Machines.—VII., **Lubricating Greases.** Rosin Oil, Anthracene Oil, Making Greases, Testing and Analysis of Greases.—VIII., **Lubrication.** Friction and Lubrication, Lubricant, Lubrication of Ordinary Machinery, Spontaneous Combustion of Oils, Stainless Oils, Lubrication of Engine Cylinders, Cylinder Oils.—**Appendices.** A. Table of Baume's Hydrometer.—B. Table of Thermometric Degrees.—C. Table of Specific Gravities of Oils.—Index.

THE UTILISATION OF WASTE PRODUCTS. A Treatise on the Rational Utilisation, Recovery and Treatment of Waste Products of all kinds. By Dr. THEODOR KOLLER. Translated from the Second Revised German Edition. Twenty-two Illustrations. Demy 8vo. 280 pp. 1902. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Introduction. - Chapters I., The Waste of Towns. - II., Ammonia and Sal-Ammoniac - Rational Processes for Obtaining these Substances by Treating Residues and Waste. - III., Residues in the Manufacture of Aniline Dyes. - IV., Amber Waste. - V., Brewers' Waste. - VI., Blood and Slaughter-House Refuse. - VII., Manufactured Fuels. - VIII., Waste Paper and Bookbinders' Waste. - IX., Iron Slags. - X., Excrement. - XI., Colouring Matters from Waste. - XII., Dyers' Waste Waters. - XIII., Fat from Waste. - XIV., Fish Waste. - XV., Cud-mine Sludge. - XVI., Tannery Waste. - XVII., Gold and Silver Waste. - XVIII., India-rubber and Camelliole Waste. - XIX., Residues in the Manufacture of Rosin Oil. - XX., Wood Waste. - XXI., Horn Waste. - XXII., Infusorial Earth. - XXIII., Iridium from Goldsmith's Sweepings. - XXIV., Jute Waste. - XXV., Cork Waste. - XXVI., Leather Waste. - XXVII., Glue Makers' Waste. - XXVIII., Illuminating Gas from Waste and the By-Products of the Manufacture of Coal Gas. - XXIX., Meersenum. - XXX., Molasses. - XXXI., Metal Waste. - XXXII., By-Products in the Manufacture of Mineral Waters. - XXXIII., Fruit. - XXXIV., The By-Products of Paper and Paper Pulp Works. - XXXV., By-Products in the Treatment of Coal Tar Oils. - XXXVI., Fur Waste. - XXXVII., The Waste Matter in the Manufacture of Parchment Paper. - XXXVIII., Mother of Pearl Waste. - XXXIX., Petroleum Residues. - XL., Platinum Residues. - XLI., Broken Porcelain, Earthenware and Glass. - XLII., Salt Waste. - XLIII., Slate Waste. - XLIV., Sulphur. - XLV., Burnt Pyrites. - XLVI., Silk Waste. - XLVII., Soap Makers' Waste. - XLVIII., Alkali Waste and the Recovery of Soda. - XLIX., Waste Produced in Grinding Mirrors. - L., Waste Products in the Manufacture of Starch. - LI., Stearic Acid. - LII., Vegetable Ivory Waste. - LIII., Turf. - LIV., Waste Waters of Cloth Factories. - LV., Wine Residues. - LVI., Tinsplate Waste. - LVII., Wool Waste. - LVIII., Wool Sweat. - LIX., The Waste Liquids from Sugar Works. - Index.

AMMONIA AND ITS COMPOUNDS: Their Manufacture and Uses. By CAMILLE VINCENT, Professor at the Central School of Arts and Manufactures, Paris. Translated from the French by M. J. SALTER. Royal 8vo. 113 pp. 1901. Thirty-two Illustrations. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Chapters I., General Considerations: Sections 1. Various Sources of Ammoniacal Products; 2. Human Urine as a Source of Ammonia. II., Extraction of Ammoniacal Products from Sewage: Sections 1. Preliminary Treatment of Excreta in the Settling Tanks - The Lencauchez Process, The Bilange Process, The Kuentz Process; 2. Treatment of the Clarified Liquors for the Manufacture of Ammonium Sulphate - The Figuera Process and Apparatus, Apparatus of Margueritte and Sourdeval, The Lair Apparatus, Apparatus of Sintier and Muhé, Apparatus of Bilange, The Kuentz Process, Process and Apparatus of Hennebutte and De Vauréal; 3. Treatment of Entire Sewage - Chevalet's Apparatus, Paul Mallet's Apparatus, Lencauchez's Apparatus. III., Extraction of Ammonia from Gas Liquor: Sections 1. Clarification of Gas Liquor; 2. Manufacture of Ammonium Sulphate - A. Mallet's Apparatus, A. Mallet's Modified Apparatus, Paul Mallet's Apparatus, Chevalet's Apparatus, Grüneberg's Apparatus; 3. Concentration of Gas Liquor - Solvay's Apparatus, Kuentz's Apparatus, Grüneberg's Apparatus. IV., Manufacture of Ammoniacal Compounds from Bones, Nitrogenous Waste, Beetroot Wash and Peat: Sections 1. Ammonia from Bones; 2. Ammonia from Nitrogenous Waste Materials; 3. Ammonia from Beetroot Wash (Vinassee); 4. Ammonia from Peat - Treatment of the Ammoniacal Liquors. V., Manufacture of Caustic Ammonia, and Ammonium Chloride, Phosphate and Carbonate: Sections 1. Manufacture of Caustic Ammonia; 2. Manufacture of Ammonium Chloride - From Fermented Urine, Process of the Lesage Company, Kuentz's Process; From Gas Liquor, English Process, Kuentz's Process; From the Dry Distillation of Animal Matter; From Ammonium Sulphate, Sublimation; 3. Ammonium Phosphates; 4. Carbonates of Ammonium - Sesquicarbonate from Animal Matter, English Process, Uses. VI., Recovery of Ammonia from the Ammonia Soda Mother Liquors: Sections 1. General Considerations; 2. Apparatus of Schloësing and Rolland; 3. Apparatus of the Société Anonyme de l'Est. - Index.

THE MANUFACTURE OF VARNISHES, OIL REFINING AND BOILING, AND KINDRED INDUSTRIES. Describing the Manufacture of Spirit Varnishes and Oil Varnishes; Raw Materials: Resins, Solvents and Colouring Principles; Drying Oils: their Properties, Applications and Preparation by both Hot and Cold Processes; Manufacture, Employment and Testing of Different Varnishes. Translated from the French of ACH. LIVACHE, Ingénieur Civil des Mines. Greatly Extended and Adapted

to English Practice, with numerous Original Recipes by JOHN GEDDES MCINTOSH, Lecturer on Oils, Colours and Varnishes, Regent Street Polytechnic. Twenty-seven Illustrations. 400 pp. Demy 8vo. 1899. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

I. Resins: Gum Resins, Oleo Resins and Balsams, Commercial Varieties, Source, Collection, Characteristics, Chemical Properties, Physical Properties, Hardness, Adulterations Appropriate Solvents, Special Treatment, Special Use.—II. Solvents: Natural, Artificial, Manufacture, Storage, Special Use.—III. Colouring: Principles, (1) Vegetable, (2) Coal Tar, (3) Coloured Resinates, (4) Coloured Oleates and Linoleates.—Gum Running: Furnaces, Bridges, Flues, Chimney Shafts, Melting Pots, Condensers, Boiling or Mixing Pans, Copper Vessels, Iron Vessels (Cast), Iron Vessels (Wrought), Iron Vessels (Silvered), Iron Vessels (Enamelled), Steam Superheated Plant, Hot-air Plant.—Spirit Varnish Manufacture: Cold Solution Plant, Mechanical Agitators, Hot Solution Plant, Jacketted Pans, Mechanical Agitators, Clarification and Filtration, Bleaching Plant, Storage Plant.—Manufacture, Characteristics and Uses of the Spirit Varnishes yielded by: Amber, Copal, Dammar, Shellac, Mastic, Sandarac, Rosin, Asphalt, India Rubber, Gutta Percha, Collodion, Celluloid, Resinates, Oleates.—Manufacture of Varnish Stains.—Manufacture of Lacquers.—Manufacture of Spirit Enamels.—Analysis of Spirit Varnishes.—Physical and Chemical Constants of Resins.—Table of Solubility of Resins in different Menstrua.—Systematic qualitative Analysis of Resins, Hirschop's tables.—Drying Oils: Oil Crushing Plant, Oil Extraction Plant, Individual Oils, Special Treatment of Linseed Oil, Poppyseed Oil, Walnut Oil, Hempseed Oil, Llamantia Oil, Japanese Wood Oil, Gurjun Balsam, Climatic Influence on Seed and Oil.—Oil Refining: Processes.—Oil Boiling: Pale Boiled Oil, Double Boiled Oil, Hartley and Blenkinsop's Process.—Driers: Manufacture, Special Individual Use of (1) Litharge, (2) Sugar of Lead, (3) Red Lead, (4) Lead Borate, (5) Lead Linoleate, (6) Lead Resinate, (7) Black Oxide of Manganese, (8) Manganese Acetate, (9) Manganese Borate, (10) Manganese Resinate, (11) Manganese Linoleate, Mixed Resinates and Linoleates, Manganese and Lead, Zinc Sulphate, Terebine, Liquid Driers.—Solidified Boiled Oil.—Manufacture of Linoleum.—Manufacture of India Rubber Substitutes.—Printing Ink Manufacture.—Lithographic Ink Manufacture.—Manufacture of Oil Varnishes.—Running and Special Treatment of Amber, Copal, Kauri, Manilla.—Addition of Oil to Resin.—Addition of Resin to Oil.—Mixed Processes.—Solution in Cold of previously Fused Resin.—Dissolving Resins in Oil, etc., under pressure.—Filtration.—Clarification.—Storage.—Ageing.—Coachmakers' Varnishes and Japans.—Oak Varnishes.—Japanners' Stoving Varnishes.—Japanners' Gold Size.—Brunswick Black.—Various Oil Varnishes.—Oil-Varnish Stains.—Varnishes for "Enamels".—India Rubber Varnishes.—Varnishes Analysis: Processes, Matching.—Faults in Varnishes: Cause, Prevention.—Experiments and Exercises.

THE MANUFACTURE OF LAKE PIGMENTS FROM ARTIFICIAL COLOURS. By FRANCIS H. JENNISON, F.I.C., F.C.S. Sixteen Coloured Plates, showing Specimens of Eighty-nine Colours, specially prepared from the Recipes given in the Book. 136 pp. Demy 8vo. 1900. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., Introduction.—II., The Groups of the Artificial Colouring Matters.—III., The Nature and Manipulation of Artificial Colours.—IV., Lake-forming Bodies for Acid Colours.—V., Lake-forming Bodies' Basic Colours.—VI., Lake Bases.—VII., The Principles of Lake Formation.—VIII., Red Lakes.—IX., Orange, Yellow, Green, Blue, Violet and Black Lakes.—X., The Production of Insoluble Azo Colours in the Form of Pigments.—XI., The General Properties of Lakes Produced from Artificial Colours.—XII., Washing, Filtering and Finishing.—XIII., Matching and Testing Lake Pigments.—Index.

Press Opinions.

"Evidently the result of prolonged research. A valuable consulting work."—*Derby Mercury*.
 "The practical portion of the volume is the one which will especially commend itself, as that is the part of the subject which most readers would buy the book for."—*Chemist and Druggist*.

"A very valuable treatise on the manufacture of lake pigments of the coal-tar series principally."—*Chemical Trade Journal*.

"This is undoubtedly a book which will occupy a very high place amongst technical works, and will prove of exceptional value to all whom it immediately concerns."—*Eastern Morning News*.

THE TESTING AND VALUATION OF RAW MATERIALS USED IN PAINT AND COLOUR MANUFACTURE. By M. W. JONES, F.C.S. A Book for the Laboratories of Colour Works. 88 pp. Crown 8vo. 1900. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Aluminium Compounds, China Clay, Iron Compounds, Potassium Compounds, Sodium Compounds, Ammonium Hydrate, Acids, Chromium Compounds, Tin Compounds, Copper Compounds, Lead Compounds, Zinc Compounds, Manganese Compounds, Arsenic Compounds, Antimony Compounds, Calcium Compounds, Barium Compounds, Cadmium Compounds, Mercury Compounds, Ultramarine, Cobalt and Carbon Compounds, Oils Index.

THE CHEMISTRY OF ESSENTIAL OILS AND ARTIFICIAL PERFUMES. By ERNEST J. PARRY, B.Sc. (Lond.), F.I.C., F.C.S. Illustrated with Twenty Engravings. 411 pp. 1899. Demy 8vo. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

Chapters I., The General Properties of Essential Oils.—II., Compounds occurring in Essential Oils.—III., The Preparation of Essential Oils.—IV., The Analysis of Essential Oils.—V., Systematic Study of the Essential Oils.—VI., Terpeneless Oils.—VII., The Chemistry of Artificial Perfumes.—Appendix: Table of Constants.

Press Opinions.

"Will take a high place in the list of scientific text-books."—*London Argus*.

"We can heartily recommend this volume."—*British and Colonial Druggist*.

"Mr. Parry has done good service in carefully collecting and marshalling the results of the numerous researches published in various parts of the world."—*Pharmaceutical Journal*.

"At various times monographs have been printed by individual workers, but it may safely be said that Mr. Parry is the first in these latter days to deal with the subject in an adequate manner. His book is well conceived and well written. . . ."—*Chemist and Druggist*.

COSMETICS. Translated from the German of Dr. THEODOR KOLLER. Crown 8vo. 262 pp. 1902. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s. net.

Contents.

Preface.—Chapters I., Purposes and Uses of, and Ingredients used in the Preparation of Cosmetics.—II., Preparation of Perfumes by Pressure, Distillation, Maceration, Absorption or Enflourage, and Extraction Methods.—III., Chemical and Animal Products used in the Preparation of Cosmetics.—IV., Oils and Fats used in the Preparation of Cosmetics.—V., General Cosmetic Preparations.—VI., Mouth Washes and Tooth Pastes.—VII., Hair Dyes, Hair Restorers and Depilatories.—VIII., Cosmetic Adjuncts and Specialties—Colouring Cosmetic Preparations.—IX., Antiseptic Washes and Soaps—Toilet and Hygienic Soaps.—X., Secret Preparations for Skin, Complexion, Teeth, Mouth, etc.—XI., Testing and Examining the Materials Employed in the Manufacture of Cosmetics.—Index.

INK MANUFACTURE: Including Writing, Copying, Lithographic, Marking, Stamping, and Laundry Inks. By SIGMUND LEHNER. Three Illustrations. Crown 8vo. 162 pp. 1902. Translated from the German of the Fifth Edition. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s. net.

Contents.

Chapters I., Introduction.—II., Varieties of Ink.—III., Writing Inks.—IV., Raw Materials of Tannin Inks.—V., The Chemical Constitution of the Tannin Inks.—VI., Recipes for Tannin Inks.—VII., Logwood Tannin Inks.—VIII., Ferric Inks.—IX., Alizarine Inks.—X., Extract Inks.—XI., Logwood Inks.—XII., Copying Inks.—XIII., Hektographs.—XIV., Hektograph Inks.—XV., Safety Inks.—XVI., Ink Extracts and Powders.—XVII., Preserving Inks.—XVIII., Changes in Ink and the Restoration of Faded Writing.—XIX., Coloured Inks.—XX., Red Inks.—XXI., Blue Inks.—XXII., Violet Inks.—XXIII., Yellow Inks.—XXIV., Green Inks.—XXV., Metallic Inks.—XXVI., Indian Ink.—XXVII., Lithographic Inks and Pencils.—XXVIII., Ink Pencils.—XXIX., Marking Inks.—XXX., Ink Specialties.—XXXI., Sympathetic Inks.—XXXII., Stamping Inks.—XXXIII., Laundry or Washing Blue.—Index.

DRYING OILS, BOILED OIL AND SOLID AND LIQUID DRIERS. By L. E. ANDÉS. A Practical Work for Manufacturers of Oils, Varnishes, Printing Inks, Oilcloth and Linoleum, Oilcakes, Paints, etc. Expressly Written for this Series of Special Technical Books, and the Publishers hold the Copyright for English and Foreign Editions. Forty-two Illustrations. 342 pp. 1901. Demy 8vo. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

8 Contents.

Chapters I., General Chemical and Physical Properties of the Drying Oils; Cause of the Drying Property; Absorption of Oxygen; Behaviour towards Metallic Oxides, etc.—II., The Properties of and Methods for obtaining the Drying Oils.—III., Production of the Drying Oils by Expression and Extraction; Refining and Bleaching; Oil Cakes and Meal; The Refining and Bleaching of the Drying Oils: The Bleaching of Linseed Oil.—IV., The Manufacture of Boiled Oil: The Preparation of Drying Oils for Use in the Grinding of Paints and Artists' Colours and in the Manufacture of Varnishes by Heating over a Fire or by Steam, by the Cold Process, by the Action of Air, and by Means of the Electric Current; The Driers used in Boiling Linseed Oil: The Manufacture of Boiled Oil and the Apparatus therefor; Livache's Process for Preparing a Good Drying Oil and its Practical Application.—V., The Preparation of Varnishes for Letterpress, Lithographic and Copperplate Printing, for Oilcloth and Waterproof Fabrics; The Manufacture of Thickened Linseed Oil, Burnt Oil, Stand Oil by Fire Heat, Superheated Steam, and by a Current of Air.—VI., Behaviour of the Drying Oils and Boiled Oils towards Atmospheric Influences, Water, Acids and Alkalies.—VII., Boiled Oil Substitutes.—VIII., The Manufacture of Solid and Liquid Driers from Linseed Oil and Rosin; Linolic Acid Compounds of the Driers.—IX., The Adulteration and Examination of the Drying Oils and Boiled Oil.

REISSUE OF CHEMICAL ESSAYS OF C. W. SCHEELE. First Published in English in 1786. Translated from the Academy of Sciences at Stockholm, with Additions. 300 pp. Demy 8vo. 1901. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Memoir: C. W. Scheele and his work (written for this edition).—Chapters I., On Fluor Mineral and its Acid.—II., On Fluor Mineral.—III., Chemical Investigation of Fluor Acid, with a View to the Earth which it Yields, by Mr. Wiegler.—IV., Additional Information Concerning Fluor Minerals.—V., On Manganese, Magnesium, or Magnesia Vitriariorum.—VI., On Arsenic and its Acid.—VII., Remarks upon Salts of Benzoin.—VIII., On Silex, Clay and Alum.—IX., Analysis of the Calculus Vesical.—X., Method of Preparing Mercurius Dulcis Via Humida.—XI., Cheaper and more Convenient Method of Preparing Pulvis Algarothi.—XII., Experiments upon Molybdæna.—XIII., Experiments on Plumbago.—XIV., Method of Preparing a New Green Colour.—XV., Of the Decomposition of Neutral Salts by Unslaked Lime and Iron.—XVI., On the Quantity of Pure Air which is Daily Present in our Atmosphere.—XVII., On Milk and its Acid.—XVIII., On the Acid of Saccharum Lactis.—XIX., On the Constituent Parts of Lapis Ponderosus or Tungsten.—XX., Experiments and Observations on Ether.—Index.

Press Opinions.

"We would recommend the book as a refreshing recreation to the hard-worked student or investigator of the present day."—*Chemical News*.

"The present reissue renders accessible a very complete record of the researches of the great chemist."—*Chemical Trade Journal*.

"Written in a manner that cannot fail to entertain even those whose knowledge of chemical lore is of a very elementary character."—*Widnes Weekly News*.

GLUE AND GLUE TESTING. By SAMUEL RIDEAL, D.Sc. Lond., F.I.C. Fourteen Engravings. 144 pp. Demy 8vo. 1900. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Chapters I., **Constitution and Properties:** Definitions and Sources, Gelatine, Chondrin and Allied Bodies, Physical and Chemical Properties, Classification, Grades and Commercial Varieties.—II., **Raw Materials and Manufacture:** Glue Stock, Lining, Extraction, Washing and Clarifying, Filter Presses, Water Supply, Use of Alkalies, Action of Bacteria and of Antiseptics, Various Processes, Cleansing, Forming, Drying, Crushing, etc., Secondary Products.—III., **Uses of Glue:** Selection and Preparation for Use, Carpentry, Veneering, Paper-Making, Bookbinding, Printing Rollers, Hectographs, Match Manufacture, Sandpaper, etc., Substitutes for other Materials, Artificial Leather and Caoutchouc.—IV., **Gelatine:** General Characters, Liquid Gelatine. Photographic Uses, Size, Tanno-, Chrome and Formo-gelatine, Artificial Silk, Cements, Pneumatic Tyres, Culinary, Meat Extracts, Isinglass, Medicinal and other Uses, Bacteriology.—V., **Glue Testing:** Review of Processes, Chemical Examination, Adulteration, Physical Tests, Valuation of Raw Materials.—VI., **Commercial Aspects.**

Press Opinions.

"This work is of the highest technical character."—*Carpenter and Builder*.

"Dr. Rideal's book must be regarded as a valuable contribution to other technical literature, which manufacturers, merchants and users may study with profit."—*British Trade Journal*.

TECHNOLOGY OF PETROLEUM: Oil Fields of the World—Their History, Geography and Geology—Annual Production and Development—Oil-well Drilling—Transport. By HENRY NEUBERGER and HENRY NOALHAT. Translated from the French by J. G. MCINTOSH. 550 pp. 153 Illustrations. 26 Plates. Super Royal 8vo. 1901. Price 21s.; India and Colonies, 22s.; Other Countries, 23s. 6d.; strictly net.

Contents.

Part I. Study of the Petroliferous Strata—Chapters I., Petroleum—Definition.—II., The Genesis or Origin of Petroleum.—III., The Oil Fields of Galicia, their History.—IV., Physical Geography and Geology of the Galician Oil Fields.—V., Practical Notes on Galician Land Law—Economic Hints on Working, etc.—VI., Roumania—History, Geography, Geology.—VII., Petroleum in Russia—History.—VIII., Russian Petroleum (*continued*).—X., The Secondary Oil Fields of Europe, Northern Germany and Turkestan Territory—Turkestan—Persia—British India and Burma—British Burma or Lower Burma—China—Chinese Tibet—Japan, Formosa and Saghalien.—XIII., Petroleum in Oceania—Sumatra, Java, Borneo—Isle of Timor—Philippine Isles—New Zealand.—XIV., The United States of America—History.—XV., Physical Geology and Geography of the United States Oil Fields.—XVI., Canadian and other North American Oil Fields.—XVII., Economic Data of Work in North America.—XVIII., Petroleum in the West Indies and South America.—XIX., Petroleum in the French Colonies.

Part II. Excavations.—Chapter XX., Hand Excavation or Hand Digging of Oil Wells.

Part III. Methods of Boring.—Chapters XXI., Methods of Oil-well Drilling or Boring.—XXII., Boring Oil Wells with the Rope.—XXIII., Drilling with Rigid Rods and a Free-fall—Fabian System.—XXIV., Free-fall Drilling by Steam Power.—XXV., Oil-well Drilling by the Canadian System.—XXVI., Drilling Oil Wells on the Combined System.—XXVII., Comparison between the Combined Fauch System and the Canadian.—XXVIII., The American System of Drilling with the Rope.—XXIX., Hydraulic Boring with the Drill by Hand and Steam Power.—XXX., Rotary Drilling of Oil Wells, Bits, Steel-crowned Tools, Diamond Tools—Hand Power and Steam Power—Hydraulic Sand-pumping.—XXXI., Improvements in and different Systems of Drilling Oil Wells.

Part IV. Accidents.—Chapters XXXII., Boring Accidents—Methods of preventing them—Methods of remedying them.—XXXIII., Explosives and the use of the "Torpedo" Levigation.—XXXIV., Storing and Transport of Petroleum.—XXXV., General Advice—Prospecting, Management and carrying on of Petroleum Boring Operations.

Part V., General Data.—Customary Formulae.—Memento. Practical Part. General Data bearing on Petroleum.—Glossary of Technical Terms used in the Petroleum Industry.—Copious Index.

Press Opinions.

"The book is undoubtedly one of the most valuable treatises that can be placed in the hands of all who desire a knowledge of petroleum."—*Liverpool Journal of Commerce*.

"The book will undoubtedly take a high place in the literature of petroleum."—*Liverpool Post*.

"We should like to say that the translator has accomplished his difficult task, so full of technical difficulties, with a great amount of success."—*Petroleum*.

A DICTIONARY OF CHEMICALS AND RAW PRODUCTS USED IN THE MANUFACTURE OF PAINTS, COLOURS, VARNISHES AND ALLIED PREPARATIONS. By GEORGE H. HURST, F.C.S. Demy 8vo. 380 pp. 1901. Price 7s. 6d.; India and Colonies. 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

The names of the Chemicals and Raw Products are arranged in alphabetical order, and the description of each varies in length from half to eight pages. The following are some of the articles described and explained: Acetates—Acetic Acid—Acidimetry—Alcohol—Alum—Ammonia—Amber—Animi—Arsenic—Beeswax—Benzol—Bichromates of Potash and Soda—Bleaching Powder—Bone Black—Boric Acid—Brunswick Green—Cadmium Yellow—Carbonates—Carmine—Carnauba Wax—Caustic Potash and Soda—Chrome Colours—Clay—Coal Tar Colours—Copal—Dammar—Drying Oils—Emerald Green—Gamboge—Glue—Glycerine—Gums—Gypsum—Indian Red—Japanese Lacquer—Lac—Lakes—Lamp Black—Lead Compounds—Linseed Oil—Magnesia—Manganese Compounds—Mica—Nitric Acid—Ochres—Orange Lead—Orr's White—Paraffin—Prussian Blue—Rosin Oil—Sepia—Sienna—Smalts—Sodium Carbonate—Sublimed White Lead—Sulphuric Acid—Terra Verte—Testing Pigments—Turpentine—Ultramarine—Umbers—Vermilionettes—White Lead—Whiting—Zinc Compounds.—Appendix: Comparison of Baumé Hydrometer and Specific Gravity for Liquids Lighter than Water—Hydrometer Table for Liquids Heavier than Water—Comparison of Temperature Degrees—Tables for Converting French Metric Weights and Measures into English—Table of the Elements—etc., etc.—Copious Index.

Press Opinions.

"This treatise will be welcomed by those interested in this industry who have not secured the full advantage of a course of scientific training."—*Chemical Trade Journal*.

"In concise and lucid terms almost every ingredient used in paint and colour manufacture is described together with the methods of testing their intrinsic and chemical value."—*Pontefract Express*.

"Such a book of reference for paint, colour and varnish manufacturers has long been needed."—*Manchester Courier*.

RECIPES FOR THE COLOUR, PAINT, VARNISH, OIL, SOAP AND DRYSALTERY TRADES. Compiled by AN ANALYTICAL CHEMIST. 350 pp. 1902. Demy 8vo. Price 7s. 6d; India and British Colonies, 8s.; Other Countries, 8s. 6d; strictly net.

Contents.

Compiler's Preface.—Sections I., Pigments or Colours for Paints, Lithographic and Letterpress Printing Inks, etc.—II., Mixed Paints and Preparations for Paint-making, Painting, Lime-washing, Paperhanging, etc.—III., Varnishes for Coach-builders, Cabinet-makers, Wood-workers, Metal-workers, Photographers, etc.—IV., Soaps for Toilet, Cleansing, Polishing, etc.—V., Perfumes.—VI., Lubricating Greases, Oils, etc.—VII., Cements, Pastes, Glues and Other Adhesive Preparations.—VIII., Writing, Marking, Endorsing and Other Inks.—Sealing-wax and Office Requisites.—IX., Preparations for the Laundry, Kitchen, Stable and General Household Uses.—X., Disinfectant Preparations.—XI., Miscellaneous Preparations.—Index.

PURE AIR, OZONE AND WATER. A Practical Treatise of their Utilisation and Value in Oil, Grease, Soap, Paint, Glue and other Industries. By W. B. COWELL. Twelve Illustrations. Crown 8vo. 85 pp. 1900. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Chapters I., Atmospheric Air; Lifting of Liquids; Suction Process; Preparing Blown Oils; Preparing Siccativ Drying Oils.—II., Compressed Air; Whitewash.—III., Liquid Air; Retrocession.—IV., Purification of Water; Water Hardness.—V., Fleshings and Bones.—VI., Ozonised Air in the Bleaching and Deodorising of Fats, Glues, etc.; Bleaching Textile Fibres.—Appendix: Air and Gases; Pressure of Air at Various Temperatures; Fuel; Table of Combustibles; Saving of Fuel by Heating Feed Water; Table of Solubilities of Scale Making Minerals; British Thermal Units Tables; Volume of the Flow of Steam into the Atmosphere; Temperature of Steam.—Index.

THE MANUFACTURE OF MINERAL AND LAKE PIGMENTS. Containing Directions for the Manufacture of all Artificial, Artists and Painters' Colours, Enamel, Soot and Metallic Pigments. A Text-book for Manufacturers, Merchants, Artists and Painters. By Dr. JOSEF BERSCH. Translated from the Second Revised Edition by ARTHUR C. WRIGHT, M.A. (Oxon.), B.Sc. (Lond.), formerly Assistant Lecturer and Demonstrator in Chemistry at the Yorkshire College, Leeds. Forty-three Illustrations. 476 pp., demy 8vo. 1901. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

Chapters I., Introduction.—II., Physico-chemical Behaviour of Pigments.—III., Raw Materials Employed in the Manufacture of Pigments.—IV., Assistant Materials.—V., Metallic Compounds.—VI., The Manufacture of Mineral Pigments.—VII., The Manufacture of White Lead.—VIII., Enamel White.—IX., Washing Apparatus.—X., Zinc White.—XI., Yellow Mineral Pigments.—XII., Chrome Yellow.—XIII., Lead Oxide Pigments.—XIV., Other Yellow Pigments.—XV., Mosaic Gold.—XVI., Red Mineral Pigments.—XVII., The Manufacture of Vermilion.—XVIII., Antimony Vermilion.—XIX., Ferric Oxide Pigments.—XX., Other Red Mineral Pigments.—XXI., Purple of Cassius.—XXII., Blue Mineral Pigments.—XXIII., Ultramarine.—XXIV., Manufacture of Ultramarine.—XXV., Blue Copper Pigments.—XXVI., Blue Cobalt Pigments.—XXVII., Smalts.—XXVIII., Green Mineral Pigments.—XXIX., Emerald Green.—XXX., Verdigris.—XXXI., Chromium Oxide.—XXXII., Other Green Chromium Pigments.—XXXIII., Green Cobalt Pigments.—XXXIV., Green Manganese Pigments.—XXXV., Compounded Green Pigments.—XXXVI., Violet Mineral Pigments.—XXXVII., Brown Mineral Pigments.—XXXVIII., Brown Decomposition Products.—XXXIX., Black Pigments.—XL., Manufacture of Soot Pigments.—XLI., Manufacture of Lamp Black.—XLII., The Manufacture of Soot Black without Chambers.—XLIII., Indian Ink.—XLIV., Enamel Colours.—XLV., Metallic Pigments.—XLVI., Bronze Pigments.—XLVII., Vegetable Bronze Pigments.

PIGMENTS OF ORGANIC ORIGIN.—Chapters XLVIII., Lakes.—XLIX., Yellow Lakes.—L., Red Lakes.—LI., Manufacture of Carmine.—LII., The Colouring Matter of Lac.—LIII., Safflower or Carthamine Red.—LIV., Madder and its Colouring Matters.—LV., Madder Lakes.—LVI., Manjit (Indian Madder).—LVII., Lichen Colouring Matters.—LVIII., Red Wood Lakes.—LIX., The Colouring Matters of Sandal Wood and Other Dye Woods.—LX., Blue Lakes.—LXI., Indigo Carmine.—LXII., The Colouring Matter of Log Wood.—LXIII., Green Lakes.—LXIV., Brown Organic Pigments.—LXV., Sap Colours.—LXVI., Water Colours.—LXVII., Crayons.—LXVIII., Confectionery Colours.—LXIX., The Preparation of Pigments for Printing.—LXX., The Examination of Pigments.—LXXI., Examination of Lakes.—LXXII., The Testing of Dye Woods.—LXXIII., The Design of a Colour Works.—LXXIV.—Commercial Names of Pigments. Appendix: Conversion of Metric to English Weights and Measures.—Centigrade and Fahrenheit Thermometer Scales.—Index.

BONE PRODUCTS AND MANURES: An Account of the most recent Improvements in the Manufacture of Fat, Glue, Animal Charcoal, Size, Gelatine and Manures. By THOMAS LAMBERT, Technical and Consulting Chemist. Illustrated by Twenty-one Plans and Diagrams. 162 pp., demy 8vo. 1901. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I. Chemical Composition of Bones—Arrangement of Factory—Crushing of Bones—Treatment with Benzene—Benzene in Crude Fat—Analyses of Clarified Fats—Mechanical Cleansing of Bones—Animal Charcoal—Tar and Ammoniacal Liquor, Char and Gases, from good quality Bones—Method of Retorting the Bones—Analyses of Chars—"Spent" Chars—Cooling of Tar and Ammoniacal Vapours—Value of Nitrogen for Cyanide of Potash—Bone Oil—Marrow Bones—Composition of Marrow Fat—Premier Juice—Buttons.—II., Properties of Glue—Glutin and Chondrin—Skin Glue—Liming of Skins—Washing—Boiling of Skins—Clarification of Glue Liquors—Acid Steeping of Bones—Water System of Boiling Bones—Steam Method of Treating Bones—Nitrogen in the Treated Bones—Glue—Boiling and Clarifying—House—Plan showing Arrangement of Clarifying Vats—Plan showing Position of Evaporators—Description of Evaporators—Sulphurous Acid Generator—Clarification of Liquors—Section of Drying—House—Specification of a Glue—Size—Uses and Preparation and Composition of Size—Concentrated Size.—III., Properties of Gelatine—Preparation of Skin Gelatine—Washing—Bleaching—Boning—Clarification—Evaporation—Drying—Bone Gelatine—Selecting Bones—Crushing—Dissolving—Bleaching—Boiling—Properties of Glutin and Chondrin—Testing of Glues and Gelatines.—IV., The Uses of Glue, Gelatine and Size in Various Trades—Soluble and Liquid Glues—Steam and Waterproof Glues.—V., Manures—Importation of Food Stuffs—Soils—Germination—Plant Life.—VI., Natural Manures—Water and Nitrogen in Farmyard Manure—Full Analysis of Farmyard Manure—Action on Crops—Water-Closet System—Sewage Manure—Green Manures.—VII., Artificial Manures—Bones—Boiled and Steamed Bones—Mineral Phosphates—English Coprolites—French and Spanish Phosphorites—German and Belgian Phosphates—Basic Slag—Guano Proper—Guano Phosphates.—VIII., Mineral Manures—Common Salt—Potash Salts—Calcareous Manures—Prepared Nitrogenous Manures—Ammoniacal Compounds—Sodium Nitrate—Potassium Nitrate—Organic Nitrogenous Matters—Shoddy—Hoofs and Horns—Leather Waste—Dried Meat—Dried Blood—Superphosphates—Composition—Manufacture—Section of Manure-Shed—First and Second Floor Plans of Manure-Shed—Quality of Acid Used—Mixings—Special Manures—Potato Manure—Dissolved Bones—Dissolved Bone Compound—Enriched Peruvian Guano—Special Manure for Garden Stuffs, etc.—Special Manure for Grass Lands—Special Tobacco Manures—Sugar-Cane Manure—Compounding of Manures—Valuation of Manures.—IX., Analyses of Raw and Finished Products—Common Raw Bones—Degreased Bones—Crude Fat—Refined Fat—Degelatinised Bones—Animal Charcoal—Bone Superphosphates—Guano—Dried Animal Products—Potash Compounds—Sulphate of Ammonia—Extraction in Vacuum—Description of a Vacuum Pan—French and British Gelatines compared.—Index.

Press Opinion.

"We can with confidence recommend the perusal of the book to all persons interested in the manufacture of artificial manures, and also to the large number of farmers and others who are desirous of working their holdings on the most up-to-date methods, and obtaining the best possible results, which scientific research has placed within their reach."—*Hugan Observer*.

ANALYSIS OF RESINS AND BALSAMS. Translated from the German of Dr. KARL DIETERICH. Demy 8vo. 340 pp. 1901. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Part I., Definition of Resins in General—Definition of Balsams, and especially the Gum Resins—External and Superficial Characteristics of Resinous Bodies—Distinction between Resinous Bodies and Fats and Oils—Origin, Occurrence and Collection of Resinous Substances—Classification—Chemical Constituents of Resinous Substances—Resinols—Resinot Annols—Behaviour of Resin Constituents towards the Cholesterol Reactions—Uses and Identification of Resins—Melting-point—Solvents—Acid Value—Saponification Value—Resin Value—Ester and Ether Values—Acetyl and Carbonyl Value—Methyl Value—Resin Acid—Systematic Résumé of the Performance of the Acid and Saponification Value Tests.

Part II., Balsams—Introduction—Definitions—Canada Balsam—Copaiba Balsam—Angostura Copaiba Balsam—Balaia Copaiba Balsam—Carthagenia Copaiba Balsam—Maracabo Copaiba Balsam—Maturin Copaiba Balsam—Gurjun Copaiba Balsam—Para Copaiba Balsam—Surinam Copaiba Balsam—West African Copaiba Balsam—Mecca Balsam—Peruvian Balsam—Tolu Balsam—Acaroid Resin—Amine—Amber—African and West Indian Kino—Bengal Kino—Labdanum—Mastic—Pine Resin—Sandarach—Scammonium—Shellac—Storax—Adulteration of Styrax Liquidus Crudus—Purified Storax—Styrax Crudus Colatus—Tacamahac—Thapsia Resin—Turpentine—Chios Turpentine—Strassburg Turpentine—Turpeth Turpentine.

Gum Resins—Ammoniacum—Bdellium—Euphorbium—Galbanum—Gamboge—Lactucarium—Myrrh—Opopanax—Sagapenum—Olibanum or Incense—Acaroid Resin—Amber—Thapsia Resin.—Index.

MANUFACTURE OF PAINT. A Practical Handbook for Paint Manufacturers, Merchants and Painters. By J. CRUICKSHANK SMITH, B.Sc. Demy 8vo. 1901. 200 pp. Sixty Illustrations and One Large Diagram. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Part I., Chapters I., Preparation of Raw Material.—II., Storing of Raw Material.—III., Testing and Valuation of Raw Material—Paint Plant and Machinery.

Part II., Chapters V., The Grinding of White Lead.—VI., Grinding of White Zinc.—VII., Grinding of other White Pigments.—VIII., Grinding of Oxide Paints.—IX., Grinding of Staining Colours. X., Grinding of Black Paints. XI., Grinding of Chemical Colours. Yellows.—XII., Grinding of Chemical Colours. Blues.—XIII., Grinding Greens.—XIV., Grinding Reds.—XV., Grinding Lakes.—XVI., Grinding Colours in Water.—XVII., Grinding Colours in Turpentine.

Part III., Chapters XVIII., The Uses of Paint.—XIX., Testing and Matching Paints.—XX., Economic Considerations.—Index.

Press Opinions.

"Will fill a place hitherto unoccupied: . . . bears all the marks of thoroughness both on the scientific and technical side. That it will take its place as the standard book of its subject may safely be predicted."—*Aberdeen Free Press*.

"This is a workmanlike manual of the methods and processes of an industry known to the world at large only by its remote results. . . . This clearly written and well-informed handbook gives a plain exposition of these matters, and cannot but prove useful to manufacturers and merchants, and to painters curious about the materials in which they work."—*The Scotsman*.

"Mr. Smith's book is practical throughout, and it will be found helpful to those engaged in the industry and those who make use of paints."—*The Architect*.

"It is an important text-book for students attending technical classes in these subjects, concisely setting forth in a most practical manner many intricate details in the preparation and production of paint during the operation of grinding. . . . The work should occupy a position on the bookshelf of every individual interested in paint."—*The Decorators' and Painters' Magazine*.

THE CHEMISTRY OF PIGMENTS. By ERNEST J. PARRY, B.Sc. (Lond.), F.I.C., F.C.S., and J. H. COSTE, F.I.C., F.C.S. Demy 8vo. Five Illustrations. 285 pp. 1902. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Chapter I., **Introductory.** Light—White Light—The Spectrum—The Invisible Spectrum—Normal Spectrum—Simple Nature of Pure Spectral Colour—The Recomposition of White Light—Primary and Complementary Colours—Coloured Bodies—Absorption Spectra.—Chapter II., **The Application of Pigments.** Uses of Pigments: Artistic, Decorative, Protective—Methods of Application of Pigments: Pastels and Crayons, Water Colour, Tempera Painting, Fresco, Encaustic Painting, Oil-colour Painting, Ceramic Art, Enamel, Stained and Painted Glass, Mosaic.—Chapter III., **Inorganic Pigments.** White Lead—Zinc White—Enamel White—Whitening—Red Lead—Litharge—Vermilion—Royal Scarlet—The Chromium Greens—Chromates of Lead, Zinc, Silver and Mercury—Brunswick Green—The Ochres—Indian Red—Venetian Red—Siennas and Umbers—Light Red—Cappagh Brown—Red Oxides—Mars Colours—Terre Verte—Prussian Brown—Cobalt Colours—Cæruleum—Smalt—Copper Pigments—Malachite—Bremen Green—Scheele's Green—Emerald Green—Verdigris—Brunswick Green—Non-arsenical Greens—Copper Blues—Ultramarine—Carbon Pigments—Ivory Black—Lamp Black—Bistre—Naples Yellow—Arsenic Sulphides: Orpiment, Realgar—Cadmium Yellow—Vandyck Brown.—Chapter IV., **Organic Pigments.** Prussian Blue—Natural Lakes—Cochineal—Carmine—Crimson—Lac Dye—Scarlet—Madder—Alizarin—Campeachy—Quercitron—Rhamnus—Brazil Wood—Alkanet—Santal Wood—Archil—Coal-tar Lakes—Red Lakes—Alizarin Compounds—Orange and Yellow Lakes—Green and Blue Lakes—Indigo—Dragon's Blood—Gamboge—Sepia—Indian Yellow, Puree—Bitumen, Asphaltum, Mummy.—Index.

LEAD AND ITS COMPOUNDS. By THOS. LAMBERT, Technical and Consulting Chemist. Demy 8vo. 200 pp. Forty Illustrations. 1902. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d. net. Plans and Diagrams.

Contents.

Chapters I., History—Ores of Lead.—II., Geographical Distribution of the Lead Industry.—III., Chemical and Physical Properties of Lead—Alloys of Lead—Compounds of Lead.—IV., Dressing of Lead Ores.—V., Smelting of Lead Ores—Smelting in the Scotch or American Ore-hearth—Smelting in the Shaft or Blast Furnace.—VI., Condensation of Lead Fume.—VII., Desilverisation, or the Separation of Silver from Argentiferous Lead—Cupellation.—VIII., The Manufacture of Lead Pipes and Sheets.—IX., Protoxide of Lead—Litharge and Massicot—Red Lead or Minium—X., Lead Poisoning.—XI., Lead Substitutes.—XII., Zinc and its Compounds.—XIII., Pumice Stone.—XIV., Drying Oils and Siccatives.—XV., Oil of Turpentine Resin.—XVI., Classification of Mineral Pigments.—XVII., Analysis of Raw and Finished Products.—Tables.—Index

NOTES ON LEAD ORES: Their Distribution and Properties.

By JAS. FAIRIE, F.G.S. Crown 8vo. 1901. 64 pages. Price 2s. 6d.;
Abroad, 3s.; strictly net.

Contents.

Chapters I., Definitions—Properties—Occurrence—II., Galena—Johnstonite—Cerussite—Ceraise (White Lead)—Minum—Red Lead.—III., Pyromorphite—Mimetene—Hedphane—Crocoise—Wolfenite—Vanadinite—IV., Bleiglätte—Anglesite—Caledonite—Linarite—Lanarkite—Leadhillite—Sassanite—Clausenite—Cotunnite.—V., Mendipte—Matlockite—Cromfordite—Nagyagite—Aurite—Melanochroite—Vauguelinite—Scheeleite.—VI., Plattnerite—Tlklorodite—Rafanositite—Deckenite—Descloezite—Dufrenoyite—Bleimierite—Moffhrasite—Geocrinite—Wibrenenite—Schalzit—Boalangerite—Heteromorphite—Meneghinite—Jamesonite—Plagionite—Zinkenite.—VII., Kobelite—Bournonite—Selenkuperfblei—Nussierite—Percylite—Wolchite—Polysphracrite—Miesite.—Index.

THE RISKS AND DANGERS OF VARIOUS OCCUPATIONS AND THEIR PREVENTION. By LEONARD A. PARRY, M.D., B.S. (Lond.). 196 pp. Demy 8vo. 1900. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., Occupations which are Accompanied by the Generation and Scattering of Abnormal Quantities of Dust.—II., Trades in which there is Danger of Metallic Poisoning.—III., Certain Chemical Trades.—IV., Some Miscellaneous Occupations.—V., Trades in which Various Poisonous Vapours are Inhaled.—VI., General Hygienic Considerations.—Index.

PRACTICAL X RAY WORK. By FRANK T. ADDYMAN, B.Sc. (Lond.), F.I.C., Member of the Roentgen Society of London; Radiographer to St. George's Hospital; Demonstrator of Physics and Chemistry, and Teacher of Radiography in St. George's Hospital Medical School. Demy 8vo. Twelve Plates from Photographs of X Ray Work. Fifty-two Illustrations. 200 pp. 1901. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Part I., **Historical**—Chapters I., Introduction.—II., Work leading up to the Discovery of the X Rays.—III., The Discovery.

Part II., **Apparatus and its Management**—Chapters I., Electrical Terms.—II., Sources of Electricity.—III., Induction Coils.—IV., Electrostatic Machines.—V., Tubes.—VI., Air Pumps.—VII., Tube Holders and Stereoscopic Apparatus.—VIII., Fluorescent Screens.

Part III., **Practical X Ray Work**—Chapters I., Installations.—II., Radioscopy.—III., Radiography.—IV., X Rays in Dentistry.—V., X Rays in Chemistry.—VI., X Rays in War.—Index.

List of Plates.

Frontispiece—Congenital Dislocation of Hip-Joint.—I., Needle in Finger.—II., Needle in Foot.—III., Revolver Bullet in Calf and Leg.—IV., A Method of Localisation.—V., Stellate Fracture of Patella showing shadow of "Strapping".—VI., Sarcoma.—VII., Six-weeks-old Injury to Elbow showing new Growth of Bone.—VIII., Old Fracture of Tibia and Fibula badly set.—IX., Heart Shadow.—X., Fractured Femur showing Grain of Splint.—XI., Barrell's Method of Localisation.

India-Rubber and Gutta Percha.

INDIA-RUBBER AND GUTTA PERCHA. Translated from the French of T. SULIGMANN, G. LAMY, TORVILHON and H. FALCONNET. Royal 8vo. Eighty-six Illustrations. Three Plates. About 400 pages. 1902. [In the press.]

Contents.

Part I.—Chapters I., Botanical Origin.—II., Climatology—Soil—Rational Culture and Acclimation of the Different Species of India-Rubber Plants.—III., Methods of Obtaining the Latex—Methods of Preparing Raw or Crude India-Rubber.—IV., Classification of the Commercial Species of Raw Rubber.—V., Physical and Chemical Properties of the Latex and of India-Rubber.—VI. and VII., Mechanical Transformation of Natural Caoutchouc into Washed or Normal Caoutchouc (Purification) and Normal Rubber into Masticated Rubber—Softening, Cutting, Washing, Drying—Preliminary Observations.—VIII., Vulcanisation of Normal Rubber.—IX., Chemical and Physical Properties of Vulcanised Rubber—General Considerations.—X., Hardened Rubber or Ebonite.—XI., Considerations on Mineralisation and other Mixtures—Coloration and Dyeing—Analysis of Natural or Normal Rubber and Vulcanised Rubber.—XII., Rubber Substitutes—Imitation Rubber.

Part II.—**Gutta Percha**.—Chapters I., Botanical Origin.—II., Climatology—Soil—Rational Culture.—III., Methods of Collection.—IV., Classification of the Different Species of Commercial Gutta Percha.—V., Physical and Chemical Properties.—VI., Mechanical Transformation.—VII., Methods of Analysing.—VIII., Gutta Percha Substitutes.—Index.

DRYING BY MEANS OF AIR AND STEAM. Explanations, Formulæ, and Tables for Use in Practice. Translated from the German of E. HAUSBRAND. Two folding Diagrams and Thirteen Tables. Crown 8vo. 1901. 72 pp. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Preface.—British and Metric Systems Compared.—Centigrade and Fahr. Thermometers.—Chapters I., Introduction.—II., Estimation of the Maximum Weight of Saturated Aqueous Vapour which can be contained in 1 kilo. of Air at Different Pressure and Temperatures.—III., Calculation of the Necessary Weight and Volume of Air, and of the Least Expenditure of Heat, per Drying Apparatus with Heated Air, at the Atmospheric Pressure: A, With the Assumption that the Air is Completely Saturated with Vapour both before Entry and after Exit from the Apparatus.—B, When the Atmospheric Air is Completely Saturated before entry, but at its exit is only $\frac{2}{3}$, $\frac{1}{2}$ or $\frac{1}{4}$ Saturated.—C, When the Atmospheric Air is not Saturated with Moisture before Entering the Drying Apparatus.—IV., Drying Apparatus, in which, in the Drying Chamber, a Pressure is Artificially Created, Higher or Lower than that of the Atmosphere.—V., Drying by Means of Superheated Steam, without Air.—VI., Heating Surface, Velocity of the Air Current, Dimensions of the Drying Room, Surface of the Drying Material, Losses of Heat.—Index.

EVAPORATING, CONDENSING AND COOLING APPARATUS. Explanations, Formulæ and Tables for Use in Practice. By E. HAUSBRAND, Engineer. Translated from the Second Revised German Edition by A. C. WRIGHT, M.A. (Oxon.), B.Sc. (Lond.), formerly Assistant Lecturer and Demonstrator in Chemistry at the Yorkshire College, Leeds. With Twenty-one Illustrations and Seventy-six Tables. Demy 8vo. 1902. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s. net.

Contents.

Prefaces.—List of Tables.—Metric and British Systems and Thermometers compared.—Symbols and Contractions.—Introduction.—Chapters I., Coefficient of Transmission of Heat, k , and the Mean Temperature Difference, θ/m .—II., Parallel and Opposite Currents.—III., Apparatus for Heating with Direct Fire.—IV., The Injection of Saturated Steam.—V., Superheated Steam.—VI., Evaporation by Means of Hot Liquids.—VII., The Transference of Heat in General, and Transference by means of Saturated Steam in Particular.—VIII., The Transference of Heat from Saturated Steam in Pipes (Coils) and Double Bottoms.—IX., Evaporation in a Vacuum.—X., The Multiple-effect Evaporator.—XI., Multiple-effect Evaporators from which Extra Steam is Taken.—XII., The Weight of Water which must be Evaporated from 100 Kilos. of Liquid in order its Original Percentage of Dry Materials from 1.25 per cent. up to 20.70 per cent.—XIII., The Relative Proportion of the Heating Surfaces in the Elements of the Multiple Evaporator and their Actual Dimensions.—XIV., The Pressure Exerted by Currents of Steam and Gas upon Floating Drops of Water.—XV., The Motion of Floating Drops of Water upon which Press Currents of Steam.—XVI., The Splashing of Evaporating Liquids.—XVII., The Diameter of Pipes for Steam, Alcohol, Vapour and Air.—XVIII., The Diameter of Water Pipes.—XIX., The Loss of Heat from Apparatus and Pipes to the Surrounding Air, and Means for Preventing the Loss.—XX., Condensers.—XXI., Heating Liquids by Means of Steam.—XXII., The Cooling of Liquids.—XXIII., The Volumes to be Exhausted from Condensers by the Air-pumps.—XXIV., A Few Remarks on Air pumps and the Vacua they Produce.—XXV., The Volumetric Efficiency of Air-pumps.—XXVI., The Volumes of Air which must be Exhausted from a Vessel in order to Reduce its Original Pressure to a Certain Lower Pressure.—Index.

Leather Trades.

THE LEATHER WORKER'S MANUAL. Being a Compendium of Practical Recipes and Working Formulæ for Curriers, Bootmakers, Leather Dressers, Blacking Manufacturers, Saddlers, Fancy Leather Workers, and all Persons engaged in the Manipulation of Leather. By H. C. STANDAGE. 165 pp. 1900. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., Blackings, Polishes, Glosses, Dressings, Renovators, etc., for Boot and Shoe Leather.—II., Harness Blackings, Dressings, Greases, Compositions, Soaps, and Boot top Powders and Liquids, etc., etc.—III., Leather Grinders' Sundries.—IV., Currier's Seasonings, Blacking Compounds, Dressings, Finishes, Glosses, etc.—V., Dyes and Stains for Leather.—VI., Miscellaneous Information.—VII., Chrome Tannage.—Index.

Press Opinions.

"The book being absolutely unique."—*Birmingham Gazette*.
 "This is a valuable collection of practical receipts and working formulæ for the use of those engaged in the manipulation of leather."—*Liverpool Mercury*.

PRACTICAL TREATISE ON THE LEATHER INDUSTRY. By A. M. VILLON. A Translation of Villon's "Traite Pratique de la Fabrication des cuirs et du Travail des Peaux". By FRANK T. ADDYMAN, B.Sc. (Lond.), F.I.C., F.C.S.; and Corrected by an Eminent Member of the Trade. 500 pp., royal 8vo. 1901. 123 Illustrations. Price 21s.; India and Colonies, 22s.; Other Countries, 23s. 6d.; strictly net.

Contents.

Preface—Translator's Preface—List of Illustrations.

Part I. **Materials used in Tanning**—Chapter I., Skins: I., Skin and its Structure; II., Skins used in Tanning; III., Various Skins and their Uses—Chapter II., Tannin and Tanning Substances: I., Tannin; II., Barks (Oak); III., Barks other than Oak; IV., Tanning Woods; V., Tannin-bearing Leaves; VI., Excrecences; VII., Tan-bearing Fruits; VIII., Tan-bearing Roots and Bales; IX., Tanning Juices; X., Tanning Substances used in Various Countries; XI., Tannin Extracts; XII., Estimation of Tannin and Tannin Principles.

Part II. **Tanning**—Chapter I., The Installation of a Tannery: I., Tan Furnaces; II., Chimneys, Boilers, etc.; III., Steam Engines—Chapter II., Grinding and Trituration of Tanning Substances: I., Cutting up Bark; II., Grinding Bark; III., The Grinding of Tan Woods; IV., Powdering Fruit, Galls and Grains; V., Notes on the Grinding of Bark—Chapter III., Manufacture of Sole Leather: I., Soaking; II., Sweating and Unhairing; III., Plumping and Combing; IV., Handling; V., Tanning; VI., Tanning Elephants' Hides; VII., Drying; VIII., Straining or Pinning—Chapter IV., Manufacture of Dressing Leather: I., Soaking; II., Depilation; III., New Processes for the Depilation of Skins; IV., Tanning; V., Cow Hides; VI., Horse Hides; VII., Goat Skins; Manufacture of Split Hides—Chapter V., On Various Methods of Tanning: I., Mechanical Methods; II., Physical Methods; III., Chemical Methods; IV., Tanning with Extracts—Chapter VI., Quantity and Quality: I., Quantity; II., Net Cost; III., Quality of Leather—Chapter VII., Various Manipulations of Tanned Leather: I., Second Tanning; II., Grease Stains; III., Bleaching Leather; IV., Waterproofing Leather; V., Weighting Tanned Leather; VI., Preservation of Leather—Chapter VIII., Tanning Various Skins.

Part III. **Currying**—Chapter I., Waxed Calf: I., Preparation; II., Shaving; III., Stretching or Slaking; IV., Oiling the Grain; V., Oiling the Flesh Side; VI., Whitening and Graining; VII., Waxing; VIII., Finishing; IX., Dry Finishing; X., Finishing in Colour; XI., Cost—Chapter II., White Calf: I., Finishing in White—Chapter III., Cow Hide for Upper Leathers: I., Black Cow Hide; II., White Cow Hide; III., Coloured Cow Hide—Chapter IV., Smooth Cow Hide—Chapter V., Black Leather—Chapter VI., Miscellaneous Hides: I., Horse; II., Goat; III., Waxed Goat Skin; IV., Matt Goat Skin—Chapter VII., Russia Leather: I., Russia Leather; II., Artificial Prussia Leather.

Part IV. **Enamelled, Hungary and Chamov Leather, Morocco, Parchment, Furs and Artificial Leather**—Chapter I., Enamelled Leather: I., Varnish Manufacture; II., Application of the Enamel; III., Enamelling in Colour—Chapter II., Hungary Leather: I., Preliminary; II., Wet Work or Preparation; III., Aluming; IV., Dressing or Loft Work; V., Tallowing; VI., Hungary Leather from Various Hides—Chapter III., Tawing: I., Preparatory Operations; II., Dressing; III., Dyeing Tawed Skins; IV., Rugs—Chapter IV., Chamov Leather—Chapter V., Morocco: I., Preliminary Operations; II., Morocco Tanning; III., Mordants used in Morocco Manufacture; IV., Natural Colours used in Morocco Dyeing; V., Artificial Colours; VI., Different Methods of Dyeing; VII., Dyeing with Natural Colours; VIII., Dyeing with Aniline Colours; IX., Dyeing with Metallic Salts; X., Leather Printing; XI., Finishing Morocco; XII., Shagreen; XIII., Bronzed Leather—Chapter VI., Gilding and Silvering: I., Gilding; II., Silvering; III., Nickel and Cobalt—Chapter VII., Parchment—Chapter VIII., Furs and Furriery: I., Preliminary Remarks; II., Indigenous Furs; III., Foreign Furs from Hot Countries; IV., Foreign Furs from Cold Countries; V., Furs from Birds' Skins; VI., Preparation of Furs; VII., Dressing; VIII., Colouring; IX., Preparation of Birds' Skins; X., Preservation of Furs—Chapter IX., Artificial Leather: I., Leather made from Scraps; II., Compressed Leather; III., American Cloth; IV., Papier Mâché; V., Linoleum; VI., Artificial Leather.

Part V. **Leather Testing and the Theory of Tanning**—Chapter I., Testing and Analysis of Leather: I., Physical Testing of Tanned Leather; II., Chemical Analysis—Chapter II., The Theory of Tanning and the other Operations of the Leather and Skin Industry: I., Theory of Soaking; II., Theory of Unhairing; III., Theory of Swelling; IV., Theory of Handling; V., Theory of Tanning; VI., Theory of the Action of Tannin on the Skin; VII., Theory of Hungary Leather Making; VIII., Theory of Tawing; IX., Theory of Chamov Leather Making; X., Theory of Mineral Tanning.

Part VI. **Uses of Leather**—Chapter I., Machine Belts: I., Manufacture of Belting; II., Leather Chain Belts; III., Various Belts; IV., Use of Belts—Chapter II., Boot and Shoemaking: I., Boots and Shoes; II., Laces—Chapter III., Saddlery: I., Composition of a Saddle; II., Construction of a Saddle—Chapter IV., Harness: I., The Pack Saddle; II., Harness—Chapter V., Military Equipment—Chapter VI., Glove Making—Chapter VII., Carriage Building—Chapter VIII., Mechanical Uses.

Appendix, **The World's Commerce in Leather**—I., Europe; II., America; III., Asia; IV., Africa; Australasia—Index.

Press Opinions.

"The book is well and lucidly written."—*Shoe Manufacturers' Monthly Journal*.

"This book cannot fail to be of great value to all engaged in the leather trades. . . ."—*Midland Free Press*.

"Gives much useful and interesting information concerning the various processes by which the skins of animals are converted into leather."—*Leeds Mercury*.

"It can thus be confidently recommended to all who are more or less practically interested in the technology of a very important subject."—*Leicester Post*.

"This is, in every respect, an altogether admirable, practical, clear and lucid treatise on the various and numerous branches of the great leather industry, of which it deals in an exhaustive, highly intelligent, workmanlike and scientific manner. . . ."—*Shoe and Leather Trader*.

"It certainly should be worth the while of English tanners to see what lessons they can learn from French practice, and French practice, we should imagine, could hardly have a better exponent than the author of this large volume."—*Western Daily Press and Bristol Times*.

"... As a work of reference the volume will be extremely useful in the trade, and where leisure affords sufficient opportunity a careful perusal and study of it would afford ample reward."—*Kettering Guardian*.

Books on Pottery, Bricks, Tiles, Glass, etc.

THE MANUAL OF PRACTICAL POTTING. Revised and Enlarged. Third Edition. 200 pp. 1901. Price 17s. 6d.; India and Colonies, 18s. 6d.; Other Countries, 20s.; strictly net.

Contents.

Introduction. The Rise and Progress of the Potter's Art.—**Chapters I., Bodies.** China and Porcelain Bodies, Parian Bodies, Semi-porcelain and Vitreous Bodies, Mortar Bodies, Earthenwares Granite and C.C. Bodies, Miscellaneous Bodies, Sagger and Crucible Clays, Coloured Bodies, Jasper Bodies, Coloured Bodies for Mosaic Painting, Encaustic Tile Bodies, Body Stains, Coloured Dips.—**II., Glazes.** China Glazes, Ironstone Glazes, Earthenware Glazes, Glazes without Lead, Miscellaneous Glazes, Coloured Glazes, Majolica Colours.—**III., Gold and Cold Colours.** Gold, Purple of Cassius, Marone and Ruby, Enamel Coloured Bases, Enamel Colour Fluxes, Enamel Colours, Mixed Enamel Colours, Antique and Vellum Enamel Colours, Underglaze Colours, Underglaze Colour Fluxes, Mixed Underglaze Colours, Flow Powders, Oils and Varnishes.—**IV., Means and Methods.** Reclamation of Waste Gold, The Use of Cobalt, Notes on Enamel Colours, Liquid or Bright Gold.—**V., Classification and Analysis.** Classification of Clay Ware, Lord Playfair's Analysis of Clays, The Markets of the World, Time and Scale of Firing, Weights of Potter's Material, Decorated Goods Count.—**VI., Comparative Loss of Weight of Clays.**—**VII., Ground Felspar Calculations.**—**VIII., The Conversion of Slop Body Recipes into Dry Weight.**—**IX., The Cost of Prepared Earthenware Clay.**—**X., Forms and Tables.** Articles of Apprenticeship, Manufacturer's Guide to Stocktaking, Table of Relative Values of Potter's Materials, Hourly Wages Table, Workman's Settling Table, Comparative Guide for Earthenware and China Manufacturers in the use of Slop Flint and Slop Stone, Foreign Terms applied to Earthenware and China Goods, Table for the Conversion of Metrical Weights and Measures on the Continent of South America.—**Index.**

CERAMIC TECHNOLOGY: Being some Aspects of Technical Science as Applied to Pottery Manufacture. Edited by CHARLES F. BINNS. 100 pp. Demy 8vo. 1897. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

Preface.—Introduction.—Chapters I., The Chemistry of Pottery.—**II., Analysis and Synthesis.**—**III., Clays and their Components.**—**IV., The Biscuit Oven.**—**V., Pyrometry.**—**VI., Glazes and their Composition.**—**VII., Colours and Colour-making.**—**Index.**

RECIPES FOR FLINT GLASS MAKING. By a British Glass Master and Mixer. Sixty Recipes. Being Leaves from the Mixing Book of several experts in the Flint Glass Trade, containing up-to-date recipes and valuable information as to Crystal, Demi-crystal and Coloured Glass in its many varieties. It contains the recipes for cheap metal suited to pressing, blowing, etc., as well as the most costly crystal and ruby. British manufacturers have kept up the quality of this glass from the arrivals of the Venetians to Hungry Hill, Stourbridge, up to the present time. The book also contains remarks as to the result of the metal as it left the pots by the respective metal mixers, taken from their own memoranda upon the originals. 1900. Price for United Kingdom, 10s. 6d.; Abroad, 15s.; United States, \$4; strictly net.

Contents.

Ruby—Ruby from Copper—Flint for using with the Ruby for Coating—A German Metal—Cornelian, or Alabaster—Sapphire Blue—Crysophus—Opal—Turquoise Blue—Gold Colour—Dark Green—Green (common)—Green for Malachite—Blue for Malachite—Black for Malachite—Black—Common Canary Batch—Canary—White Opaque Glass—Sealing-wax Red—Flint—Flint Glass—Crystal and Demo—Achromatic Glass—Paste Glass—White Enamel—Firestone—Dead White (for moons)—White Agate—Canary—Canary Enamel—Index.

COLOURING AND DECORATION OF CERAMIC

WARE. By ALEX. BRONGNIART. With Notes and Additions by ALPHONSE SALVETAT. Translated from the French. 200 pp. 1898. Price 7s. 6d.: India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

The Pastes, Bodies or Ceramic Articles Capable of being Decorated by Vitrifiable Colours—The Chemical Preparation of Vitrifiable Colours—Composition and Preparation of Vitrifiable Colours—The Oxides—Preparation of Oxides—Preparation of Chromates—Preparation of other Colours—Composition and Preparation of Fluxes—Muffle Colours—Recipes for Colours—Use of Metals—Lustres—Preparation and Application of Colours—Composition of Coloured Pastes—Underglaze Colours—Colours in the Glaze—Overglaze Colours—Painting in Vitrifiable Colours—Gilding—Burnishing—Printing—Enlarging and Reducing Gelatine Prints—Muffle Kilns for Vitrifiable Colours—Influence of the Material on the Colour—Changes Resulting from the Actions of the Fire—Alterations Resulting from the Colours—Alterations in Firing.

HOW TO ANALYSE CLAY. Practical Methods for Practical Men. By HOLDEN M. ASHBY, Professor of Organic Chemistry, Harvey Medical College, U.S.A. Twenty Illustrations. 1898. Price 2s. 6d.; Abroad, 3s.; strictly net.

Contents

List of Apparatus—List of Atomic Weights—Use of Balance, and Burette, Sand Bath, and Water Bath—Dessicator—Drying Oven—Filtering—Fusion—Determination of Water, Organic Matter, Iron, Calcium, Alkalies, Limestone, Silica, Alumina—Magnesium etc.—Mechanical Analysis—Rational Analysis—Standard Solutions—Volumetric Analysis—Standards for Clay Analysis—Sampling.

ARCHITECTURAL POTTERY. Bricks, Tiles, Pipes, Enamelled Terra-cottas, Ordinary and Incrusted Quarries, Stoneware Mosaics, Faïences and Architectural Stoneware. By LEON LEFEVRE. With Five Plates. 950 Illustrations in the Text, and numerous estimates. 500 pp., royal 8vo. 1900. Translated from the French by K. H. BIRD, M.A., and W. MOORE BINNS. Price 15s.; India and Colonies, 16s.; Other Countries, 17s. 6d.; strictly net.

Contents.

Part I. **Plain Undecorated Pottery.** Chapter I., Clays : 1. Classification, General Geological Remarks.—Classification, Origin, Locality : 2. General Properties and Composition : Physical Properties, Contraction, Analysis, Influence of Various Substances on the Properties of Clays : 3. Working of Clay-Pits—I. Open Pits: Extraction, Transport, Cost—II. Underground Pits—Mining Laws. Chapter II., Preparation of the Clay: Weathering, Mixing, Cleaning, Crushing and Pulverising—Crushing Cylinders and Mills, Pounding Machines—Damping. Damping Machines—Soaking, Shortening, Pugging: Horse and Steam Pug-Mills, Rolling Cylinders—Particulars of the Above Machines. Chapter III., Bricks: § 1, Manufacture—(1) Hand and Machine Moulding.—I. Machines Working by Compression: on Soft Clay, on Semi-Firm Clay, on Firm Clay, on Dry Clay.—II. Expression Machines: with Cylindrical Propellers, with Screw Propellers—Dies—Cutting-tables—Particulars of the Above Machines—General Remarks on the Choice of Machines—Types of Installations—Estimates—Plemishing, Hand and Steam Presses, Particulars—(2) Drying, by Exposure to Air, Without Shelter, and Under Sheds—Drying-rooms in Tiers, Closed Drying-rooms, in Tunnels, in Galleries—Detailed Estimates of the Various Drying-rooms, Comparison of Prices—Transport from the Machines to the Drying-rooms, Barrows, Trucks, Plain or with Shelves, Lifts—(3) Firing—I. In Clamps—II. In Intermittent Kilns. A, Open: a, using Wood; b, Coal; b', in Clamps; b'', Flame—B, Closed: c, Direct Flame; c', Rectangular; c'', Round; d, Reverberatory—III. Continuous Kilns: C, with Solid Fuel: Round Kiln, Rectangular Kiln, Chimneys (Plans and Estimates)—D, With Gas Fuel, Pillard Kiln (Plans and Estimates), Schneider Kiln (Plans and Estimates), Water-gas Kiln—Heat Production of the Kilns: § 2, Dimensions, Shapes, Colours, Decoration, and Quality of Bricks—Hollow Bricks, Dimensions and Prices of Bricks, Various Shapes, Qualities—Various Hollow Bricks, Dimensions, Resistance, Qualities: § 3, Applications—History—Asia, Africa, America, Europe: Greek, Roman, Byzantine, Turkish, Romanesque, Gothic, Renaissance, Architecture—Architecture of the Nineteenth Century: in Germany, England, Belgium, Spain, Holland, France, America—Use of Bricks—Walls, Arches, Pavements

Flues, Cornices—Facing with Coloured Bricks—Balustrades. Chapter IV., Tiles: § 1, History: § 2, Manufacture—(1) Moulding, by Hand, by Machinery: Preparation of the Clay, Soft Paste, Firm Paste, Hard Paste—Preparation of the Slabs, Transformation into Flat Tiles, into Jointed Tiles—Screw, Cam and Revolver Presses—Particulars of Tile-presses—(2) Drying—Planchettes, Shelves, Drying-barrows and Trucks—(3) Firing—Divided Kilns—Installation of Mechanical Tileworks—Estimates: § 3, Shapes, Dimensions and Uses of the Principal Types of Tile—Ancient Tiles: Flat, Round, Roman, Flemish—Modern Tiles—With Vertical Interrupted Join: Gilardoni's, Martin's: Hooked, Boulet's Villa: with Vertical Continuous Join: Muller's, Alsace, Pantile—Foreign Tiles—Special Tiles—Ridge Tiles. Coping Tiles, Border Tiles, Frontons, Gutters, Antefixes, Membrons, Angular—Roofing Accessories: Chimney-pots, Mitrons, Lanterns, Chimneys—Qualities of Tiles—Black Tiles—Stoneware Tiles—Particulars of Tiles. Chapter V., Pipes: I. Conduit Pipes—Manufacture—Moulding: Horizontal Machines, Vertical Machines, Worked by Hand and Steam—Particulars of these Machines—Drying—Firing—II. Chimney Flues—Ventiducts and "Boisseaux," "Waggon"—Particulars of these Products. Chapter VI., Quarries: 1, Plain Quarries of Ordinary Clay: 2, of Cleaned Clay—Machines, Cutting, Mixing, Polishing—Drying and Firing—Applications—Particulars of Quarries. Chapter VII., Terra-cotta: History—Manufacture—Application: Balustrades, Columns, Pilasters, Capitals, Friezes, Frontons, Medallions, Panels, Rose-windows, Ceilings—Appendix: Official Methods of Testing Terra-cottas.

Part II. **Made-up or Decorated Pottery.**—Chapter I., General Remarks on the Decoration of Pottery: Dips—Glazes: Composition, Colouring, Preparation, Harmony with Pastes—Special Processes of Decoration—Enamels, Opaque, Transparent, Colours, Under-glaze, Over-glaze—Other Processes: Crackling, Mottled, Flashing, Metallic Iridescence, Lustres. Chapter II., Glazed and Enamelled Bricks—History: Glazing—Enamelling—Applications: Ordinary Enamelled Bricks, Glazed Stoneware, Enamelled Stoneware—Enamelled Tiles. Chapter III., Decorated Quarries: I. Paving Quarries—1, Decorated with Dips—2, Stoneware: 1, Fired to Stoneware: *a*, of Slag Base—Applications: *b*, of Melting Clay—Applications—*b*, Plain or Incrusted Stoneware: *c*, of Special Clay (Stoke-on-Trent)—Manufacture—Application—*b*, of Felspar Base—Colouring, Manufacture, Moulding, Drying, Firing—Applications.—II. Facing Quarries—1, in Faience—1, of Limestone Paste—3, of Silicious Paste—*c*, of Felspar Paste—Manufacture, Firing—2, of Glazed Stoneware—3, of Porcelain—Applications of Facing Quarries.—III. Stove Quarries—Preparation of the Pastes, Moulding, Firing, Enamelling, Decoration—Applications—Faïences for Fireplaces. Chapter IV., Architectural Decorated Pottery: § 1, Faïences: § 2, Stoneware: § 3, Porcelain. Chapter V., Sanitary Pottery: Stoneware Pipes: Manufacture, Firing—Applications—Sinks—Applications—Urinals, Seats and Pans—Applications—Drinking-fountains, Washstands.—Index.

A TREATISE ON THE CERAMIC INDUSTRIES. A

Complete Manual for Pottery, Tile and Brick Works. By EMILE BOURRY, Ingénieur des Arts et Manufactures. Translated from the French by WILTON P. RIX, Examiner in Pottery and Porcelain to the City and Guilds of London Technical Institute, Pottery Instructor to the Hanley School Board. Royal 8vo. 1901. Over 700 pp. Price 21s.: India and Colonies, 22s.; Other Countries, 23s. 6d.; strictly net.

Contents.

Part I., **General Pottery Methods.** Chapters I., Definition and History. Definitions and Classification of Ceramic Products—Historic Summary of the Ceramic Art.—II., Raw Materials of Bodies. Clays: Pure Clay and Natural Clays—Various Raw Materials: Analogous to Clay—Agglomerative and Agglutinative—Opening—Fusible—Refractory—Trials of Raw Materials.—III., Plastic Bodies. Properties and Composition—Preparation of Raw Materials: Disaggregation—Purification—Preparation of Bodies: By Plastic Method—By Dry Method—By Liquid Method.—IV., Formation. Processes of Formation: Throwing—Expression—Moulding by Hand, on the Jolley, by Compression, by Slip Casting—Slapping—Slipping.—V., Drying. Drying of Bodies—Processes of Drying: By Evaporation—By Aeration—By Heating—By Ventilation—By Absorption.—VI., Glazes. Composition and Properties—Raw Materials—Manufacture and Application.—VII., Firing. Properties of the Bodies and Glazes during Firing—Description of the Kilns—Working of the Kilns.—VIII., Decoration. Colouring Materials—Processes of Decoration.

Part II., **Special Pottery Methods.** Chapters IX., Terra Cottas. Classification: Plain Ordinary, Hollow, Ornamental, Vitrified, and Light Bricks—Ordinary and Black Tiles—Paving Tiles—Pipes—Architectural Terra Cottas—Vases, Statues and Decorative Objects—Common Pottery—Pottery for Water and Filters—Tobacco Pipes—Lustre Ware—Properties and Tests for Terra Cottas.—X., Fireclay Goods. Classification: Argillaceous, Aluminous, Carboniferous, Silicious and Basic Fireclay Goods—Fireclay Mortar (Pug)—Tests for Fireclay Goods.—XI., Faïences. Varnished Faïences—Enamelled Faïences—Silicious Faïences—Pipeclay Faïences—Pebble Work—Feldspathic Faïences—Composition, Processes of Manufacture and General Arrangements of Faïence Potteries.—XII., Stoneware. Stoneware Properly So-called: Paving Tiles—Pipes—Sanitary Ware—Stoneware for Food Purposes and Chemical Productions—Architectural Stoneware—Vases, Statues and other Decorative Objects—Fine Stoneware.—XIII., Porcelain. Hard Porcelain for Table Ware and Decoration, for the Fire, for Electrical Conduits, for Mechanical Purposes: Architectural Porcelain, and Dull or Biscuit Porcelain—Soft Phosphated or English Porcelain—Soft Vitreous Porcelain, French and New Sevres—Argillaceous Soft or Seger's Porcelain—Dull Soft or Parian Porcelain—Dull Feldspathic Soft Porcelain.—Index.

THE ART OF RIVETING GLASS, CHINA AND EARTHENWARE. By J. HOWARTH. Second Edition. 1900. Price 1s. net; by post, home or abroad, 1s. 1d.

Contents.

Tools and Materials Required—Wire Used for Rivets—Soldering Solution—Preparation for Drilling—Commencement of Drilling—Cementing—Preliminaries to Riveting—Rivets to Make—To Fix the Rivets—Through-and-through Rivets—Soldering—Tinning a Soldering-Iron—Perforated Plates, Handles, etc.—Handles of Ewers, etc.—Vases and Comports—Marble and Alabaster Ware—Decorating—How to Loosen Fast Decanter Stoppers—China Cements.

NOTES ON POTTERY CLAYS. Their Distribution, Properties, Uses and Analyses of Ball Clays, China Clays and China Stone. By JAS. FAIRIE, F.G.S. 1901. 132 pp. Crown 8vo. Price 3s. 6d.; India and Colonies, 4s.; Other Countries, 4s. 6d.; strictly net.

Contents.

Definitions—Occurrence—Brick Clays—Fire Clays—Analyses of Fire Clays.—**Ball Clays**—Properties—Analyses—Occurrence—Pipe Clay—Black Clay—Brown Clay—Blue Clay—Dorsetshire and Devonshire Clays—**China Clay** or Kaolin—Occurrence—Chinese Kaolin—Cornish Clays—Hensbarrow Granite—Properties, Analyses and Composition of China Clays—Method of Obtaining China Clay—Experiments with Chinese Kaolin—Analyses of Chinese and Japanese Clays and Bodies—Irish Clays—**Chinese Stone**—Composition—Occurrence—Analyses.—Index.

PAINTING ON GLASS AND PORCELAIN AND ENAMEL PAINTING. A Complete Introduction to the Preparation of all the Colours and Fluxes used for Painting on Porcelain, Enamel, Faience and Stoneware, the Coloured Pastes and Coloured Glasses, together with a Minute Description of the Firing of Colours and Enamels. On the Basis of Personal Practical Experience of the Condition of the Art up to Date. By FELIX HERMANN, Technical Chemist. With Eighteen Illustrations. 300 pp. Translated from the German second and enlarged Edition. 1897. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

History of Glass Painting.—Chapters I., The Articles to be Painted: Glass, Porcelain, Enamel, Stoneware, Faience.—II., Pigments: 1, Metallic Pigments: Antimony Oxide, Naples Yellow, Barium Chromate, Lead Chromate, Silver Chloride, Chromic Oxide.—III., Fluxes: Fluxes, Felspar, Quartz, Purifying Quartz, Sedimentation, Quenching, Borax, Boracic Acid, Potassium and Sodium Carbonates, Rocaille Flux.—IV., Preparation of the Colours for Glass Painting.—V., The Colour Pastes.—VI., The Coloured Glasses.—VII., Composition of the Porcelain Colours.—VIII., The Enamel Colours: Enamels for Artistic Work.—IX., Metallic Ornamentation: Porcelain Gilding, Glass Gilding.—X., Firing the Colours: 1, Remarks on Firing: Firing Colours on Glass, Firing Colours on Porcelain: 2, The Muffle.—XI., Accidents occasionally Supervening during the Process of Firing.—XII., Remarks on the Different Methods of Painting on Glass, Porcelain, etc. Appendix: Cleaning Old Glass Paintings.

Press Opinions.

"Will be found of much interest to the amateur."—*Art Amateur*, New York.
 "... The whole cannot fail to be both of service and interest to glass workers and to potters generally, especially those employed upon high-class work."—*Staffordshire Sentinel*.
 "For manufacturers and students it will be a valuable work, and the recipes which appear on almost every page form a very valuable feature."—*Builders Journal*.
 "... Very careful instructions are given for the chemical and mechanical preparation of the colours used in glass-staining and porcelain-painting; indeed, to the china painter such a book as this should be of permanent value."—*Daily Chronicle*.

A TREATISE ON THE ART OF GLASS PAINTING.

Prefaced with a Review of Ancient Glass. By ERNEST R. SUFFLING. With One Coloured Plate and Thirty-seven Illustrations. Demy 8vo. 140 pp. 1902. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d. net.

Contents.

Chapters I., A Short History of Stained Glass.—II., Designing Scale Drawings.—III., Cartoons and the Cut Line.—IV., Various Kinds of Glass Cutting for Windows.—V., The Colours and Brushes used in Glass Painting.—VI., Painting on Glass, Dispersed Patterns.—VII., Diapered Patterns—Aciding—Firing.—VIII., Fret Lead Glazing.—Index.

A Reissue of

THE HISTORY OF THE STAFFORDSHIRE POTTERIES; AND THE RISE AND PROGRESS OF THE MANUFACTURE OF POTTERY AND PORCELAIN.

With References to Genuine Specimens, and Notices of Eminent Potters. By SIMEON SHAW. (Originally Published in 1829.) 265 pp. 1900. Demy 8vo. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Introductory Chapter showing the position of the Pottery Trade at the present time (1899).—**Chapters I., Preliminary Remarks.**—**II., The Potteries**, comprising Tunstall, Brownhills, Greenfield and New Field, Golden Hill, Latebrook, Green Lane, Burslem, Longport and Dale Hall, Hot Lane and Cobridge, Hanley and Shelton, Etruria, Stoke, Penkhull, Fenton, Lane Delph, Foley, Lane End.—**III., On the Origin of the Art**, and its Practice among the early Nations.—**IV., Manufacture of Pottery**, prior to 1700.—**V., The Introduction of Red Porcelain** by Messrs. Elers, of Bradwell, 1690.—**VI., Progress of the Manufacture** from 1700 to Mr. Wedgwood's commencement in 1760.—**VII. Introduction of Fluid Glaze.**—Extension of the Manufacture of Cream Colour.—Mr. Wedgwood's Queen's Ware.—Jasper, and Appointment of Potter to Her Majesty.—Black Potting.—**VIII., Introduction of Porcelain.** Mr. W. Littler's Porcelain.—Mr. Cookworthy's Discovery of Kaolin and Petuntse, and Patent.—Sold to Mr. Champion—resold to the New Hall Com.—Extension of Term.—**IX., Blue Printed Pottery.** Mr. Turner, Mr. Spode (1), Mr. Baddeley, Mr. Spode (2), Messrs. Turner, Mr. Wood, Mr. Wilson, Mr. Minton.—Great Change in Patterns of Blue Printed.—**X., Introduction of Lustre Pottery.** Improvements in Pottery and Porcelain subsequent to 1800.

Press Opinions.

"There is much curious and useful information in the work, and the publishers have rendered the public a service in reissuing it."—*Burton Mail*.

"Copies of the original work are now of considerable value, and the facsimile reprint now issued cannot but prove of considerable interest to all interested in the great industry."—*Derby Mercury*.

A Reissue of

THE CHEMISTRY OF THE SEVERAL NATURAL AND ARTIFICIAL HETEROGENEOUS COMPOUNDS USED IN MANUFACTURING PORCELAIN, GLASS AND POTTERY. By SIMEON SHAW.

(Originally published in 1837.) 750 pp. 1900. Royal 8vo. Price 14s.; India and Colonies, 15s.; Other Countries, 16s. 6d.; strictly net.

Contents.

PART I., ANALYSIS AND MATERIALS.—**Chapters I., Introduction:** Laboratory and Apparatus; **Elements:** Combinative Potencies, Manipulative Processes for Analysis and Reagents, Pulverisation, Blow-pipe Analysis, Humid Analysis, Preparatory Manipulations, General Analytic Processes, Compounds Soluble in Water, Compounds Soluble only in Acids, Compounds (Mixed) Soluble in Water, Compounds (Mixed) Soluble in Acids, Compounds (Mixed) Insoluble, Particular Analytic Processes.—**II., Temperature:** Coal, Steam Heat for Printers' Stoves.—**III., Acids and Alkalies:** Boracic Acid, Muriatic Acid, Nitric Acid, Sulphuric Acid, Potash, Soda, Lithia, Calculation of Chemical Separations.—**IV., The Earths:** Alumina, Clays, Silica, Flint, Lime, Plaster of Paris, Magnesia, Barytes, Felspar, Grauen (or China Stone), China Clay, Chert.—**V., Metals:** Reciprocal Combinative Potencies of the Metals, Antimony, Arsenic, Chromium, Green Oxide, Cobalt, Chromic Acid, Humid Separation of Nickel from Cobalt, Arsenite of Cobalt, Copper, Gold, Iron, Lead, Manganese, Platinum, Silver, Tin, Zinc.

PART II., SYNTHESIS AND COMPOUNDS.—**Chapters I., Sketch of the Origin and Progress of the Art.**—**II., Science of Mixing:** Scientific Principles of the Manufacture, Combinative Potencies of the Earths.—**III., Bodies:** Porcelain—Hard, Porcelain—Fritted Bodies, Porcelain—Raw Bodies, Porcelain—Soft, Fritted Bodies, Raw Bodies, Stone Bodies, Ironstone, Dry Bodies, Chemical Utensils, Fritted Jasper, Fritted Pearl, Fritted Drab, Raw Chemical Utensils, Raw Stone, Raw Jasper, Raw Pearl, Raw Mortar, Raw Drab, Raw Brown, Raw Fawn, Raw Cane, Raw Red Porous, Raw Egyptian, Earthenware, Queen's Ware, Cream Colour, Blue and Fancy Printed, Dipped and Mocha, Chalky, Rings, Stills, etc.—**IV., Glazes:** Porcelain—Hard Fritted, Porcelain—Soft Fritted, Porcelain—Soft Raw, Cream Colour Porcelain, Blue Printed Porcelain, Fritted Glazes, Analysis of Fritt, Analysis of Glaze, Coloured Glazes, Dips, Smears and Washes; **Glasses:** Flint Glass, Coloured Glasses, Artificial Garnet, Artificial Emerald, Artificial Amethyst, Artificial Sapphire, Artificial Opal, Plate Glass, Crown Glass, Broad Glass, Bottle Glass, Phosphoric Glass, British Steel Glass, Glass-Staining and Painting, Engraving on Glass, Dr. Faraday's Experiments.—**V., Colours:** Colour Making, Fluxes or Solvents, Components of the Colours: **Reds, etc., from Gold**, Carmine or Rose Colour, Purple, Reds, etc., from Iron, Blues, Yellows, Greens, Blacks, White, Silver for Burnishing, Gold for Burnishing, Printer's Oil, Lustres.

PART III., TABLES OF THE CHARACTERISTICS OF CHEMICAL SUBSTANCES.

Press Opinions.

"... There is an excellent historical sketch of the origin and progress of the art of pottery which shows the intimate knowledge of classical as well as (the then) modern scientific literature possessed by the late Dr. Shaw."—*Glasgow Herald*.

"The historical sketch of the origin and progress of pottery is very interesting and instructive. The science of mixing is a problem of great importance, and the query how the natural products, alumina and silica can be compounded to form the best wares may be solved by the aid of chemistry instead of by guesses, as was formerly the case. This portion of the book may be most suggestive to the manufacturer, as also the chapters devoted to the subject of glazes, glasses and colours."—*Birmingham Post*.

"Messrs. Scott, Greenwood & Co. are doing their best to place before the pottery trades some really good books, and their spirited enterprise is worthy of encouragement, for the utility of technical literature bearing upon the practical side of potting goes without saying. . . . They are to be congratulated on their enterprise in republishing it."—*Staffordshire Sentinel*.

Paper Making.

THE DYEING OF PAPER PULP. A Practical Treatise for the use of Papermakers, Paperstainers, Students and others. By JULIUS ERFURT, Manager of a Paper Mill. Translated into English and Edited with Additions by JULIUS HÜBNER, F.C.S., Lecturer on Papermaking at the Manchester Municipal Technical School. With Illustrations and **157 patterns of paper dyed in the pulp.** Royal 8vo. 180 pp. 1901. Price 15s.; India and Colonies, 16s.; Other Countries, 20s.; strictly net. Limited edition.

Contents.

I., Behaviour of the Paper Fibres during the Process of Dyeing, Theory of the Mordant.—II., Colour Fixing Mediums (Mordants).—III., Influence of the Quality of the Water Used.—IV., Inorganic Colours.—V., Organic Colours.—VI., Practical Application of the Coal Tar Colours according to their Properties and their Behaviour towards the Different Paper Fibres.—VII., Dyed Patterns on Various Pulp Mixtures.—Dyeing to Shade.—Index.

Press Opinions.

"The book is one that is of value to every one connected with the colouring of paper."—*Paper Trade Journal*.

"The great feature of the volume is undoubtedly the series of actual patterns of dyed papers, 157 in all—twelve of which, made in England, have been added to the original German series. Detailed formulæ are given for the preparation of the pulp for each, and the tints of the samples practically form a key, by means of which the accuracy of the student's or practitioner's experiments can be tested. . . ."—*World's Paper Trade Review*.

Enamelling on Metal.

ENAMELS AND ENAMELLING. An Introduction to the Preparation and Application of all Kinds of Enamels for Technical and Artistic Purposes. For Enamel Makers, Workers in Gold and Silver, and Manufacturers of Objects of Art. By PAUL RANDAU. Translated from the German. With Sixteen Illustrations. 180 pp. 1900. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

I., Introduction.—II., Composition and Properties of Glass.—III., Raw Materials for the Manufacture of Enamels.—IV., Substances Added to Produce Opacity.—V., Fluxes.—VI., Pigments.—VII., Decolorising Agents.—VIII., Testing the Raw Materials with the Blow-pipe Flame.—IX., Subsidiary Materials.—X., Preparing the Materials for Enamel Making.—XI., Mixing the Materials.—XII., The Preparation of Technical Enamels, The Enamel Mass.—XIII., Appliances for Smelting the Enamel Mass.—XIV., Smelting the Charge.—XV., Composition of Enamel Masses.—XVI., Composition of Masses for Ground Enamels.—XVII., Composition of Cover Enamels.—XVIII., Preparing the Articles for Enamelling.—XIX., Applying the Enamel.—XX., Firing the Ground Enamel.—XXI., Applying and Firing the Cover Enamel or Glaze.—XXII., Repairing Defects in Enamelled Ware.—XXIII., Enamelling Articles of Sheet Metal.—XXIV., Decorating Enamelled Ware.—XXV., Specialties in Enamelling.—XXVI., Dial-plate Enamelling.—XXVII., Enamels for Artistic Purposes, Recipes for Enamels of Various Colours.—Index.

Press Opinions.

"Should prove of great service to all who are either engaged in or interested in the art of enamelling."—*Jewellers and Watchmakers' Trade Advertiser*.

"I must inform you that this is the best book ever I have come across on enamels, and it is worth double its cost."—J. MINCHIN, Jr., Porto, Portugal, 22nd July, 1900.

"This is a very useful and thoroughly practical treatise, and deals with every branch of the enameller's art."—*Invention*.

THE ART OF ENAMELLING ON METAL. By W. NORMAN BROWN. Twenty-eight Illustrations. Crown 8vo. 60 pp. 1900. Price 2s. 6d.; Abroad, 3s.; strictly net.

Contents.

Chapters I., History—Cloisonné—Champs Léve—Translucent Enamel—Surface Painted Enamels.—II., Cloisonné—Champs Léves—Translucent—Painted.—III., Painted Enamel—Apparatus—Furnaces and Muffles for Firing.—IV., The Copper Base or Plate—Planishing—Cloisons—Champ Léve Plates.—V., Enamels—Trituration—Washing—Coating a Plate with Enamel—Firing Ordinary Plaques for Painting—Designing—Squaring off.—VI., Designs for Cloisonné—Designs for Painted Enamels—Technical Processes—Brushes, etc.—Colours—Grisaille—Full-coloured Designs.

Press Opinion.

"The information conveyed in *The Art of Enamelling on Metal* is as complete as can be expected in a manual of ordinary length, and is quite ample in all respects to start students in a most interesting branch of decorative art."—*Hardware Metals and Machinery*.]

Books on Textile and Dyeing Subjects.

THE TECHNICAL TESTING OF YARNS AND TEXTILE FABRICS. With Reference to Official Specifications. Translated from the German of Dr. J. HERZFELD. Second Edition. Sixty-nine Illustrations. 200 pp. Demy 8vo. 1902. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Yarn Testing. III., Determining the Yarn Number.—IV., Testing the Length of Yarns.—V., Examination of the External Appearance of Yarn.—VI., Determining the Twist of Yarn and Twist.—VII., Determination of Tensile Strength and Elasticity.—VIII., Estimating the Percentage of Fat in Yarn.—IX., Determination of Moisture (Conditioning).—Appendix

Press Opinions.

"It would be well if our English manufacturers would avail themselves of this important addition to the extensive list of German publications which, by the spread of technical information, contribute in no small degree to the success, and sometimes to the supremacy, of Germany in almost every branch of textile manufacture."—*Manchester Courier*.

"This is probably the most exhaustive book published in English on the subject."—*Textile Recorder*.

"A careful study of this book enables one to say with certainty that it is a standard work on the subject."—*Glasgow Herald*.

"... For the first time all the data relating to both physical and chemical tests as used throughout the whole of the textile industry, so that not only the commercial and textile chemist, who has frequently to reply to questions on these matters, but also the practical manufacturer of textiles and his subordinates, whether in spinning, weaving, dyeing, and finishing, are catered for. . . . The book is profusely illustrated, and the subjects of these illustrations are clearly described."—*Textile Manufacturer*.

DECORATIVE AND FANCY TEXTILE FABRICS.

With Designs and Illustrations. By R. T. LORD. A Valuable Book for Manufacturers and Designers of Carpets, Damask, Dress and all Textile Fabrics. 200 pp. 1898. Demy 8vo. 132 Designs and Illustrations. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., A Few Hints on Designing Ornamental Textile Fabrics.—II., A Few Hints on Designing Ornamental Textile Fabrics (continued).—III., A Few Hints on Designing Ornamental Textile Fabrics (continued).—IV., A Few Hints on Designing Ornamental Textile Fabrics (continued).—V., Hints for Ruled-paper Draughtsmen.—VI., The Jacquard Machine.—VII., Brussels and Wilton Carpets.—VIII., Tapestry Carpets.—IX., Ingrain Carpets.—X., Axminster Carpets.—XI., Damask and Tapestry Fabrics.—XII., Scarf Silks and Ribbons.—XIII., Silk Handkerchiefs.—XIV., Dress Fabrics.—XV., Mantle Cloths.—XVI., Figured Plush.—XVII., Bed Quilts.—XVIII., Calico Printing.

Press Opinions.

"The book can be strongly recommended to students and practical men."—*Textile Colourist*.

"Those engaged in the designing of dress, mantle tapestry, carpet and other ornamental textiles will find this volume a useful work of reference."—*Leeds Mercury*.

"To be commended as a model manual."—*Dundee Advertiser*.

"Designers especially, who desire to make progress in their calling, will do well to take the hints thrown out in the first four chapters on 'Designing Ornamental Textile Fabrics'."—*Nottingham Daily Guardian*.

POWER-LOOM WEAVING AND YARN NUMBERING,

According to Various Systems, with Conversion Tables. An Auxiliary and Text-book for Pupils of Weaving Schools, as well as for Self-Instruction and for General Use by those engaged in the Weaving Industry. Translated from the German of ANTHON GRUNER. With **Twenty-six Diagrams in Colours.** 150 pp. 1900. Crown 8vo. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

I., Power-Loom Weaving in General. Various Systems of Looms.—II., Mounting and Starting the Power-Loom. English Looms.—Tappet or Treadle Looms. Dobbies.—III., General Remarks on the Numbering, Reeling and Packing of Yarn.—Appendix.—Useful Hints. Calculating Warps.—Wett Calculations.—Calculations of Cost Price in Hanks.

Press Opinions.

"A long-felt want in the weaving industry,"—*Belfast Evening Telegraph*.
 "The author has dealt very practically with the subject,"—*Bradford Daily Telegraph*.
 "The book . . . should prove invaluable to the student,"—*Cotton Factory Times*.
 "It is a capital text-book for use in the weaving schools or for self-instruction, while all engaged in the weaving industry will find its suggestions helpful,"—*Northern Daily Telegraph*.
 "Yarn numbering according to various systems, with conversion tables and numerous coloured diagrams, materially assist to a clear comprehension of the subject,"—*Northern White*.
 "The 'inside' managers of our textile mills in which the work is complex or greatly varied, and where yarns of different materials are in use, will find this work convenient for reference,"—*Textile Mercury*.
 "The author attempts to fill a gap in weaving literature caused by the neglect of many obscure points connected with the industry,"—*Cheshire County News*.
 "It is clear and concise, and gives just that knowledge in quality and amount which any student of the weaving industry ought to consider as a minimum necessary for his thorough comprehension of his future profession,"—*North British Daily Mail*.
 "... The work should prove of much value, as it is in every sense practical, and is put before the reader in such a clear manner that it can be easily understood,"—*Textile Industries*.
 "The smallest details of loom-setting are entered into, and a full explanation of problems, which are a source of anxiety to many engaged in overlooking, is given. Students will find the work an admirable text-book, and all who are interested in weaving will see in it a valuable addition to the literature on this subject,"—*Bradford Observer*.

THE CHEMICAL TECHNOLOGY OF TEXTILE

FIBRES: Their Origin, Structure, Preparation, Washing, Bleaching, Dyeing, Printing and Dressing. By Dr. GEORG VON GEORGIEVICS. Translated from the German by CHARLES SALTER. 320 pp. Forty-seven illustrations. Royal 8vo. 1902. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s. net.

Contents.

Chapters I., **The Textile Fibres.**—Artificial Fibres—Mineral Fibres—Vegetable Fibres—Cellulose—Cotton—Bombax Cotton—Vegetable Silk—Flax—Hemp—Jute—Ramie, Rhea, China Grass, Nettle Fibre—Distinguishing Tests for the Various Fibres—Animal Fibres: Silk—Animal Hairs—Sheep's Wool—Goat Wool and Camel Wool—Artificial Wool (Wool Substitutes)—Conditioning. —II., **Washing, Bleaching, Carbonising.** Washing and Bleaching (Definition)—Bleaching Agents—Cotton Bleaching—Linen Bleaching—Jute Bleaching—Hemp Bleaching—Ramie Bleaching—Scouring and Bleaching Silk—Washing and Bleaching Wool—Blueing or White Dyeing—Carbonising. —III., **Mordants and Mordanting.** Mordants: Mordanting Wool—Mordanting Silk—Mordanting Cotton—Alumina Mordants—Iron Mordants—Chrome Mordants—Tin Mordants—Copper and other Mordants—The Fixing Agents (Acid Mordants): Tannic Acids—Oleic Acids.—IV., **Dyeing.** (1) Theory of Colour—Combination of Colours: Dyeing to Pattern—(2) Theory of Dyeing—(3) Classification of Dye Stuffs—Methods of Dyeing—Application of Acid Dye Stuffs—Application of Basic Dye Stuffs—Application of Direct or Substantive Cotton Dyes—Application of the Mordant Dyes: Dyeing with Cochineal—Black and Blue Dyeings with Logwood on Wool—Turkey-Red Dyeing—Dyeing with Catechu—Black-Dyeing Cotton with Logwood—Application of the Vat Dyes—Application of the Developing Dyes—(4) Dyeing on a Manufacturing Scale—Selection of Dye Stuffs for Dyeing—Silk Dyeing—Wool Dyeing—Cotton Dyeing—Dyeing Mixed Fabrics—(5) Sample Dyeings, Colorimetric Determinations, Reactions of Dye Stuffs on the Fibre, Tests for Fastness. V., **Printing.**—Hand Printing—The Perrotine Press—The Cylinder Press—Calico Printing: (1) Reproduction of Pattern by Direct Printing: Thickening Agents—Employment of Mordant Dye Stuffs, Basic, Albumin, Direct, Developing, Vat, Acid—Treatment of the Goods when Printed—(2) Combined Printing and Dyeing—(3) Discharge Style Printing: Discharging the Mordant—Discharging Antimony Tannate—Discharging the Finished Dye—Turkey-Red Discharge Style—(4) Reserve Style Printing—(5) Topping Printing—Wool Printing—Silk Printing—Printing Yarns, Warps, and Combed Sliver.—VI., **Dressing and Finishing.**—Dressing and Finishing Substances used in Finishing: (1) Starch, Gum, etc.—(2) Fatty Substances—(3) Hygroscopic Materials—(4) Loading Ingredients—(5) Colouring for the Dressing Preparations—(6) Metals or their Sulphites—(7) Waterproofing—(8) Fireproofing—(9) Antiseptics for Prevention of Mould—Application of Dressings—Drying—Stretching—Finishing: Shearing, Damping, Calendering, Beetling, Moiré or Watered Effects, Stamping—Finishing Woollens.—Index.

COLOUR: A HANDBOOK OF THE THEORY OF COLOUR. By GEORGE H. HURST, F.C.S. **With Ten Coloured Plates** and Seventy-two Illustrations. 160 pp. Demy 8vo. 1900. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., **Colour and Its Production.** Light, Colour, Dispersion of White Light, Methods of Producing the Spectrum, Glass Prism and Diffraction Grating Spectroscopes, The Spectrum, Wave Motion of Light, Recomposition of White Light, Hue, Luminosity, Purity of Colours, The Polaroscope, Phosphorescence, Fluorescence, Interference. II., **Cause of Colour in Coloured Bodies.** Transmitted Colours, Absorption Spectra of Colouring Matters. III., **Colour Phenomena and Theories.** Mixing Colours, White Light from Coloured Lights, Effect of Coloured Light on Colours, Complementary Colours, Young-Helmholtz Theory, Brewster Theory, Supplementary Colours, Maxwell's Theory, Colour Photography. IV., **The Physiology of Light.** Structure of the Eye, Persistence of Vision, Subjective Colour Phenomena, Colour Blindness. V., **Contrast.** Contrast, Simultaneous Contrast, Successive Contrast, Contrast of Tone, Contrast of Colours, Modification of Colours by Contrast, Colour Contrast in Decorative Design. VI., **Colour in Decoration and Design.** Colour Harmonies, Colour Equivalents, Illumination and Colour, Colour and Textile Fabrics, Surface Structure and Colour. VII., **Measurement of Colour.** Colour Patch Method, The Tintometer, Chromometer.

Press Opinions.

"This useful little book possesses considerable merit."—*Birmingham Post*.

"It will be found to be of direct service to the majority of dyers, calico printers and colour mixers, to whom we confidently recommend it."—*Chemical Trade Journal*.

"It is thoroughly practical, and gives in simple language the why and wherefore of the many colour phenomena which perplex the dyer and the colourist."—*Dyer and Calico Printer*.

TEXTILE RAW MATERIALS AND THEIR CONVERSION INTO YARNS. (The Study of the Raw Materials and the Technology of the Spinning Process.) Text-book for Textile, Trade and Higher Technical Schools. By JULIUS ZIPSER. Translated from German by CHARLES SALTER. 302 Illustrations. 480 pp. Demy 8vo. 1901. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Introduction.

PART I.—The Raw Materials Used in the Textile Industry.

GROUP I. MINERAL RAW MATERIALS. Asbestos—Glass—Metals.

GROUP II. VEGETABLE RAW MATERIALS. Seed Fibres—Cotton—Bombax Wool—Asclepias Wool—Poplar, Cotton Grass and Bulrush Wool—Stem Fibres. Flax—Hemp—Jute—Nettle Fibres—Sunn Hemp—Leaf Fibres—New Zealand Hemp—Manila Hemp—Sisal and Domingo Hemp—Aloe Fibre—Pineapple Fibre—Vegetable Wool—Fruit Fibres—Cocoanut Fibre—Other Vegetable Raw Materials employed in Weaving—Straw—Wood—Cane—Cacoutchouc.

GROUP III. ANIMAL RAW MATERIALS. Animal Wool and Hair—Sheep's Wool—Goat Wool—Camel Wool—Llama and similar Wools—Cow Hair—Poodle Hair—Hare, Rabbit, Cat and Beaver Fur—Horse Hair—Bristles—Silk—Natural Silk—Artificial Silk—Byssus Silk—Detection and Estimation of Textile Raw Materials in Yarns and Fabrics—Characteristics of Mineral Raw Materials—Detecting and Methods of separating Vegetable and Animal Raw Materials in general—The Combustion Test—The Mandarin Test—The Picric Acid Test—The Sulphuric Acid Test—The Alkali Test—Differentiation of Animal and Vegetable Fibres in detail—Characteristics of Cotton—Characteristics of Flax—Characteristics of Sheep's Wool—Characteristics of True Silk—Characteristics of Artificial Silk—Determining the Constituents of Textile Fabrics—The Detection of Cotton in Linen Fabrics—The Detection of Cotton in Woollen Fabrics—The Detection of Cotton in Silk Fabrics—The Detection of Wool in Silk Fabrics—The Detection of Wild Silk in True Silk Fabrics—The Detection of Artificial Silk in Silk Fabrics—The Detection of Byssus Silk in Silk Fabrics.

PART II.—The Technology of Spinning or the Conversion of Textile Raw Materials into Yarn.

Spinning—Sequence of Operations in Spinning—General Observations on the Machinery and Process of Spinning—Fine Spinning Machines—Requisite Properties for a Well-spun Thread.

GENERAL REVIEW OF THE VARIOUS BRANCHES OF THE SPINNING PROCESS.

GROUP I. SPINNING VEGETABLE RAW MATERIALS. Cotton Spinning—Preparatory Processes—Mixing—Opening and Cleaning—Carding—Combing—Drawing and Doubling—Roving—Roving Frames producing a Permanent Twist—Machines producing Temporary Twist—Fine Spinning—The Throstle or Water Frame—The Mule Frame—Supplementary Treatment: Finishing—Finishing Processes producing no change in the Character of the Yarn—Converting Yarns into New Forms—Additional Finishing Processes—Packing—Installation of a Cotton Mill—Humidifiers—Spinning Waste Cotton and Waste Cotton Yarns—Preliminary Operations

—Carding—Slubbing—Fine Spinning—Supplementary and Finishing Operations—Flax Spinning—Preparatory Treatment—Gilling—Doubling and Drawing—Roving—The Roving Frame—Fine Spinning—Supplementary Treatment—Operations leaving the Yarn unchanged—Operations for producing New Combinations of Threads—Packing—Tow Spinning—Preparatory Treatment—Roving—Fine Spinning—Hemp Spinning—Spinning Hackled or Line Hemp—Spinning Hemp Tow String—Jute Spinning—Spinning Jute Line Yarn—Spinning Jute Tow Yarns—Supplementary Treatment—The Production of Mixed Yarns—Utilising Jute Waste.

PART III.—Spinning Animal Raw Materials.

Spinning Carded Woolen Yarn—Preparatory Treatment—Scouring, Washing and Rinsing—Draining and Drying the Wool—Barr Extraction—Opening and Willowing—Oiling the Wool—Carding—Condensing—Belt Condensers—Steel Band Condenser—Fine Spinning—The Mule—The Throstle Frame—Supplementary Treatment—Supplementary Treatment leaving the Yarn unaltered—Treatment with a view to producing Novel Effects—Finishing Yarn—Packing Yarn—Worsted Spinning—Manufacture of True Worsted Yarn—Preliminary Treatment—Sorting—Steeping, Washing and Rinsing—Drying—Oiling—Carding—Gilling—Combing—Drawing—Smoothing—Drawing and Slubbing—Roving—Fine Spinning—The Worsted Mule Frame—Worsted Throstle Frames—Supplementary Treatment—Processes leaving the Yarn unchanged—Producing New Types of Yarn—Finishing Worsted Yarn—Packing—Semi-Worsted Yarns—Artificial Wool or Shoddy Spinning—Shoddy and Mungo Manufacture—Cleaning the Rags—Sorting the Rags—Trimming the Rags—Scouring—Disintegration—Extract Wool—Shaking—Scouring—Carbonising—Disintegration—Spinning Shoddy and other Wool Substitutes—Spinning Waste Silk—Chappe Silk—Preliminary Treatment—Sorting—Steeping—Maceration—Scouring and Beetling—Drying—Sprinkling—Beating—Opening—Dressing—Spreading—Doubling and Drawing—Slubbing and Roving—Fine Spinning—Supplementary Operations—Operations leaving the Yarn unaltered—Producing New Forms of Yarn—Finishing off Chappe Silk—Packing—Bourette Spinning—Index.

DYERS' MATERIALS: An Introduction to the Examination, Evaluation and Application of the most important Substances used in Dyeing, Printing, Bleaching and Finishing. By PAUL HEERMAN, Ph.D. Translated from the German by ARTHUR C. WRIGHT, M.A. (Oxon.), B.Sc. (Lond.). With Two Plates, containing Twenty-four Illustrations. Crown 8vo. 150 pp. 1901. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s; strictly net.

Contents.

Chapter I., General: Table I., *Indicators—Standard Solutions*.—Table II., *Solutions and Reagents in General Use*.—Chapter II., *Primary Materials: Water—Textile Fibres*. Chapter III., *Inorganic Materials: Hydrochloric Acid—Chlorides of Sodium, Magnesium, Ammonium, Barium, Zinc, Copper, Manganese and Aluminium, Stannous Chloride, Stannic Chloride, Tin Spirits, Chromium Chloride—Fluorides and Bifluorides, Alkaline Bifluorides, Chromium Fluoride, Chromium Oxylfluoride, Copper Fluoride, Antimony Fluoride, Aniline Hydrofluoride, Antimony Fluoride, Double Salts—Sulphuric Acid, Fuming or Nordhausen Sulphuric Acid—Sulphates, Sodium Sulphate, Sodium Bisulphate, Calcium Sulphate, Magnesium Sulphate, Lead Sulphate, Ferrous Sulphate, Aluminium Sulphate, Ferric Sulphate, Copper Sulphate, Alums—Nitric Acid and Nitrates: Nitric Acid, Sodium, Silver, Lead, Ferrous, Ferric and Rarer Nitrates—Chlorine-Oxygen Compounds: Bleaching Powder, Alkaline Hypochlorites, other Hypochlorites, Potassium, Sodium, Aluminium, Chromium and Aniline Chlorates—Sulphite Compounds: Sulphurous Acid, Sulphites, Sodium Bisulphite, Hydro- or Hypo-sulphurous Acid, Hydro- or Hypo-sulphites, Sodium Thiosulphate—Miscellaneous Compounds: Sodium Nitrite, Sodium Phosphate, Water-glass, Sodium Arsenate, Sodium Tungstate, Sodium Stannate, Sodium Aluminate, Borax, Potassium Permanganate, Potassium Bichromate, Sodium Bichromate, Vanadates—Alkalies: Ammonia, Ammonium Salts, Caustic Soda, Caustic Potash, Sodium Carbonate, Calcium Carbonate—Peroxides: Hydrogen Peroxide, Barium Peroxide, Sodium Peroxide—Zinc Dust*.—Chapter IV., *Organic Compounds: Fatty Acids and Their Salts—Acetic Acid, Acetates, Ferrous Acetate, Oxalic Acid and Oxalates, Tartaric Acid, Tartar, Tartar Emetic, Citric Acid, Lactic Acid—Cyanogen Compounds: Sulphocyanides, Potassium Ferrocyanide, Potassium Ferricyanide—Derivatives of the Iats: Soap, "Boiled-off Liquor," Turkey Red Oil—Tannins—Aniline and Aniline Salts—Thickening and Stiffening Materials: Starch, Prepared and Soluble Starch, Dextrine, Gum Arabic, Gum Senegal, Gum Tragacanth, Glue, Size—Dyes*.—Appendix: Atomic Weights of the Elements—Molecular Weights of Certain Compounds—Gravimetric Equivalents—Volumetric Equivalents. Plate I., *Microscopic Appearance of the Textile Fibres* (11 Illustrations).—Plate II., *Microscopic Appearance of the Different Varieties of Starch* (13 Illustrations).—Index.

Press Opinions.

"To those engaged in any branches of dyeing, printing, bleaching and finishing it ought to prove a valuable addition to existing works."—*Textile Industries*.

"Cannot fail to be of the greatest value."—*Huddersfield Examiner*.

THE COLOUR PRINTING OF CARPET YARNS. A Useful Manual for Colour Chemists and Textile Printers. By DAVID PATERSON, F.C.S. Seventeen Illustrations. 132 pp. Demy 8vo. 1900. Price 7s. 6d.; India and Colonies, 8s. Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., Structure and Constitution of Wool Fibre.—II., Yarn Scouring.—III., Scouring Materials.—IV., Water for Scouring.—V., Bleaching Carpet Yarns.—VI., Colour Making for Yarn Printing.—VII., Colour Printing Pastes.—VIII., Colour Recipes for Yarn Printing.—IX., Science of Colour Mixing.—X., Matching of Colours.—XI., "Hank" Printing.—XII., Printing Tapestry Carpet Yarns.—XIII., Yarn Printing.—XIV., Steaming Printed Yarns.—XV., Washing of Steamed Yarns.—XVI., Aniline Colours Suitable for Yarn Printing.—XVII., Glossary of Dyes and Dye-wares used in Wood Yarn Printing.—Appendix.

Press Opinions.

- "The book is worthy the attention of the trade."—*Worcester Herald*.
 "The treatise is arranged with great care, and follows the processes described in a manner at once clear and convincing."—*Glasgow Record*.
 "A most useful manual dealing in an intelligible and interesting manner with the colour printing of carpet yarns."—*Kidderminster Times*.
 "The author has evidently strained every effort in order to make his work the standard guide of its class."—*Leicester Post*.
 "The book, which is admirably printed and illustrated, should fulfil the need of a practical guide in the colour printing of carpet yarns."—*Nottingham Express*.
 "The subject is very exhaustively treated in all its branches. . . ."—*Northern Whig*.
 "It gives an account of its subject which is both valuable and instructive."—*Scotsman*.
 "The work shows a thorough grasp of the leading characteristics as well as the minutæ of the industry, and gives a lucid description of its chief departments. . . . As a text-book in technical schools where this branch of industrial education is taught, the book is valuable."—*Dundee Courier*.
 "The book bears every mark of an extensive practical knowledge of the subject in all its bearings, and supplies a real want in technical literature."—*Dyer and Calico Printer*.
 "It is thoroughly practical, and contains much information which has not hitherto appeared in book form."—*Journal of the Society of Dyers and Colourists*.

A PRACTICAL TREATISE ON THE BLEACHING OF LINEN AND COTTON YARN AND FABRICS. By

L. TAILFER, Chemical and Mechanical Engineer. Translated from the French by JOHN GEDDES MCINTOSH, Lecturer on Chemical Technology, London. Demy 8vo. 303 pp. Twenty Illusts. 1901. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

Chapter I. General Considerations on Bleaching. Chapter II. Steeping. Chapter III. Washing: Its End and Importance—Roller Washing Machines—Wash Wheel (Dash Wheel)—Stocks or Wash Mill—Squeezing. Chapter IV. Lye Boiling—Lye Boiling with Milk of Lime—Lye Boiling with Soda Lye—Description of Lye Boiling Keirs—Operations of Lye Boiling—Concentration of Lyes. Chapter V. Maier and Platt's Keir—Description of the Keir—Saturation of the Fabrics—Alkali used in Lye Boiling—Examples of Processes. Chapter VI. Soap—Action of Soap in Bleaching—Quality and Quantity of Soaps to use in the Lye—Soap Lyes or Scalds—Soap Scouring Stocks. Chapter VII. Bleaching on Grass or on the Bleaching Green or Lawn. Chapter VIII. Chemicking—Remarks on Chlorides and their Decolourising Action—Chemicking Cisterns—Chemicking—Strengths, etc. Chapter IX. Sours—Properties of the Acids—Effects Produced by Acids Souring Cisterns. Chapter X. Drying—Drying by Steam—Drying by Hot Air—Drying by Air. Chapter XI. Damages to Fabrics in Bleaching—Yarn Mildew—Fermentation—Iron Rust Spots—Spots from Contact with Wood—Spots incurred on the Bleaching Green—Damages arising from the Machines. Chapter XII. Examples of Methods used in Bleaching—Linen—Cotton. Chapter XIII. The Valuation of Caustic and Carbonated Alkali (Soda) and General Information Regarding these Bodies—Object of Alkalimetry—Titration of Carbonate of Soda—Comparative Table of Different Degrees of Alkalimetric Strength—Five Problems relative to Carbonate of Soda—Caustic Soda, its Properties and Uses—Mixtures of Carbonated and Caustic Alkali—Note on a Process of Manufacturing Caustic Soda and Mixtures of Caustic and Carbonated Alkali (Soda). Chapter XIV. Chlorimetry—Titration—Wagner's Chlorimetric Method—Preparation of Standard Solutions—Apparatus for Chlorine Valuation—Alkali in Excess in Decolourising Chlorides. Chapter XV. Chlorine and Decolourising Chlorides—Synopsis—Chlorine—Chloride of Lime—Hypochlorite of Soda—Brocholi's Chlorozone—Various Decolourising Hypochlorites—Comparison of Chloride of Lime and Hypochlorite of Soda. Chapter XVI. Water—Qualities of Water—Hardness—Dervaux's Purifier—Testing the Purified Water—Different Plant for Purification—Filters. Chapter XVII. Bleaching of Yarn—Weight of Yarn—Lye Boiling—Chemicking—Washing—Bleaching of Cotton Yarn. Chapter XVIII. The Installation of a Bleach Works—Water Supply—Steam Boilers—Steam Distribution Pipes—Engines—Keirs—Washing Machines—Stocks—Wash Wheels—Chemicking and Souring Cisterns—Various—Buildings. Chapter XIX. Addenda—Energy of Decolourising Chlorides and Bleaching by Electricity and Ozone—Energy of Decolourising Chlorides—Chlorides—Production of Chlorine and Hypochlorites by Electrolysis—Lunge's Process for increasing the intensity of the Bleaching Power of Chloride of Lime—Triller's Process for Removing the Excess of Lime or Soda from Decolourising Chlorides—Bleaching by Ozone.

THE SCIENCE OF COLOUR MIXING. A Manual intended for the use of Dyers, Calico Printers and Colour Chemists. By DAVID PATERSON, F.C.S. Forty-one Illustrations. **Five Coloured Plates, and Four Plates showing Eleven Dyed Specimens of Fabrics.** 132 pp. Demy 8vo. 1900. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., Colour a Sensation; Colours of Illuminated Bodies; Colours of Opaque and Transparent Bodies; Surface Colour.—II., Analysis of Light; Spectrum; Homogeneous Colours: Ready Method of Obtaining a Spectrum.—III., Examination of Solar Spectrum; The Spectroscope and its Construction. Chemists' Use of the Spectroscope.—IV., Colour by Absorption; Solutions and Dyed Fabrics; Dichroic Coloured Fabrics in Gaslight.—V., Colour Primaries of the Scientist *versus* the Dyer and Artist; Colour Mixing by Rotation and Lye Dyeing; Hue, Purity, Brightness; Tints; Shades, Scales, Tones, Sad and Sombre Colours.—VI., Colour Mixing; Pure and Impure Greens, Orange and Violets; Large Variety of Shades from few Colours; Consideration of the Practical Primaries: Red, Yellow and Blue.—VII., Secondary Colours; Nomenclature of Violet and Purple Group; Tints and Shades of Violet; Changes in Artificial Light.—VIII., Tertiary Shades; Broken Hues; Absorption Spectra of Tertiary Shades.—Appendix: Four Plates with Dyed Specimens Illustrating Text.—Index.

Press Opinions.

"The work has evidently been prepared with great care."—*Halifax Courier*.

"The volume, which is clearly and popularly written, should prove of the utmost service to all who are concerned with the practical use of colours, whether as dyers or painters."—*Scotsman*.

"We have no hesitation in advising the purchase of the present volume by dyers and calico printers, as containing a mass of most useful information at a nominal price."—*Irish Textile Journal*.

"Mr. Paterson's work . . . will be found exceedingly helpful, not only to the practical colourist, but also to students in our textile colleges, by forming a useful complement to their class lectures."—*Wakefield Express*.

" . . . The author is a dyer, and in his concluding chapters keeps well before him the special wants and requirements of dyers. He writes pleasantly and lucidly, and there is no difficulty in following him, although here and there a lapse into ambiguousness occurs."—*Textile Mercury*.

COLOUR MATCHING ON TEXTILES. A Manual intended for the use of Students of Colour Chemistry, Dyeing and Textile Printing. By DAVID PATERSON, F.C.S. Coloured Frontispiece. Twenty-nine Illustrations and Fourteen Specimens of Dyed Fabrics Illustrating Text. Demy 8vo. 132 pp. 1901. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., Colour Vision and Structure of the Eye—Perception of Colour—Primary and Complementary Colour Sensations.—II., Daylight for Colour Matching—Selection of a Good Pure Light—Diffused Daylight, Direct Sunlight, Blue Skylight, Variability of Daylight, etc., etc.—III., Matching of Hues—Purity and Luminosity of Colours—Matching Bright Hues—Aid of Tinted Films—Matching Difficulties Arising from Contrast.—IV., Examination of Colours by Reflected and Transmitted Lights—Effect of Lustre and Transparency of Fibres in Colour Matching.—V., Matching of Colours on Velvet Pile—Optical Properties of Dye-stuffs, Dichroism, Fluorescence.—VI., Use of Tinted Mediums—Orange Film—Defects of the Eye—Yellowing of the Lens—Colour Blindness, etc.—VII., Matching of Dyed Silk Trimmings and Linings and Bindings—Its Difficulties—Behaviour of Shades in Artificial Light—Colour Matching of Old Fabrics, etc.—VIII., Examination of Dyed Colours under the Artificial Lights—Electric Arc, Magnesium and Dufton, Gardner Lights, Welsbach, Acetylene, etc.—Testing Qualities of an Illuminant.—IX., Influence of the Absorption Spectrum in Changes of Hue under the Artificial Lights—Study of the Causes of Abnormal Modifications of Hue, etc.

Press Opinions.

"It should form a part of the library of every dyer and colourist in the United Kingdom and indeed of every English-speaking country."—*Dyer and Calico Printer*.

"We recommend it to every one who has anything to do with colour matching, even to merchants dealing in colouring goods."—*Indian Textile Journal*.

Reissue of

THE ART OF DYEING WOOL, SILK AND COTTON.

Translated from the French of M. HELLOT, M. MACQUER and M. LE PILEUR D'APLIGNY. First Published in English in 1789. Six Plates. Demy 8vo. 446 pp. 1901. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Part I., **The Art of Dyeing Wool and Woollen Cloth, Stuffs, Yarn, Worsted, etc.**

Part II., **The Art of Dyeing Silk.**

Part III., **The Art of Dyeing Cotton and Linen Thread, together with the Method of Stamping Silks, Cottons, etc.**

Press Opinions.

"The book has been produced in excellent style and should be of great assistance to dyers."
—*Drapers' Record*.

"Its reissue cannot fail to be of deep interest to all engaged in textile manufacture."
—*Macclesfield Courier*.

THE DYEING OF COTTON FABRICS: A Practical Handbook for the Dyer and Student. By FRANKLIN BEECH, Practical Colourist and Chemist. 272 pp. Forty-four Illustrations of Bleaching and Dyeing Machinery. Demy 8vo. 1901. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Chapters I., Structure and Chemistry of the Cotton Fibre.—II., Scouring and Bleaching of Cotton.—III., Dyeing Machinery and Dyeing Manipulations.—IV., Principals and Practice of Cotton Dyeing—1, Direct Dyeing; 2, Direct Dyeing followed by Fixation with Metallic Salts; 3, Direct Dyeing followed by Fixation with Developers; 4, Direct Dyeing followed by Fixation with Couplers; 5, Dyeing on Tannic Mordant; 6, Dyeing on Metallic Mordant; 7, Production of Colour Direct upon Cotton Fibres; 8, Dyeing Cotton by Impregnation with Dye-stuff Solution.—V., Dyeing Union (Mixed Cotton and Wool) Fabrics.—VI., Dyeing Half Silk (Cotton-Silk, Satin) Fabrics.—VII., Operations following Dyeing—Washing, Soaping, Drying.—VIII., Testing of the Colour of Dyed Fabrics.—IX., Experimental Dyeing and Comparative Dye Testing.—Index.

The book contains numerous recipes for the production on Cotton Fabrics of all kinds of a great range of colours, thus making it of great service in the Dyehouse, while to the Student it is of value in that the scientific principles which underlie the operations of dyeing are clearly laid down.

THE DYEING OF WOOLLEN FABRICS. By FRANKLIN BEECH, Practical Colourist and Chemist. Thirty-three Illustrations. Demy 8vo. 228 pp. 1902. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d. net.

Contents.

Chapters I., The Wool Fibre—Structure, Composition and Properties.—II., Processes Preparatory to Dyeing—Scouring and Bleaching of Wool.—III., Dyeing Machinery and Dyeing Manipulations—Loose Wool Dyeing, Yarn Dyeing and Piece Dyeing Machinery.—IV., The Principles and Practice of Wool Dyeing—Properties of Wool Dyeing—Methods of Wool Dyeing—Groups of Dyes—Dyeing with the Direct Dyes—Dyeing with Basic Dyes—Dyeing with Acid Dyes—Dyeing with Mordant Dyes—Level Dyeing—Blacks on Wool—Reds on Wool—Mordanting of Wool—Orange Shades on Wool—Yellow Shades on Wool—Green Shades on Wool—Blue Shades on Wool—Violet Shades on Wool—Brown Shades on Wool—Mode Colours on Wool.—V., Dyeing Union (Mixed Cotton Wool) Fabrics.—VI., Dyeing of Gloria.—VII., Operations following Dyeing—Washing, Soaping, Drying.—VIII., Experimental Dyeing and Comparative Dye Testing.—IX., Testing of the Colour of Dyed Fabrics.—Index.

COTTON SPINNING (First Year). By THOMAS THORNLEY, Spinning Master, Bolton Technical School. 160 pp. Eighty-four Illustrations. Crown 8vo. 1901. Price 3s.; Abroad, 3s. 6d.; strictly net.

Contents.

Syllabus and Examination Papers of the City and Guilds of London Institute.—Chapters I., Cultivation, Classification, Ginning, Baling and Mixing of the Raw Cotton.—II., Bale-Breakers, Mixing Lattices and Hopper Feeders.—III., Opening and Scutching.—IV., Carding.—Index to Illustrations.—General Index.

COTTON SPINNING (Intermediate, or 'Second Year). By THOMAS THORNLEY. 180 pp. Seventy Illustrations. Crown 8vo. 1901. Price 5s.; India and British Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Syllabuses and Examination Papers of the City and Guilds of London Institute.—Chapters I., The Combing Process.—II., The Drawing Frame.—III., Bobbin and Fly Frames.—IV., Mule Spinning.—V., Ring Spinning.—Index to Illustrations.—General Index.

COTTON SPINNING (Honours, or Third Year). By THOMAS THORNLEY. 216 pp. Seventy-four Illustrations. Crown 8vo. 1901. Price 5s.; India and British Colonies, 5s. 6d.; Other Countries, 6s.; strictly net.

Contents.

Syllabuses and Examination Papers of the City and Guilds of London Institute.—Chapters I., Cotton.—II., The Practical Manipulation of Cotton Spinning Machinery.—III., Doubling and Winding.—IV., Reeling.—V., Warping.—VI., Production and Costs.—VII., Main Driving.—VIII., Arrangement of Machinery and Mill Planning.—IX., Waste and Waste Spinning.—Index to Illustrations.—General Index.

Opinions of Spinning Teachers.

"The work (Vol. I.) contains a large amount of valuable information."—Mr. Jas. Tasher, Preston.

"They are certainly the best published on the subject."—Mr. John Kerfoot, Leigh.

"Admirably fulfils the object in view, viz., a concise guide to the students preparing for the City and Guilds Examination Course."—Mr. Jas. W. Lomax, Bolton.

"I have carefully read the book, and do not hesitate in saying that I consider it will undoubtedly be a boon to cotton spinning students for three, among other, reasons: (1) The store of information on different makers' machines; (2) it shows the student how he should consider the questions proposed at the examinations; and (3) the methods he should adopt in answering same."—Samuel Ward, Teacher in Cotton Spinning, Glossop and Openshaw.

COTTON COMBING MACHINES. By THOS. THORNLEY, Spinning Master. Technical School. Bolton. Demy 8vo. 117 Illustrations. 300 pp. 1902. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d. net.

Contents.

Chapters I., The Sliver Lap Machine and the Ribbon Cap Machine.—II., General Description of the Heilmann Comber.—III., The Cam Shaft.—IV., On the Detaching and Attaching Mechanism of the Comber.—V., Resetting of Combers.—VI., The Erection of a Heilmann Comber.—VII., Stop Motions: Various Calculations.—VIII., Various Notes and Discussions.—IX., Cotton Combing Machines of Continental Make.—Index.

Books for Mining Engineers and Steam Users.

RECOVERY WORK AFTER PIT FIRES. A Description of the Principal Methods Pursued, especially in Fiery Mines, and of the Various Appliances Employed, such as Respiratory and Rescue Apparatus, Dams, etc. By ROBERT LAMPRECHT, Mining Engineer and Manager. Translated from the German. Illustrated by Six large Plates, containing Seventy-six Illustrations. 175 pp., demy 8vo. 1901. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s.; strictly net.

Contents.

Preface.—I., **Causes of Pit Fires:** 1, Fires Resulting from the Spontaneous Ignition of Coal; 2, Fires Caused by Burning Timber; 3, Fires Caused by Fire-damp Explosions.—II., **Preventive Regulations:** 1, The Outbreak and Rapid Extension of a Shaft Fire can be most reliably prevented by Employing little or no Combustible Material in the Construction of the Shaft; 2, Precautions for Rapidly Localising an Outbreak of Fire in the Shaft; 3, Precautions to be Adopted in case those under 1 and 2 Fail or Prove Inefficient. Precautions against Spontaneous Ignition of Coal. Precautions for Preventing Explosions of Fire-damp and Coal Dust. Employment of Electricity in Mining, particularly in Fiery Pits. Experiments on the Ignition of Fire-damp Mixtures and Clouds of Coal Dust by Electricity.—III., **Indications of an Existing or Incipient Fire.**—IV., **Appliances for Working in Irrespirable Gases:** 1, Respiratory Apparatus; 2, Apparatus with Air Supply Pipes, (a) The Bremen Smoke Helmet, (b) The Müller Smoke Helmet, (c) The Stolz Rescue Mask; 3, Reservoir Apparatus; 4, Oxygen Apparatus. The Schwann Respiratory Apparatus. The Fleuss Respiratory Apparatus. The Improved Walcher-Gartner Pneumatophor, (a) The Single Bottle Apparatus, Instructions for Using the Pneumatophor, Taking to Pieces and Resetting the Apparatus ready for Use; (b) Two Bottle Apparatus (Shamrock Type). The Neupert Rescue Apparatus (The Mayer-Pilar System).—V., **Extinguishing Pit Fires:** (a) Chemical Means; (b) Extinction with Water. Dragging down the Burning Masses and Packing with Clay; (c) Insulating the Seat of the Fire by Dams. Dam Building. Dam Work in the Fiery Pits of Southern Hungary: (a) Cross-dams of Clay; (b) Masonry Dams, Gallery Linings. Wagner's Portable Safety Dam. Analyses of Fire Gases. Isolating the Seat of a Fire with Dams: Working in Irrespirable Gases ("Gas-diving"): 1, Air-Lock Work (Horizontal Advance) on the Mayer System as Pursued at Karwin in 1894; 2, Air-Lock Work (Horizontal Advance) by the Mauerhofer Modified System. Vertical Advance. Mayer System. Complete Isolation of the Pit. Flooding a

Burning Section isolated by means of Dams. Wooden Dams: (a) Upright Balk Dams; (b) Horizontal Balk Dams; (c) Wedge Dams, Masonry Dams. Examples of Cylindrical and Dome-shaped Dams. Dam Doors; Flooding the Whole Pit.—VI., **Rescue Stations:** (a) Stations above Ground; (b) Underground Rescue Stations.—VII., **Spontaneous Ignition of Coal in Bulk.**—Index.

Illustrations.

Sheet I., Respiratory and Rescue Appliances—Precautions against Fire. Sheet II., Respiratory and Rescue Apparatus. Sheet III., Respiratory and Rescue Apparatus Stretchers. Sheet IV., Dams. Sheet V., Signalling Appliances—Dam Construction—Cable Laying. Sheet VI., Working with Diving Gear in Irrespirable Gases—Gallery Work. Sheet VII., Working with Diving Gear in Irrespirable Gases (Mayer System)—Appliances in the Shaft.

Press Opinions.

"A work of this extremely valuable character deserves to be made widely known amongst colliery managers and mining engineers at home and abroad."—*Coal and Iron*.

"This book is, in a manner, unique. The literature of mining accidents is fairly extensive, but it consists largely of departmental Blue Books."—*Sheffield Daily Telegraph*.

"A concise and lucid description of the principal methods pursued, especially in fiery mines, and of the various appliances employed, such as respiratory and rescue apparatus, dams, etc."—*Staffs Advertiser*.

"The prevention of spontaneous combustion in collieries and the extinction of underground fires are duties that fall heavily on many colliery managers. They should, therefore, welcome this translation of Mr. Lamprecht's German treatise."—*Ironmonger*.

THE PREVENTION OF SMOKE. Combined with the Economical Combustion of Fuel. By W. C. POPPLEWELL, M.Sc., A.M.Inst., C.E., Consulting Engineer. Forty-six Illustrations. 190 pp. 1901. Demy 8vo. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Introductory.—Chapters I., Fuel and Combustion.—II., Hand Firing in Boiler Furnaces.—III., Stoking by Mechanical Means.—IV., Powdered Fuel.—V., Gaseous Fuel.—VI., Efficiency and Smoke Tests of Boilers.—VII., Some Standard Smoke Trials.—VIII., The Legal Aspect of the Smoke Question.—IX., The Best Means to be adopted for the Prevention of Smoke.—Index.

Press Opinions.

"Everybody interested in smoke prevention will derive the greatest benefit from Mr. Popplewell's treatise, and will learn much that is new to them."—*Public Health Engineer*.

"The Manchester expert who writes this book is thoroughly equipped for the task, and he has produced a work which ought to be in the hands of all Sanitary Inspectors and Health Committees, and it would be a useful present from manufacturers to stokers, instead of possibly spending the value of the volume in payment of fines."—*Sheffield Independent*.

GAS AND COAL DUST FIRING. A Critical Review of the Various Appliances Patented in Germany for this purpose since 1885. By ALBERT PÜTSCH. 130 pp. Demy 8vo. 1901. Translated from the German. With 103 Illustrations. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Generators—Generators Employing Steam—Stirring and Feed Regulating Appliances—Direct Generators—Burners—Regenerators and Recuperators—Glass Smelting Furnaces—Metallurgical Furnaces—Pottery Furnace—Coal Dust Firing.—Index.

Press Opinions.

"The work is worthy of perusal by all consumers of fuel. It is exceedingly well printed and illustrated."—*Chemical Trade Journal*.

"The book will appeal with force to the manufacturer as well as to the technical student, whilst it is also of far more than average interest to the general reader."—*Halifax Guardian*.

"The importance that gas and coal dust firing have attained of recent years, and especially the great interest attaching of late to the question of coal dust firing, makes the appearance of the present volume most opportune."—*Iron and Coal Trades Review*.

Books on Plumbing, Decorating, Metal Work, etc., etc.

EXTERNAL PLUMBING WORK. A Treatise on Lead Work for Roofs. By JOHN W. HART, R.P.C. 180 Illustrations. 270 pp. Demy 8vo. Second Edition Revised. 1902. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d. strictly net.

Contents.

Chapters I., Cast Sheet Lead.—II., Milled Sheet Lead.—III., Root Cesspools.—IV., Socket Pipes.—V., Drops.—VI., Gutters.—VII., Gutters (continued).—VIII., Breaks.—IX., Circular Bricks.—X., Flats.—XI., Flats (continued).—XII., Rolls on Flats.—XIII., Roll Ends.—XIV., Roll Intersections.—XV., Seam Rolls.—XVI., Seam Rolls (continued).—XVII., Tack Fixings.—XVIII., Step Flashings.—XIX., Step Flashings (continued).—XX., Secret Gutters.—XXI., Soakers.—XXII., Hip and Valley Soakers.—XXIII., Dormer Windows.—XXIV., Dormer Windows (continued).—XXV., Dormer Tops.—XXVI., Internal Dormers.—XXVII., Skylights.—XXVIII., Hips and Ridging.—XXIX., Hips and Ridging (continued).—XXX., Fixings for Hips and Ridging.—XXXI., Ornamental Ridging.—XXXII., Ornamental Curb Rolls.—XXXIII., Curb Rolls.—XXXIV., Cornices.—XXXV., Towers and Finials.—XXXVI., Towers and Finials (continued).—XXXVII., Towers and Finials (continued).—XXXVIII., Domes.—XXXIX., Domes (continued).—XL., Ornamental Lead Work.—XLI., Rain Water Heads.—XLII., Rain Water Heads (continued).—XLIII., Rain Water Heads (continued).

Press Opinions.

"This is an eminently practical and well-illustrated volume on the management of external lead work."—*Birmingham Daily Post*.

"It is thoroughly practical, containing many valuable hints, and cannot fail to be of great benefit to those who have not had large experience."—*Sanitary Journal*.

"Works on sanitary plumbing are by no means rare, but treatises dealing with external plumbing work are sufficiently scarce to ensure for Mr. Hart's new publication a hearty reception."—*The Ironmonger*.

HINTS TO PLUMBERS ON JOINT WIPING, PIPE BENDING AND LEAD BURNING. Third Edition, Revised and Corrected. By JOHN W. HART, R.P.C. 184 Illustrations. 313 pp. Demy 8vo. 1891. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d.; strictly net.

Contents.

Introduction. Chapters I., Pipe Bending.—II., Pipe Bending (continued).—III., Pipe Bending (continued).—IV., Square Pipe Bendings.—V., Half-circular Elbows.—VI., Curved Bends on Square Pipe.—VII., Bessed Bends.—VIII., Curved Plinth Bends.—IX., Rain-water Shoes on Square Pipe.—X., Corner and Angle Bends.—XI., Square Pipe Fixings.—XII., Joint-wiping.—XIII., Secretures for Wiped Joints.—XIV., Preparing Wiped Joints.—XV., Joint Fixings.—XVI., Plumbing Irons.—XVII., Joint Fixings.—XVIII., Use of "Touch" in Soldering.—XIX., Underland Joints.—XX., Blown and Copper Bit Joints.—XXI., Branch Joints.—XXII., Branch Joints (continued).—XXIII., Block Joints.—XXIV., Block Joints (continued).—XXV., Block Fixings.—XXVI., Astragal Joints.—XXVII., Large Branch Joints.—XXVIII., Large Underland Joints.—XXIX., Solders.—XXX., Autogenous Soldering or Lead Burning.—Index.

Press Opinions.

"Rich in useful diagrams as well as in hints."—*Liverpool Mercury*.

"The papers are eminently practical, and go much farther into the mysteries they describe than the title 'Hints' properly suggests."—*Scotsman*.

"The articles are apparently written by a thoroughly practical man. As a practical guide the book will doubtless be of much service."—*Glasgow Herald*.

"So far as the practical hints in this work are concerned, it will be useful to apprentices and students in technical schools, as it deals mainly with the most important or difficult branches of the plumber's craft, *viz.* joint wiping, pipe bending and lead burning. . . . 'Hints' are the most useful things to an apprentice, and there are many in this work which are not to be found in some of the text-books."—*English Mechanic*.

"22 PRYME STREET, HULL, 24th November, 1894.

"Gentlemen,—Your books to hand for which accept my best thanks, also for circulars. I myself got one of J. W. Hart's books on Plumbing from your traveller, and having looked through the same I can safely recommend it as being the best book I have seen. Mr. J. W. Hart treats exhaustively upon soldering and pipe bending, which are two of the most essential branches in the plumbing trade."

THE PRINCIPLES AND PRACTICE OF DIPPING, BURNISHING, LACQUERING AND BRONZING BRASS WARE. By W. NORMAN BROWN. 35 pp. Crown 8vo. 1900. Price 2s.; Abroad, 2s. 6d.; strictly net.

Contents.

Chapters I., Cleansing and Dipping; Boiling up and Cleansing; Dipping.—II., Scratch-brushing and Burnishing; Polishing; Burnishing.—III., Lacquering; Tools; Lacquers.—IV., Bronzing; Black Bronzing; Florentine Red Bronzing; Green Bronzing.—Index.

Press Opinions.

"Mr. Brown is clearly a master of his craft, and has also the immense advantage of being able to convey his instructions in a manner at once clear and concise."—*Leicester Post*.

"A thoroughly practical little treatise on the subject in all its branches, and one which should be in the hands of every tradesman or amateur who has lacquering to do."—*Irish Builder*.

WORKSHOP WRINKLES for Decorators, Painters, Paper-hangers and Others. By W. N. BROWN. Crown 8vo. 128 pp. 1901. Price 2s. 6d. ; Abroad, 3s. ; strictly net.

Contents.

Parts I., Decorating.—II., Painting.—III., Paper-hanging.—IV., Miscellaneous.
Arranged in alphabetical order.

Press Opinion.

"Decorators, painters and amateurs will find this a comprehensive work of reference on nearly every subject they are in need of."—*Building News*.

HOUSE DECORATING AND PAINTING. By W. NORMAN BROWN. Eighty-eight Illustrations. 150 pp. Crown 8vo. 1900. Price 3s. 6d. ; India and Colonies, 4s. ; Other Countries, 4s. 6d. strictly net.

Contents.

Chapters I., Tools and Appliances.—II., Colours and Their Harmony.—III., Pigments and Media.—IV., Pigments and Media.—V., Pigments and Media.—VI., Pigments and Media.—VII., Preparation of Work, etc.—VIII., Application of Ordinary Colour.—IX., Graining.—X., Graining.—XI., Graining.—XII., Gilding.—XIII., Writing and Lettering.—XIV., Sign Painting.—XV., Internal Decoration.—Index.

Press Opinion.

"The author is evidently very thoroughly at home in regard to the technical subjects he has set himself to elucidate, from the mechanical rather than the artistic point of view, although the matter of correctness of taste is by no means ignored. Mr. Brown's style is directness itself, and there is no tyro in the painting trade, however mentally ungifted, who could fail to carry away a clearer grasp of the details of the subject after going over the performance."—*Building Industries*.

A HISTORY OF DECORATIVE ART. By W. NORMAN BROWN. Thirty-nine Illustrations. 96 pp. Crown 8vo. 1900. Price 2s. 6d. ; Abroad, 3s. ; strictly net.

Contents.

Chapters I., Primitive and Prehistoric Art.—II., Egyptian Art.—III., Assyrian Art.—IV., The Art of Asia Minor.—V., Etruscan Art.—VI., Greek Art.—VII., Roman Art.—VIII., Byzantine Art.—IX., Lombard or Romanesque Art.—X., Gothic Art.—XI., Renaissance Art.—XII., The Victorian Period.—Index.

Press Opinion.

"In the course of a hundred pages with some forty illustrations Mr. Brown gives a very interesting and comprehensive survey of the progress and development of decorative art. It cannot, of course, be pretended that in the limited space named the subject is treated exhaustively and in full detail, but it is sufficiently complete to satisfy any ordinary reader; indeed, for general purposes, it is, perhaps, more acceptable than a more elaborate treatise."—*Midland Counties Herald*.

A HANDBOOK ON JAPANING AND ENAMELLING FOR CYCLES, BEDSTEADS, TINWARE, ETC. By WILLIAM NORMAN BROWN. 52 pp. and Illustrations. Crown 8vo. 1901. Price 2s. ; Abroad, 2s. 6d. ; net.

Contents.

A Few Words on Enamelling—Appliances and Apparatus—Japans or Enamels—To Test Enamel for Lead—Japaning or Enamelling Metals—Japaning Tin, such as Tea Trays, and similar Goods—Enamelling Old Work—Enamel for Cast Iron—Enamel for Copper Cooking Utensils—The Enamelling Stove—Enamelling Bedsteads, Frames and similar large pieces—Paints and Varnishes for Metallic Surfaces—Varnishes for Ironwork—Blacking for Iron—Processes for Tin Plating—Galvanising—Metal Polishes—Colours for Polished Brass—A Golden Varnish for Metal—Painting on Zinc—Carriage Varnish—Japanese Varnish and its Application.—Index

THE PRINCIPLES OF HOT WATER SUPPLY. By JOHN W. HART, R.P.C. With 129 Illustrations. 1900. 177 pp., demy 8vo. Price 7s. 6d. ; India and Colonies, 8s. ; Other Countries, 8s. 6d. ; strictly net.

Contents.

Chapters I., Water Circulation.—II., The Tank System.—III., Pipes and Joints.—IV., The Cylinder System.—V., Boilers for the Cylinder System.—VI., The Cylinder System.—VII., The Combined Tank and Cylinder System.—VIII., Combined Independent and Kitchen Boiler.—IX., Combined Cylinder and Tank System with Duplicate Boilers.—X., Indirect Heating and Boiler Explosions.—XI., Pipe Boilers.—XII., Safety Valves.—XIII., Safety Valves.—XIV., The American System.—XV., Heating Water by Steam.—XVI., Steam Kettles and Jets.—XVII., Heating Power of Steam.—XVIII., Covering for Hot Water Pipes.—Index.

Brewing and Botanical.

HOPS IN THEIR BOTANICAL, AGRICULTURAL AND TECHNICAL ASPECT, AND AS AN ARTICLE OF COMMERCE. By EMMANUEL GROSS, Professor at the Higher Agricultural College, Tetschen-Liebwerd. Translated from the German. Seventy-eight Illustrations. 1900. 340 pp. Demy 8vo. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s.; strictly net.

Contents.

PART I., HISTORY OF THE HOP.

PART II., THE HOP PLANT. Introductory.—The Roots.—The Stem and Leaves.—Inflorescence and Flower.—Inflorescence and Flower of the Male Hop; Inflorescence and Flower of the Female Hop.—The Fruit and its Glandular Structure: The Fruit and Seed.—Propagation and Selection of the Hop.—Varieties of the Hop: (a) Red Hops; (b) Green Hops; (c) Pale Green Hops.—Classification according to the Period of Ripening: 1. Early August Hops, 2. Medium Early Hops, 3. Late Hops.—Injuries to Growth: Malformations: Diseases Produced by Conditions of Soil and Climate: 1. Leaves Turning Yellow, 2. Summer or Sun-brand, 3. Cones Drooping Off, 4. Honey Dew, 5. Damage from Wind, Hail and Rain: Vegetable Enemies of the Hop: Animal Enemies of the Hop.—Beneficial Insects on Hops.

PART III., CULTIVATION. The Requirements of the Hop in Respect of Climate, Soil and Situation. Climate: Soil. Situation.—Selection of Variety and Cuttings.—Planting a Hop Garden: Drainage: Preparing the Ground: Marking-out for Planting: Planting: Cultivation and Cropping of the Hop Garden in the First Year.—Work to be Performed Annually in the Hop Garden: Working the Ground: Cutting: The Non-cutting System: The Proper Performance of the Operation of Cutting. I. Method of Cutting: Close Cutting, Ordinary Cutting, The Long Cut, The Topping Cut; II. Proper Season for Cutting: Autumn Cutting, Spring Cutting: Manuring: Training the Hop Plant: Poled Gardens, Frame Training: Principal Types of Frames: Pruning, Cropping, Topping, and Leaf Stripping the Hop Plant: Picking, Drying and Bagging.—Principal and Subsidiary Utilisation of Hops and Hop Gardens.—Life of a Hop Garden: Subsequent Cropping.—Cost of Production, Yield and Selling Prices.

PART IV.—Preservation and Storage.—Physical and Chemical Structure of the Hop Cone.—Judging the Value of Hops.

PART V.—Statistics of Production.—The Hop Trade.—Index.

Press Opinions.

"The subject is dealt with fully in every little detail; consequently, even the veriest tyro can take away some useful information from its pages."—*Irish Farming World*.

"Like an oasis in the desert comes a volume upon the above subject."—*Hereford Times*.

"This is, in our opinion, the most scholarly and exhaustive treatise on the subject of hops that has been published."—*Brewers' Journal*.

Wood Waste Utilisation.

THE UTILISATION OF WOOD WASTE. Translated from the German of ERNST HUBBARD. Crown 8vo. 1902. Fifty Illustrations. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s. net.

Contents.

Chapters I., General Remarks on the Utilisation of Sawdust.—II., Employment of Sawdust as Fuel, with and without Simultaneous Recovery of Charcoal and the Products of Distillation.—III., Manufacture of Oxalic Acid from Sawdust.—(1) Process with Soda Lye; (2) Thorn's Process; (3) Bohlig's Process.—IV., Manufacture of Spirit (Ethyl Alcohol) from Wood Waste—Patent Dyes (Organic Sulphides, Sulphur Dyes, or Mercapto Dyes).—V., Artificial Wood and Plastic Compositions from Sawdust—Production of Artificial Wood Compositions for Moulded Decorations.—VI., Employment of Sawdust for Blasting Powders and Gunpowders.—VII., Employment of Sawdust for Briquettes—Employment of Sawdust in the Ceramic Industry and as an Addition to Mortar—Manufacture of Paper Pulp from Wood—Casks.—VIII., Various Applications of Sawdust and Wood Refuse—Calcium Carbide—Manure—Wood Mosaic Plaques—Bottle Stoppers—Parquetry—Fire-lighters—Carborunum.—IX., The Production of Wood Wool—Bark.—Index

Foods and Sweetmeats.

THE MANUFACTURE OF PRESERVED FOODS AND SWEETMEATS: A Handbook of all the Processes for the Preservation of Flesh, Fruit and Vegetables, and for the Preparation of Dried Fruit, Dried Vegetables, Marmalades, Fruit-Syrups and Fermented Beverages, and of all kinds of Candies, Candied Fruit, Sweetmeats, Rocks, Drops, Dragées, Pralines, etc. By A. HAUSNER. With Twenty-eight Illustrations. Translated from the German of the third enlarged Edition. Crown 8vo. 225 pp. 1902. Price 7s. 6d.; India and Colonies, 8s.; Other Countries, 8s. 6d. net.

Contents.

Part I. **The Manufacture of Conserves.**—Chapters I., Introduction.—II., The Causes of the Putrefaction of Food.—III., The Chemical Composition of Foods.—IV., The Products of Decomposition.—V., The Causes of Fermentation and Putrefaction.—VI., Preservative Bodies.—VII., The Various Methods of Preserving Food.—VIII., The Preservation of Animal Food.—IX., Preserving Meat by Means of Ice.—X., The Preservation of Meat by Charcoal.—XI., Preservation of Meat by Drying.—XII., The Preservation of Meat by the Exclusion of Air.—XIII., The Appert Method.—XIV., Preserving Flesh by Smoking.—XV., Quick Smoking.—XVI., Preserving Meat with Salt.—XVII., Quick Salting by Air Pressure.—XVIII., Quick Salting by Liquid Pressure.—XIX., Gamgee's Method of Preserving Meat.—XX., The Preservation of Eggs.—XXI., Preservation of White and Yolk of Egg.—XXII., Milk Preservation.—XXIII., Condensed Milk.—XXIV., The Preservation of Fat.—XXV., Manufacture of Soup Tablets.—XXVI., Meat Biscuits.—XXVII., Extract of Beef.—XXVIII., The Preservation of Vegetable Foods in General.—XXIX., Compressing Vegetables.—XXX., Preservation of Vegetables by Appert's Method.—XXXI., The Preservation of Fruit.—XXXII., Preservation of Fruit by Storage.—XXXIII., The Preservation of Fruit by Drying.—XXXIV., Drying Fruit by Artificial Heat.—XXXV., Roasting Fruit.—XXXVI., The Preservation of Fruit with Sugar.—XXXVII., Boiled Preserved Fruit.—XXXVIII., The Preservation of Fruit in Spirit, Acetic Acid or Glycerine.—XXXIX., Preservation of Fruit without Boiling.—XL., Jam Manufacture.—XLI., The Manufacture of Fruit Jellies.—XLII., The Making of Gelatine Jellies.—XLIII., The Manufacture of "Sulzen."—XLIV., The Preservation of Fermented Beverages.

Part II. **The Manufacture of Candies.**—Chapters XLV., Introduction.—XLVI., The Manufacture of Candied Fruit.—XLVII., The Manufacture of Boiled Sugar and Caramel.—XLVIII., The Candying of Fruit.—XLIX., Caramelised Fruit.—L., The Manufacture of Sugar-Sticks, or Barley Sugar.—LI., Bonbon Making.—LII., Fruit Drops.—LIII., The Manufacture of Dragées.—LIV., The Machinery and Appliances used in Candy Manufacture.—LV., Dyeing Candies and Bonbons.—LVI., Essential Oils used in Candy Making.—LVII., Fruit Essences.—LVIII., The Manufacture of Filled Bonbons, Liqueur Bonbons and Stamped Lozenges.—LIX., Recipes for Jams and Jellies.—LX., Recipes for Bonbon Making.—LXI., Dragées.—Appendix.—Index.

Timber Trades.

TIMBER: A Comprehensive Study of Wood in all its Aspects (Commercial and Botanical), showing the Different Applications and Uses of Timber in Various Trades, etc. Translated from the French of PAUL CHARPENTIER, Expert Chemical Engineer, Assayer of the French Mint, etc., by JOSEPH KENNEL. Royal 8vo. 437 pp. 178 Illustrations. 1902. Price 12s. 6d.; India and Colonies, 13s. 6d.; Other Countries, 15s. net.

Contents.

Preface.—Part I., **Physical and Chemical Properties of Timber.**—Chapters I., Composition of the Vegetable Bodies—Chief Elements—M. Fremy's Researches.—II., Elementary Organs of Plants and especially of Forests.—III., Different Parts of Wood Anatomically and Chemically Considered.—IV., General Properties of Wood.

Part II., **Description of the Different Kinds of Wood.**—Chapters V., Principal Essences with Caducous Leaves.—VI., Coniferous Resinous Trees.

Part III., **Division of the Useful Varieties of Timber in the Different Countries of the Globe.**—Chapters VII., European Timber.—VIII., African Timber.—IX., Asiatic Timber.—X., American Timber.—XI., Timber of Oceania.

Part IV., **Forests.**—Chapters XII., General Notes as to Forests; their Influence.—XIII., Opinions as to Sylviculture—Improvement of Forests.—XIV., Unwooding and Rewooding.—XV., Exploitation of Forests.—XVI., Damage caused to Forests—Different Alterations.

Part V., **The Preservation of Timber.**—Chapters XVII., Generalities—Causes and Progress of Deterioration—History of Different Proposed Processes.—XVIII., Dessication—Superficial Carbonisation of Timber.—XIX., Processes by Immersion—Generalities as to

Antiseptics Employed.—XX. In éction Processes in Closed Vessels.—XXI., The Boucherie System, Based upon the Displacement of the Sap.—XXII., Processes for Making Timber Uninflammable.

Part VI., Applications of Timber. Chapters XXIII., Generalities—Working Timber—Paving Timber for Mines—Railway-Traverses.—XXIV., Accessory Products—Gums—Works of M. From—Resins—Barks—Tan—Application of Cork.—XXV., The Application of Wood to Art and Dyeing.—XXVI., Different Applications of Wood—Hard Wood—Distillation of Wood—Pyroligneous Acid—Oil of Wood—Distillation of Resins.—Index.

[See also "Wood Waste Utilisation," p. 33.]

Fancy Goods Manufacture.

THE ART OF DYEING AND STAINING MARBLE, ARTIFICIAL STONE, BONE, HORN, IVORY AND WOOD. AND OF IMITATING ALL SORTS OF WOOD. A Practical Handbook for the Use of Joiners, Turners, Manufacturers of Fancy Goods, Stick and Umbrella Makers, Comb Makers, etc. Translated from the German of D. H. SOXHLET, Technical Chemist. Crown 8vo. 168 pp. 1902. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s. net.

Contents.

Preface.—Introduction.—Chapters I., Mordants and Stains. Acids, Alkalies, Iron Salts, Copper Salts, Aluminium Salts, Chromium Salts, Tin Salts, Lead Salts, Manganese Salts, Silver and Gold Salts.—II., Natural Dyes: Redwood, Red Sandalwood, Madder, Orchil, Cudbear, Lac-Dye, Cochineal, Saffron, Annatto, Safflower, Fustic, Fustet, Quercitron, Flavin, Turmeric, Weld and its substitutes, Persian Berries, Barberry Root, Indigo, Logwood, Cutch, Galls, Sumach, Knoppert.—III., Artificial Pigments: White Lead, Naples Yellow, Red Lead, Smalts, Ultramarine, Cinnabar, Prussian Blue, Orpiment, Realgar, Chrome Green, Chrome Yellow, Chrome Red, Chrome Orange, Mosaic Gold, Green Mineral Colours, Red Ochres, Rouge, Cadmium Yellow.—IV., Coal Tar Dyes: Reds, Yellows and Oranges, Blues, Violets, Greens, Browns, Grey and Black.—Aniline Dyes Soluble in Fat: Resinate Colours, Aniline Lakes.—V., Staining Marble and Artificial Stone: Red, Violet, Blue, Green, Yellow, Orange, Brown, Black, Execution of Parti-Coloured Designs.—VI., Dyeing, Bleaching and Imitation of Bone, Horn and Ivory—Bone Bleaching—Dyeing Bone: Black, Red, Yellow, Blue, Violet, Green, Grey and Brown—Horn, Bleaching and Whitening—Dyeing Black, Grey, Brown, Blue, Green, Violet and Red—Imitation of Tortoiseshell for Combs: Yellows, Dyeing Nuts.—Ivory: Dyeing Black, Red, Yellow, Blue, Violet, Green, Grey and Brown—Further Remarks on Ivory Dyeing.—VII., Wood Dyeing: Black, Grey, Brown, Violet, Blue, Red, Yellow, Green—Imitation of Mahogany: Dark Walnut, Oak, Birch-Bark, Elder-Marquetry, Walnut, Walnut-Marquetry, Mahogany, Spanish Mahogany, Palsander and Rose Wood, Tortoiseshell, Oak, Ebony, Pear Tree—Black Dyeing Processes with Penetrating Colours.—VIII., Varnishes and Polishes: English Furniture Polish, Vienna Furniture Polish, Amber Varnish, Copal Varnish, Composition for Preserving Furniture.—Index.

Building and Architecture.

THE PREVENTION OF DAMPNESS IN BUILDINGS; with Remarks on the Causes, Nature and Effects of Saline, Efflorescences and Dry-rot, for Architects, Builders, Overseers, Plasterers, Painters and House Owners. By ADOLF WILHELM KEIM. Translated from the German of the second revised Edition by M. J. SALTER, F.I.C., F.C.S., Member of the German Chemical Society of Berlin. Eight Coloured Plates and Thirteen Illustrations. Crown 8vo. 115 pp. 1902. Price 5s.; India and Colonies, 5s. 6d.; Other Countries, 6s. net.

Contents.

Part I.—Chapters I., The Various Causes of Dampness and Decay of the Masonry of Buildings, and the Structural and Hygienic Evils of the Same.—II., Precautionary Measures during Building against Dampness and Efflorescence.—III., Methods of Remedying Dampness and Efflorescences in the Walls of Old Buildings.—IV., The Artificial Drying of New Houses, as well as Old Damp Dwellings, and the Theory of the Hardening of Mortar.—V., New, Certain and Permanently Efficient Methods for Drying Old Damp Walls and Dwellings.

Part II.—Chapters I., The Cause and Origin of Dry-rot: its Injurious Effect on Health, its Destructive Action on Buildings, and its Successful Repression.—II., Methods of Preventing Dry-rot to be Adopted During Construction.—III., Old Methods of Preventing Dry-rot.—IV., Recent and More Efficient Remedies for Dry-rot.—Index

Iron.

SIDEROLOGY: THE SCIENCE OF IRON (The Constitution of Iron Alloys and Slags). Translated from German of HANNS FREIHERR V. ZÜPTNER. 350 pp. Demy 8vo. Eleven Plates and Ten Illustrations. 1902. Price 10s. 6d.; India and Colonies, 11s.; Other Countries, 12s. net.

Contents.

Book I., **The Theory of Solution.**—Chapters I., Solutions.—II., Molten Alloys—Varieties of Solutions.—III., Osmotic Pressure.—IV., Relation between Osmotic Pressure and other Properties of Solutions.—V., Osmotic Pressure and Molecular Weight of the Dissolved Substance.—VI., Solutions of Gases.—VII., Solid Solutions.—VIII., Solubility.—IX., Diffusion.—X., Electrical Conductivity—Constitution of Electrolytes and Metals.—XI., Thermal Expansion.

Book II., **Micrography.**—Chapters I., General.—II., Microstructure.—III., The Micrographic Constituents of Iron.—IV., Relation between Micrographical Composition, Carbon-Content, and Thermal Treatment of Iron Alloys.—V., The Microstructure of Slags.

Book III., **Chemical Composition of the Alloys of Iron.**—Chapters I., Introduction.—II., Constituents of Iron Alloys—Carbon.—III., Constituents of the Iron Alloys, Carbon—Opinions and Researches on Combined Carbon.—IV., Opinions and Researches on Combined Carbon (*Continuation*).—V., Opinions and Researches on Combined Carbon (*Conclusion*).—VI., Applying the Curves of Solution deduced from the Curves of Recalescence to the Determination of the Chemical Composition of the Carbon present in Iron Alloys.—VII., The Constituents of Iron—Iron.—VIII., The Constituents of Iron Alloys—Manganese.—IX., Remaining Constituents of Iron Alloys—A Silicon.—X., Gases.

Book IV., **The Chemical Composition of Slag.**—Chapters I., Introductory.—II., Silicate Slags.—III., Calculating the Composition of Silicate Slags.—IV., Phosphate Slags.—V., Oxide Slags.—Appendix.—Index.

WORKS IN PREPARATION.

TREATISE ON CLOTH FINISHING. By ROBERT BEAUMONT, of Yorkshire College, Leeds.

WEAVING MACHINERY. Three Vols. By HARRY NISBET.

COLOUR TERMS: THEIR PROPER USE AND MEANING. By DAVID PATERSON.

USE OF WATER IN THE INDUSTRIAL ARTS.

CHEMISTRY OF DYE STUFFS. Translated from German of Dr. GEORG VON GEORGIEVICS. [*In the press.*]

SUGAR REFINERS' HANDBOOK.

ART OF BOOKBINDING.

DENTAL METALLURGY.

PRELIMINARY COTTON SPINNING AND WEAVING.

By WALTER BAILEY.

VENTILATION OF MINES. By ROBERT WABNER. Royal 8vo. Thirty Plates. About 250 pp.

SIMPLE METHODS FOR TESTING PAINTERS' MATERIALS. By A. C. WRIGHT, M.A., B.Sc. Crown 8vo. Eight Illustrations. [*In the press.*]

SEALING WAXES AND OTHER ADHESIVES. By H. C. STANDAGE.

HANDY GUIDES TO THE CHOICE OF BOOKS.

Vol. I. PROSE FICTION.

Vol. II. CLASSIFIED GUIDE TO TECHNICAL, TRADE AND COMMERCIAL LITERATURE.

Others to follow.

[*In preparation.*]

The Publishers will advise when any of the above books are ready to firms sending their addresses.

27.C.33.

Cosmetics; a handbook of the ma1902

Countway Library

BES6909



3 2044 045 984 705

27.C.33.

Cosmetics; a handbook of the ma1902

Countway Library

BES6909



3 2044 045 984 705